

A stepwise approach

IDENTIFY target behaviours AND ADDRESS possible triggers and unmet needs¹

- ▶ Pain
- ▶ Dehydration
- ▶ Boredom
- ▶ Ambient temperature too hot or cold
- ▶ Fatigue
- ▶ Lack of personal belongings
- ▶ Infections (eg, skin or urinary tract)
- ▶ Overstimulation

IMPLEMENT non-pharmacological interventions and MONITOR response^{2,3} MAINTAIN effective interventions long term

- ▶ Facilitate social contact
- ▶ Reduce excess noise, people or clutter
- ▶ Provide reminiscence therapy
- ▶ Facilitate physical activities
- ▶ Provide safe wandering areas
- ▶ Establish routines (eg, for dressing and bathing)

IF an antipsychotic is trialed to control severe symptoms (psychosis or aggression), MONITOR WEEKLY for changes in response and LOOK OUT for adverse effects^{2,4}

- ▶ Observe patients closely for adverse effects including sedation, postural hypotension, and anticholinergic effects
- ▶ Antipsychotic use is associated with serious adverse effects, including cerebrovascular events and death^{2,4}

Avoid prescribing antipsychotics for patients who have atrial fibrillation, hypertension, diabetes or have had a previous stroke²

REVIEW response to therapy weekly. Consider REDUCING dose AND STOPPING at no later than 12 WEEKS^{2,5}

Discontinue treatment if symptoms do not improve within 1-2 weeks

For more information visit: www.nps.org.au/professionals/antipsychotic-medicines

References

1. Assessment and management of behavioural and psychological symptoms of dementia. BMJ 2015;350:h369.
2. Therapeutic Guidelines: Psychotropic version 7. West Melbourne: TG Ltd, 2013.
3. Behaviour Management. A guide to good practice. Sydney: Dementia Collaborative Research Centre, 2012.
4. Australian Medicines Handbook Aged Care Companion. Adelaide: AMH Pty Ltd, 2019.
5. Clinical practice guidelines and principles of care for people with dementia. Sydney: Guideline Adaptation Committee, 2016.

nps.org.au

Level 7/418A Elizabeth Street Surry Hills NSW 2010
PO Box 1147 Strawberry Hills NSW 2012
☎ 02 8217 8700 📠 02 9211 7578 @info@nps.org.au

Independent, not-for-profit and evidence-based, NPS MedicineWise enables better decisions about medicines, medical tests and other health technologies. We receive funding from the Australian Government Department of Health. National Prescribing Service Limited ABN 61 082 034 393

© 2019 NPS MedicineWise. This information is intended for health professionals. Reasonable care is taken to ensure that this information is accurate at the date of creation. Health professionals must rely on their own expertise and inquiries taking into account the individual circumstances of each patient when providing medical advice or treatment. Where permitted by law, NPS MedicineWise disclaims all liability (including for negligence) for any loss, damage or injury resulting from reliance on or use of this information. November 2019