

# COVID-19 and the quality use of medicines: evidence, risks and fads

The COVID-19 pandemic is rapidly evolving and determining the appropriate response is complex. The severity of COVID-19 and the limited evidence for any treatment have added to the complexity of clinical decision making and prescribing. A vaccine is not yet available, but decisions about treatment are needed now.

Fear in the community has resulted in people trying unproven remedies. These fads include consuming bleach,<sup>1</sup> and gargling warm salt water or vinegar.<sup>2</sup> Each fad has varying toxicity and none is of likely benefit. Consuming chloroquine from an aquarium product<sup>3</sup> and drinking methanol have been fatal.<sup>4</sup> High-dose vitamin C has been discouraged as a treatment for COVID-19,<sup>5</sup> but reports that it is being prescribed to,<sup>6</sup> and studied in,<sup>7</sup> patients with COVID-19 could confuse the public about its place in therapy. These examples highlight the need for balanced discussions of the harms and benefits of each proposed treatment.

Australian healthcare workers have the opportunity to learn from colleagues overseas with advice being received daily. There are many anecdotes about treatment, usually conveying a brief and narrow perspective. Each can consciously and subconsciously influence our clinical decisions.

Information about the purported effects of drugs in COVID-19 is rapidly changing. Most reports focus on what is new, rather than summarising what has been learnt to date. It is easy to miss when a treatment claim becomes discredited or raises new safety concerns.

Currently, supportive care is the mainstay of treatment for COVID-19. Suggested drug treatments have been based mostly on in vitro studies or biomarkers from observational studies. At best, preliminary data from these studies should be considered hypothesis generating and prompt more research, rather than guiding clinical management. Many clinicians are not experts in research methods, so we may not appreciate the limitations of the results based on the shortcomings of the methods used in some of these studies.

Over 300 trials including more than 50 drug or biological treatments for COVID-19 have been registered.<sup>8</sup> These will generate both hope and uncertainty. Currently the main approaches include inhibiting viral replication with chloroquine, hydroxychloroquine or antiviral drugs,<sup>9</sup> immune modulation by corticosteroids, tocilizumab or stem cells, and administration of convalescent sera.

Some studies that have been popularised by the media have been uploaded to online preprint servers without the rigour of peer review. This may not be apparent from an abstract or media report so a high level of scepticism is required.

A global survey of physicians in early April 2020 found that hydroxychloroquine and azithromycin were prescribed or seen to be prescribed by nearly 50% of respondents. However, only 38% perceived efficacy in COVID-19.<sup>10</sup> Some healthcare workers have prescribed hydroxychloroquine for themselves and their families. This represents extrapolation of the very low-quality evidence for treatment<sup>11</sup> to experimental use for prophylaxis.<sup>12</sup> The decision may have been their own, but it has been rumoured that some doctors were advised by their employer to self-prescribe hydroxychloroquine due to their increased risk of being infected by patients with COVID-19. This highlights the ethical questions about prescribing experimental treatments.<sup>13</sup> It is important to distinguish off-label from experimental prescribing. Given the rate that the pandemic is evolving, the processes required to start a clinical trial may appear prolonged. However, these processes are necessary to develop a protocol, allocate resources and ensure patients are monitored to avoid treatment-related deaths. An experimental treatment should only be prescribed after informed consent is obtained.

The doses of some drugs being prescribed for COVID-19 are high compared to those used for their approved indications. Clearly, the greater the dose the greater risk of harm, and this is likely to be compounded in older patients with multiple comorbidities or those with viral myocarditis.

Some studies have used combinations of drugs which makes it difficult to assess their individual efficacy and toxicity. The combination of hydroxychloroquine and azithromycin is associated with cardiotoxicity, including a newly prolonged QTc interval of over 500 ms in 10–20% of participants.<sup>14,15</sup> A preprint publication of a retrospective study reported higher mortality in patients receiving hydroxychloroquine than those who did not.<sup>16</sup> Cardiotoxicity including ventricular tachycardia and death with higher doses of chloroquine prompted the early cessation of a Brazilian study.<sup>17</sup> Preventable drug-induced toxicity due to overdosage may have occurred in other

## Darren M Roberts

Clinical pharmacologist and Nephrologist, Departments of Clinical Pharmacology and Toxicology, and Renal Medicine and Transplantation, St Vincent's Hospital, Sydney

Conjoint associate professor, St Vincent's Clinical School, University of New South Wales, Sydney

## Alexandra Bennett

Executive officer, NSW Therapeutic Advisory Group, Sydney

## Keywords

coronavirus, COVID-19, pandemic

First published 5 May 2020

<https://doi.org/10.18773/ausprescr.2020.031>

Related article:

[Principles of ethical prescribing for self and others: hydroxychloroquine in the COVID-19 pandemic](#)

pandemics, including aspirin for influenza in 1918–19<sup>18</sup> and ribavirin for severe acute respiratory syndrome in 2003.<sup>19</sup>

In March 2020 there was much discussion that renin–angiotensin system inhibitors may increase the severity of COVID-19.<sup>20–22</sup> Much of this concern then subsided and it was recommended for patients taking these drugs to continue them.<sup>20,22</sup> The harm from stopping these drugs in patients with heart failure or other high-risk cardiovascular conditions is probably far greater than the unproven risk of severe COVID-19.<sup>23,24</sup> Subsequent studies confirmed that there was no increased risk from COVID-19 with renin–angiotensin system inhibitors,<sup>25–28</sup> confirming the earlier advice. There have also been concerns about non-steroidal anti-inflammatory drugs (NSAIDs) notably ibuprofen.<sup>21</sup> The current position is that NSAIDs can be used when indicated, but paracetamol is likely to be an acceptable alternative.<sup>29–32</sup>

Another risk from the increased prescribing of unproven drugs is that it creates a shortage of these drugs for patients who rely on them. For example, there has been a shortage of hydroxychloroquine for systemic lupus erythematosus, and reports of possible ivermectin efficacy in COVID-19 led to shortages within days. The shortages also impact on the supply

of medicines for clinical trials. Regulators, funders and policymakers have needed to enforce or introduce regulations to prevent inappropriate prescribing and stockpiling. The Therapeutic Goods Administration and Pharmaceutical Benefits Scheme have now restricted who can prescribe hydroxychloroquine.<sup>33,34</sup>

COVID-19 is presenting a number of challenges. We should not compound the crisis by inappropriate prescribing based on inadequate evidence, which increases the risk of harm and causes drug shortages. At present all prescribing for COVID-19 is experimental. Healthcare professionals must constantly analyse the literature and stay up to date using trusted resources. We need to explain clearly the challenge of balancing harm and benefit to our patients, friends and family. The COVID-19 pandemic is an opportunity to improve the health literacy of the public and to emphasise the principles of the quality use of medicines to ensure drugs are used safely and effectively. ◀

*Darren Roberts is the Chair of the Editorial Executive Committee of Australian Prescriber.*

*Darren Roberts acknowledges support of the Clinician 'Buy-Out' Program, St Vincent's Centre for Applied Medical Research.*

## REFERENCES

- Spencer SH. Fake coronavirus cures, part 1: MMS is industrial bleach. FactCheck.org 2020 Feb 11. <https://www.factcheck.org/2020/02/fake-coronavirus-cures-part-1-mms-is-industrial-bleach> [cited 2020 May 4]
- False claim: gargling with warm water and salt or vinegar will kill the coronavirus. Reuters Fact Check 2020 Mar 18. <https://www.reuters.com/article/uk-factcheck-coronavirus-salt-vinegar-idUSKBN2142MW> [cited 2020 May 4]
- Beasley D. Arizona man dies after taking chloroquine for coronavirus. Reuters 2020 Mar 24. <https://www.reuters.com/article/us-health-coronavirus-usa-chloroquine-idUSKBN21A3Y2> [cited 2020 May 4]
- Trew B. Coronavirus: hundreds dead in Iran from drinking methanol amid fake reports it cures disease. Independent 2020 Mar 27. <https://www.independent.co.uk/news/world/middle-east/iran-coronavirus-methanol-drink-cure-deaths-fake-a9429956.html> [cited 2020 May 4]
- Spencer SH. Fake coronavirus cures, part 3: vitamin C isn't a shield. FactCheck.org 2020 Feb 12. <https://www.factcheck.org/2020/02/fake-coronavirus-cures-part-3-vitamin-c-isnt-a-shield> [cited 2020 May 4]
- Mongelli L, Golding B. New York hospitals treating coronavirus patients with vitamin C. New York Post Metro 2020 Mar 24 <https://nypost.com/2020/03/24/new-york-hospitals-treating-coronavirus-patients-with-vitamin-c> [cited 2020 May 4]
- Vitamin C infusion for the treatment of severe 2019-nCoV infected pneumonia. ClinicalTrials.gov. NCT04264533 <https://clinicaltrials.gov/ct2/show/NCT04264533> [cited 2020 May 4]
- Burk-Rafel J. Updated summary of all current #COVID19 clinical trials. Incredibly, 102 trials were posted last week! [Twitter post]. 2020 Apr 6. <https://twitter.com/jbrafel/status/1247058257341972482> [cited 2020 May 4]
- Belhadi D, Peiffer-Smadja N, Lescure F-X, Yazdanpanah Y, Mentre F, Laouenan C. A brief review of antiviral drugs evaluated in registered clinical trials for COVID-19. MedRxiv [preprint] 2020 Mar 27. <https://www.medrxiv.org/content/10.1101/2020.03.18.20038190v2>
- Sermo reports hydroxychloroquine usage expands globally, plasma is perceived as more effective than hydroxychloroquine, remdesivir efficacy perceptions on the rise: data from week two of largest ongoing international market research study on COVID-19. Businesswire 2020 Apr 8. <https://www.businesswire.com/news/home/20200408005387/en/Sermo-Reports-Hydroxychloroquine-Usage-Expands-Globally-Plasma> [cited 2020 May 4]
- Hydroxychloroquine and COVID-19. Updated 2020 Apr 29. Sydney: NPS MedicineWise; 2020. <https://www.nps.org.au/hcq-and-covid-19> [cited 2020 May 4]
- Mannix L. Dentists banned from prescribing potential COVID-19 drug to self, family. Sydney Morning Herald 2020 Mar 24. <https://www.smh.com.au/national/dentists-banned-from-prescribing-potential-covid-19-drug-to-self-family-20200324-p54db2.html> [cited 2020 May 4]
- Coombes I, Markwell A, Kubler P, Redmond AM, McGurk G, Roberts JA. Principles of ethical prescribing for self and others: hydroxychloroquine in the COVID-19 pandemic. Aust Prescr 2020 Apr 22 [Epub ahead of print]. <https://doi.org/10.18773/austprescr.2020.030>
- Chorin E, Dai M, Shulman E, Wadhwani L, Bar-Cohen R, Barbhaiya C, et al. The QT interval in patients with COVID-19 treated with hydroxychloroquine and azithromycin [letter]. Nat Med 2020 Apr 24 [Epub ahead of print] <https://www.nature.com/articles/s41591-020-0888-2>
- Mercuro NJ, Yen CF, Shim DJ. Risk of QT interval prolongation associated with use of hydroxychloroquine with or without concomitant azithromycin among hospitalized patients testing positive for coronavirus disease 2019 (COVID-19). JAMA Cardiol 2020 May 1 [Epub ahead of print]. <https://jamanetwork.com/journals/jamacardiology/fullarticle/2765631>
- Magagnoli J, Narendran S, Pereira F, Cummings T, Hardin JW, Sutton SS, et al. Outcomes of hydroxychloroquine usage in United States veterans hospitalized with Covid-19. MedRxiv [preprint] 2020 Apr 23. <https://www.medrxiv.org/content/10.1101/2020.04.16.20065920v2>

17. Borba MG, Val FF, Sampaio VS, Alexandre MA, Melo GC, Brito M, et al; CloroCovid-19 Team. Effect of high vs low doses of chloroquine diphosphate as adjunctive therapy for patients hospitalized with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection: a randomized clinical trial. *JAMA Netw Open* 2020;3:e208857. <https://doi.org/10.1001/jamanetworkopen.2020.8857>
18. Starko KM. Salicylates and pandemic influenza mortality, 1918–1919 pharmacology, pathology, and historic evidence. *Clin Infect Dis* 2009;49:1405-10. <https://doi.org/10.1086/606060>
19. Muller MP, Dresser L, Raboud J, McGeer A, Rea E, Richardson SE, et al; Canadian SARS Research Network. Adverse events associated with high-dose ribavirin: evidence from the Toronto outbreak of severe acute respiratory syndrome. *Pharmacotherapy* 2007;27:494-503. <https://dx.doi.org/10.1592%2Fphco.27.4.494>
20. Vaduganathan M, Vardeny O, Michel T, McMurray JJV, Pfeffer MA, Solomon SD. Renin-angiotensin-aldosterone system inhibitors in patients with Covid-19. *N Engl J Med* 2020;282:1653-9. <https://doi.org/10.1056/nejmsr2005760>
21. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med* 2020;8:e21. [https://doi.org/10.1016/s2213-2600\(20\)30116-8](https://doi.org/10.1016/s2213-2600(20)30116-8)
22. ACE2 and hypertension. #NephJC 2020 Mar 14. <http://www.nephjc.com/news/covidace2> [cited 2020 May 4]
23. Yao X, Ye F, Zhang M, Cui C, Huang B, Niu P, et al. In vitro antiviral activity and projection of optimized dosing design of hydroxychloroquine for the treatment of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). *Clin Infect Dis* 2020 Mar 9; ciae237 [Epub ahead of print]. <https://doi.org/10.1093/cid/ciae237>
24. Zaman S, MacIsaac AI, Jennings GLR, Schlaich M, Inglis SC, Arnold R, et al. Cardiovascular disease and COVID-19: Australian/New Zealand consensus statement. *Med J Aust* 2020 April 3. <https://www.mja.com.au/journal/2020/cardiovascular-disease-and-covid-19-australiannew-zealand-consensus-statement> [cited 2020 May 4]
25. Li J, Wang X, Chen J, Zhang H, Deng A. Association of renin-angiotensin system inhibitors with severity or risk of death in patients with hypertension hospitalized for coronavirus disease 2019 (COVID-19) infection in Wuhan, China. *JAMA Cardiol* 2020 Apr 23;e201624 [Epub ahead of print]. <https://doi.org/10.1001/jamacardio.2020.1624>
26. Mehra MR, Desai SS, Kuy SR, Henry TD, Patel AN. Cardiovascular disease, drug therapy, and mortality in Covid-19. *New Engl J Med* 2020 May 1 [Epub ahead of print]. <https://doi.org/10.1056/NEJMoa2007621>
27. Mancia G, Rea F, Luderghani M, Apolone G, Corrao G. Renin-angiotensin-aldosterone system blockers and the risk of Covid-19. *New Engl J Med* 2020 May 1 [Epub ahead of print]. <https://doi.org/10.1056/NEJMoa2006923>
28. Reynolds HR, Adhikari S, Pulgarin C, Troxel AB, Iturrate E, Johnson SB, et al. Renin-angiotensin-aldosterone system inhibitors and risk of Covid-19. *N Engl J Med* 2020 May 1 [Epub ahead of print]. <https://doi.org/10.1056/NEJMoa2008975>
29. Ibuprofen and COVID-19. British Pharmacological Society News 2020 Mar 18. <https://www.bps.ac.uk/news-events/news/articles/2020/ibuprofen-and-covid-19> [cited 2020 May 4]
30. World Health Organization. Could #ibuprofen worsen disease for people with #COVID19 [Twitter post]. 2020 Mar 19. <https://twitter.com/WHO/status/1240409217997189128> [cited 2020 May 4]
31. Therapeutic Goods Administration. No evidence to support claims ibuprofen worsens COVID-19 symptoms. Canberra: TGA; 2020 Mar 20. <https://www.tga.gov.au/alert/no-evidence-support-claims-ibuprofen-worsens-covid-19-symptoms> [cited 2020 May 4]
32. European Medicines Agency. EMA gives advice on the use of non-steroidal anti-inflammatories for COVID-19 [press release]. Amsterdam: EMA; 2020 Mar 18. <https://www.ema.europa.eu/en/news/ema-gives-advice-use-non-steroidal-anti-inflammatories-covid-19> [cited 2020 May 4]
33. Therapeutic Goods Administration. Amendments to the new restrictions on prescribing hydroxychloroquine for COVID-19. Canberra: TGA; 2020 Apr 6. <https://www.tga.gov.au/alert/amendments-new-restrictions-prescribing-hydroxychloroquine-covid-19> [cited 2020 May 4]
34. Pharmaceutical Benefits Scheme. PBS News: Revised hydroxychloroquine PBS listings for the treatment of autoimmune disorders and malaria from 1 May 2020. Canberra: Australian Government Department of Health; 2020 Apr 23. <http://www.pbs.gov.au/info/news/2020/05/revised-hydroxychloroquine-pbs-listings-for-the-treatment> [cited 2020 May 4]