

Pharmacor Amisulpride

contains the active ingredient amisulpride

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about PHARMACOR AMISULPRIDE.

It does not contain all of the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have benefits and risks. Your doctor has weighed the risks of you taking PHARMACOR AMISULPRIDE against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, talk to your doctor or pharmacist.

Keep this leaflet with your medicine. You may need to read it again.

What PHARMACOR AMISULPRIDE is used for

PHARMACOR AMISULPRIDE is used to treat symptoms of schizophrenia.

Schizophrenia is a condition which affects the way you think, feel and/or act. Schizophrenia may cause symptoms such as hallucinations (eg hearing, seeing or sensing things which are not there), delusions, unusual suspiciousness, emotional and social withdrawal. People with schizophrenia may also feel depressed, anxious or tense.

PHARMACOR AMISULPRIDE belongs to a group of medicines called antipsychotics.

Your doctor may have prescribed PHARMACOR AMISULPRIDE for another reason.

Ask your doctor if you have any questions about why PHARMACOR AMISULPRIDE has been prescribed for you.

PHARMACOR AMISULPRIDE must not be taken by children up to the age of puberty. There is limited information on the use of PHARMACOR AMISULPRIDE in adolescents and its use is not recommended from puberty to the age of 18 years. If you are not yet 18 years of age, ask your doctor if PHARMACOR AMISULPRIDE is right for you.

PHARMACOR AMISULPRIDE is available only with a doctor's prescription.

Before you take PHARMACOR AMISULPRIDE

When you must not take it

Do not take PHARMACOR AMISULPRIDE if you are allergic to medicines containing amisulpride or any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include skin rash, itching or hives, swelling of the face, lips or tongue which may cause difficulty in swallowing or breathing, wheezing or shortness of breath.

Do not take PHARMACOR AMISULPRIDE if you are taking the following medicines:

- medicines used to treat irregular heart rhythm such as quinidine, disopyramide, amiodarone and sotalol
- cisapride
- antibiotics such as erythromycin and pentamidene, given as an injection into the veins
- levodopa, a medicine used in Parkinson's disease
- thioridazone, an antipsychotic
- methadone, medicine used to treat pain or addiction.

Do not take PHARMACOR AMISULPRIDE if you are breastfeeding or planning to breastfeed.

Do not take PHARMACOR AMISULPRIDE if the expiry date (Exp.) printed on the pack has passed.

Do not take PHARMACOR AMISULPRIDE if the packaging is torn or shows signs of tampering.

Do not take PHARMACOR AMISULPRIDE if you have or have had any of the following medical conditions:

- phaeochromocytoma, a rare tumour of the adrenal glands which sit near the kidneys
- tumour of the pituitary gland, a small gland at the base of the brain
- breast cancer
- liver disease.

Before you start to take it

Tell your doctor if:

- you have had an allergic reaction to any medicine which have taken previously to treat your current condition.

- any other substances, such as foods, preservatives or dyes
- you suffer from lactose intolerance because
PHARMACOR AMISULPRIDE tablets contain lactose.

Tell your doctor if you are pregnant or plan to become pregnant. PHARMACOR AMISULPRIDE is not recommended for use in pregnancy.

Your doctor will discuss the risks and benefits of taking PHARMACOR AMISULPRIDE during pregnancy. Newborns and mothers who have taken PHARMACOR AMISULPRIDE during pregnancy need to be carefully monitored.

Tell your doctor if you are breastfeeding or wish to breastfeed.

Your doctor will discuss the risks and benefits of taking PHARMACOR AMISULPRIDE when breastfeeding.

Tell your doctor if you have any medical conditions, especially the following:

- kidney or liver disease, Parkinson's disease or seizures
- problems with the heart and blood vessels
- a history of blood clots
- hyperglycaemia (high sugar levels in the blood) or a family history of diabetes. Your doctor may recommend monitoring your blood sugar levels while you are taking PHARMACOR AMISULPRIDE.
- suffer from dementia
- mental/mood changes or suicidal thoughts. Patients (and care givers of patients) need to monitor for any worsening of their condition and/or the development of thoughts of suicide, suicidal behaviour or thoughts of harming themselves. Seek medical advice immediately if these symptoms present

- risk factors for stroke
- history or family history of breast cancer.

Your doctor may want to take special care if you have any of these conditions.

If you have not told your doctor about any of the above, tell them before you start taking PHARMACOR AMISULPRIDE.

Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Some medicines may be affected by PHARMACOR AMISULPRIDE, or may affect how well it works. These include:

- medicines used to treat irregular heart rhythm such as quinidine, disopyramide, amiodarone and sotalol
- other medicines used to treat heart problems such as diltiazem, verapamil, clonidine, digoxin and drugs known as beta blockers (e.g. propranolol)
- intravenous amphotericin B, an anti-fungal given by injection into the veins
- other antipsychotics such as thioridazine, chlorpromazine, trifluperazine, pimozide, haloperidol, imipramine and lithium
- diuretics
- stimulant laxatives
- glucocorticosteroids
- diagnostics drug such as tetracosactides
- medicines taken for anxiety or to help you sleep
- medicines taken for depression
- some strong pain killers
- antihistamines, medicines to treat allergies, which cause drowsiness
- some medicines taken to control blood pressure.

Your doctor can tell you what to do if you are taking any of these medicines.

If you are not sure whether you are taking any of these medicines, check with your doctor or pharmacist.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking PHARMACOR AMISULPRIDE.

How to take PHARMACOR AMISULPRIDE

How much to take

Your doctor will tell you how many tablets you need to take each day and when to take them. This depends on your condition and whether or not you are taking any other medicines. The dosage is adjusted for each individual and can range from 50 mg a day up to 800 mg a day. In some cases your doctor may increase the dose to 1200 mg a day.

Follow all directions given to you by your doctor and pharmacist carefully.

Do not take more than the dose your doctor has recommended.

How to take PHARMACOR AMISULPRIDE

Swallow the tablets whole with a glass of water.

If you forget to take PHARMACOR AMISULPRIDE

If you forget to take your medicine, take your dose as soon as you remember.

Do not take a double dose to make up for the dose you missed.

If you are not sure what to do, ask your doctor or pharmacist.

How long to take PHARMACOR AMISULPRIDE for

It is important that you do NOT stop taking PHARMACOR AMISULPRIDE unless your doctor tells you. Do not stop taking PHARMACOR AMISULPRIDE just because you feel better. It is very important to continue PHARMACOR AMISULPRIDE because it will help you stay well.

Keep taking PHARMACOR AMISULPRIDE for as long as your doctor recommends.

If you take too much PHARMACOR AMISULPRIDE (overdose)

Immediately telephone your doctor, or the Poisons Information Centre (telephone 13 11 26), or go to Accident and Emergency at the nearest hospital, if you think you or anyone else may have taken too much PHARMACOR AMISULPRIDE. Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

If you take too much PHARMACOR AMISULPRIDE, you may feel drowsy and have slurred speech.

While you are taking PHARMACOR AMISULPRIDE

Things you must do

It is very important to continue taking PHARMACOR AMISULPRIDE because it will help you stay well.

Before starting any new medicine, tell your doctor or pharmacist that you are taking PHARMACOR AMISULPRIDE.

Tell all the doctors, dentists and pharmacists who are treating you

**that you are taking
PHARMACOR AMISULPRIDE.**

Tell your doctor immediately, or go to the nearest hospital, if you have any of the following suicidal thoughts or mental/mood changes:

- thoughts or talk of death or suicide
- thoughts or talk of self-harm or harm to others
- any recent attempts of self-harm
- increase in aggressive behaviour, irritability or agitation
- depressed mood or worsening of depression.

Occasionally, the symptoms of depression may include thoughts of suicide or self-harm. These symptoms may continue to get worse during the early stages of treatment until effect of the medicine becomes apparent. All mentions of suicide or violence must be taken seriously.

Visit your doctor regularly so they can check on your progress.

Things you must not do

Do not use PHARMACOR AMISULPRIDE to treat any other conditions unless your doctor tells you to.

Do not give PHARMACOR AMISULPRIDE to anyone else, even if they have the same condition as you.

Things to be careful of

Be careful driving or operating machinery until you know how PHARMACOR AMISULPRIDE affects you.

PHARMACOR AMISULPRIDE may cause drowsiness, dizziness or lightheadedness in some people. If any of these occur, do not drive, operate machinery or do anything else that could be dangerous.

Be careful if you are elderly or unwell. Some people may experience side effects such as drowsiness, confusion, dizziness and unsteadiness, which may increase the risk of a fall.

Be careful when drinking alcohol while taking PHARMACOR AMISULPRIDE the effects may be worse. It is NOT recommended that you drink alcohol while taking PHARMACOR AMISULPRIDE.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking PHARMACOR AMISULPRIDE.

Like all other medicines, AMISULPRIDE may have unwanted side effects in some people. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects. Some side effects are dose related, so it is important that you never exceed your prescribed dose.

Do not be alarmed by this list of possible side effects.

You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor if you notice any of the following and they worry you:

- drowsiness
- weight gain
- dizziness
- increased appetite
- nausea
- vomiting
- constipation
- dry mouth
- insomnia
- anxiety
- agitation
- problems with orgasm.

These are the most common side effects of PHARMACOR AMISULPRIDE.

Some people may feel dizzy in the early stages of treatment, especially when getting up from a lying or sitting position. This side effect

usually passes after taking PHARMACOR AMISULPRIDE for a few days.

Sometimes trembling, noticeable muscle stiffness or spasm, slowness of movement, excess saliva, restlessness, an overwhelming urge to move and either distress or movements such as pacing, swinging of the legs while seated, rocking from foot to foot, or both can occur. This will usually be reduced if your dose of PHARMACOR AMISULPRIDE is lowered by your doctor or if your doctor prescribes you an additional medicine.

High blood sugar has been reported in patients taking PHARMACOR AMISULPRIDE. Symptoms of high sugar levels in the blood include passing more urine than normal, persistent excessive thirst, increased appetite with a loss in weight and weakness.

Tell your doctor immediately or go to Accident and Emergency at the nearest hospital if you notice any of the following:

- muscle twitching
- abnormal movements mainly of the face or tongue
- fever
- unexplained infections
- faster breathing
- sweating
- muscle stiffness

If this occurs, stop taking PHARMACOR AMISULPRIDE immediately and contact your doctor.

After prolonged use in women, medicines of this type can cause:

- breast pain
- milk secretion
- an absence of their monthly period
- changes in the regularity of their periods

Tell your doctor if your monthly periods are absent for six months or more.

After prolonged use in men, medicines of this type can cause breast enlargement or impotence.

Incidences of abnormal liver function have been occasionally reported.

Other side effects not listed above may also occur in some patients. Tell your doctor if you notice anything that is making you feel unwell.

After using Pharmacor Amisulpride

Storage

Keep PHARMACOR AMISULPRIDE where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Keep your tablets in the blister pack until it is time to take them.

If you take the tablets out of the blister pack they may not keep well.

Keep your tablets in a cool dry place where the temperature stays below 30°C.

Do not store PHARMACOR AMISULPRIDE or any other medicine in the bathroom or near a sink.

Do not leave PHARMACOR AMISULPRIDE in the car or on window sills.

Heat and dampness can destroy some medicines.

Disposal

If your doctor tells you to stop taking PHARMACOR AMISULPRIDE, or your tablets have passed their expiry date, ask your pharmacist what to do with any that are left over.

Product description

What it looks like

PHARMACOR AMISULPRIDE comes in 3 strengths of tablets:

- PHARMACOR AMISULPRIDE 100 - White, round, flat tablets embossed MC on one side
- PHARMACOR AMISULPRIDE 200 - White, round, flat scored on one side tablets
- PHARMACOR AMISULPRIDE 400 - White, biconvex, capsule shaped tablets, scored on both sides.

The 100 mg, 200 mg and 400 mg strengths are available in pack sizes of 30, 60 and 90 tablets.

Ingredients

The active ingredient in PHARMACOR AMISULPRIDE is amisulpride.

- each PHARMACOR AMISULPRIDE 100 tablet contains 100 mg of amisulpride
- each PHARMACOR AMISULPRIDE 200 tablet contains 200 mg of amisulpride.
- each PHARMACOR AMISULPRIDE 400 tablet contains 400 mg of amisulpride.

The tablets also contain:

- microcrystalline cellulose
- hypromellose
- magnesium stearate
- lactose
- sodium starch glycollate A.

The tablets do not contain gluten, sucrose, tartrazine or any other azo dyes.

Supplier

Southern Cross Pharma Pty Ltd
Suite 5/118 Church Street
Hawthorn VIC 3122
Australia

Australian registration numbers:
PHARMACOR AMISULPRIDE
100 AUST R 234695

PHARMACOR AMISULPRIDE

200 AUST R 234693

PHARMACOR AMISULPRIDE

400 AUST R 234694

Date of preparation:

May 2017