

AZITHROMYCIN-AFT IV

Azithromycin (a-zithro-my-sin)

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Azithromycin-AFT. It does not contain all the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you being given Azithromycin-AFT against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with you.

You may need to read it again.

What Azithromycin-AFT is used for

Azithromycin-AFT is used to treat pneumonia, a lung infection caused by certain bacteria including *Legionella pneumophila*.

Azithromycin-AFT is an antibiotic which belongs to a group of medicines called azalides.

The azalides are a sub-class of a group of antibiotics called macrolides.

Azithromycin-AFT works by killing or stopping the growth of bacteria causing your infection.

Azithromycin-AFT will not work against viral infections such as colds or flu.

Your doctor may have prescribed Azithromycin-AFT for another reason.

Ask your doctor if you have any questions about why Azithromycin-AFT has been prescribed for you.

This medicine is available only with a doctor's prescription.

This medicine is not addictive.

This medicine is not expected to affect your ability to drive a car or operate machinery.

Before you are given Azithromycin-AFT

When you must not be given it

You must not be given Azithromycin-AFT if you have an allergy to:

Azithromycin or any other ketolide or macrolide antibiotic (e.g., roxithromycin, erythromycin, clarithromycin, telithromycin) or any of the ingredients listed at the end of this leaflet

If you are not sure if you are allergic to any of the above, ask your doctor.

Symptoms of an allergic reaction may include shortness of breath, wheezing or difficulty breathing; swelling of the face, lips, tongue or to other parts of the body; skin rash, itching or hives.

Azithromycin-AFT must not be given if the expiry date (EXP) or use by date printed on the pack has passed. If you take this medicine after this date, it may not work.

Azithromycin-AFT must not be given if the packaging is torn or shows signs of tampering.

If you are not sure whether you should be given Azithromycin-AFT, talk to your doctor before you are given it.

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you have or have had any of the following medical conditions:

- any kidney problems
- any liver problems
- any heart problems, including abnormalities of the rhythm
- muscle weakness
- low levels of potassium or magnesium in your blood
- if you are pregnant or if you plan to become pregnant or are breastfeeding. Your doctor can discuss with you the risks and benefits involved.

If you have not told your doctor about any of the above, tell him/her before you are given Azithromycin-AFT.

Taking other medicines

Tell your doctor if you are taking any other medicines, including medicines you buy without a prescription from a pharmacy, supermarket or health food shop.

Some medicines may interfere with Azithromycin-AFT or increase the risk of side effects. These include:

- antacids (a medicine used to treat indigestion)
- coumarin-type oral anticoagulants (a medicine used to prevent blood clots)
- cyclosporin (a medicine used to prevent organ transplant rejection or to treat certain problems with the immune system)
- digoxin (a medicine used to treat abnormal heart rhythm or heart failure)
- ergot derivatives (such as ergotamine, which is used to treat migraines)
- terfenadine or astemizole (medicines used to treat allergies and hayfever)
- zidovudine (a medicine used to treat patients with AIDS)
- diphenoxylate (Lomotil), a medicine used to treat diarrhoea
- some medicines used to treat heart rhythm problems (heart arrhythmia) such as amiodarone, disopyramide, ibutilide and sotalol
- antipsychotic medicines used to treat schizophrenia or bipolar mania such as haloperidol, quetiapine and risperidone
- medicines used to treat depression (antidepressants) such as fluoxetine, sertraline and venlafaxine
- fluoroquinolone antibiotics such as ciprofloxacin, lomefloxacin, moxifloxacin and norfloxacin.

These medicines may be affected by Azithromycin-AFT or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines.

Your doctor or pharmacist has more information on medicines to be careful with or avoid while being given Azithromycin-AFT.

Talk to your doctor about the need for additional contraception while being given Azithromycin-AFT.

Some antibiotics may decrease the effectiveness of some birth control pills, although this has not been shown with Azithromycin-AFT.

How Azithromycin-AFT is given

Follow all directions given to you by your doctor or pharmacist carefully.

They may differ from the information contained in this leaflet.

How much is given

Your doctor will decide how much Azithromycin-AFT you should receive and the length of time for which you should receive it.

The usual dose of Azithromycin-AFT is 500 mg given as an injection into a vein for 2 to 5 days. You may then be given 500 mg oral Azithromycin-AFT once a day for another 2 - 7 days (to complete a 7 to 10 day course of antibiotics).

How it is given

Azithromycin-AFT is given as an injection by a doctor or trained nurse.

Azithromycin-AFT is a powder which is mixed with Water for Injections and then diluted by the pharmacist. It is then injected into a vein which takes approximately 60 minutes.

If too much is given (Overdose)

As Azithromycin-AFT is given under the close supervision of your doctor, it is very unlikely that you will receive too much. If you experience any side effects tell your doctor immediately.

While you are being treated

Things you must do

If the symptoms of your infection do not improve within a few days, or if they become worse, tell your doctor.

If you get severe diarrhoea, tell your doctor, pharmacist or nurse immediately. Do this even if it occurs several weeks after Azithromycin-AFT has been stopped.

Diarrhoea may mean that you have a serious condition affecting your bowel. You may need urgent medical care. Do not take any diarrhoea medicine without first checking with your doctor.

If you get a sore, white mouth or tongue while being given, or soon after stopping Azithromycin-AFT, tell your doctor. Also tell your doctor if you get vaginal itching or discharge.

This may mean you have a yeast infection called thrush. Sometimes the use of Azithromycin-AFT allows yeast to grow and the above symptoms to occur. Azithromycin-AFT does not work against yeast.

If you become pregnant while being given Azithromycin-AFT, tell your doctor.

Tell your doctor immediately if during treatment with Azithromycin-AFT your baby develops irritability with feeding or starts vomiting.

This may be a sign of a stomach disorder in the infant.

If you are about to start any new medicines, tell your doctor and pharmacist that you are being given Azithromycin-AFT.

Tell all doctors, dentists and pharmacists who are treating you that you are being given Azithromycin-AFT.

Things to be careful of

Protect your skin when you are in the sun, especially between 10am and 3pm.

Some macrolide antibiotics may cause your skin to be more sensitive to sunlight than it is normally. Exposure to sunlight may cause a skin rash, itching, redness or severe sunburn.

If outdoors, wear protective clothing and use a 30+ sunscreen. If your skin does appear to be burning tell your doctor immediately.

Things you must not do

Do not stop taking Azithromycin-AFT or lower the dosage without checking with your doctor.

If you do not complete the full course prescribed by your doctor, all the organisms causing your infection may not be killed. These organisms may continue to grow and multiply so that your infection may not clear completely or may return.

Do not give Azithromycin-AFT to anyone else, even if they have the same condition as you.

Do not use Azithromycin-AFT to treat any other medical complaints unless your doctor tells you to.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Azithromycin-AFT.

Like other medicines, Azithromycin-AFT can cause some side effects. If they occur, most are likely to be minor and temporary. However, some may be serious and need medical attention.

Ask your doctor to answer any questions you may have.

Do not be alarmed by the following list of side effects. You may not experience any of them.

While taking it

Tell your doctor if you notice any of the following and they worry you:

- pain or inflammation at the site of infusion and pain during infusion
- oral thrush - white, furry, sore tongue and mouth
- vaginal thrush - sore and itchy vagina and/or white discharge
- nausea (feeling sick), loss of appetite, vomiting, stomach pain, indigestion, wind, constipation, loose bowel motions
- dizziness, headache, spinning sensation
- tiredness, drowsiness, fatigue
- muscle or joint aches
- rash
- hearing loss or ringing in the ears
- altered taste and smell.

These side effects are usually mild.

Tell your doctor immediately if you notice any of the following:

- severe persistent diarrhoea
- fast or irregular heart beat
- symptoms of sunburn such as redness, itching, swelling or blistering which may occur more quickly than normal
- decreased feeling or sensitivity, especially in the skin
- hives, itching or skin rash
- widespread body rash, fever and swollen lymph nodes
- signs of frequent or worrying infections such as fever, severe chills, sore throat or mouth ulcers
- bleeding or bruising more easily than normal, reddish or purplish blotches under the skin
- dark urine or blood in the urine or bowel motions
- aggressive reaction, nervousness, agitation or anxiety
- severe upper stomach pain, often with nausea and vomiting.

These are serious side effects. You may need urgent medical attention. Serious side effects are rare.

If any of the following happen, stop taking Azithromycin-AFT and tell your doctor immediately or go to casualty at your nearest hospital:

- sudden signs of allergy such as rash, itching or hives on the skin, swelling of the face, lips, tongue or other parts of the body, shortness of breath, wheezing or trouble breathing
- blisters or ulcers on the skin, in the mouth or airways that may occur after a period of fever
- diarrhoea, usually with blood and mucus, stomach pain and fever
- yellowing of the eyes or skin, also called jaundice
- chest pain
- fainting
- convulsions (fits).

These are very serious side effects. You may need urgent medical attention or hospitalisation. These side effects are rare.

After finishing it

Tell your doctor immediately if you notice any of the following side effects, particularly if they occur several weeks after stopping treatment with Azithromycin-AFT:

- severe stomach cramps
- watery and severe diarrhoea, which may be bloody
- fever, in combination with one or both of the above.

Azithromycin-AFT can cause some bacteria, which are normally present in the bowel and normally harmless, to multiply and therefore cause the above symptoms. You may need urgent medical attention. However, this side effect is rare.

Do not take any medicine for this diarrhoea without first checking with your doctor.

Tell your doctor if you notice anything else that is making you feel unwell.

Other side effects not listed above may also occur in some patients.

Some of these side effects (for example certain liver conditions, and blood abnormalities) can only be found when your doctor does tests from time to time to check your progress.

Do not be alarmed at this list of possible side effects.

You may not experience any of them.

Storage using Azithromycin-AFT

Storage

Azithromycin-AFT is stored in the pharmacy or on the ward. It is kept in a cool dry place where the temperature stays below 30°C.

Disposal

The hospital staff will dispose of any leftover Azithromycin-AFT.

Product description

What it looks like

A 10 mL vial of Azithromycin-AFT contains 500 mg of azithromycin, providing 100 mg/mL solution following reconstitution.

Active Ingredient

The active ingredient is azithromycin.

Other ingredients

Sodium hydroxide

Citric acid

Supplier

Azithromycin-AFT is supplied by:

AFT Pharmaceuticals Pty Ltd
113 Wicks Road
North Ryde

Toll Free Number: 180023874276

Australian Registration Numbers

AUST R 299585

This leaflet was prepared in November 2017