

EZETIMIBE – ROSUVASTATIN SANDOZ®

ezetimibe and rosuvastatin calcium

Consumer Medicine Information

WHAT IS IN THIS LEAFLET

This leaflet answers some common questions about EZETIMIBE - ROSUVASTATIN SANDOZ.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking EZETIMIBE - ROSUVASTATIN SANDOZ against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

WHAT EZETIMIBE - ROSUVASTATIN SANDOZ IS USED FOR

EZETIMIBE - ROSUVASTATIN SANDOZ helps to lower cholesterol levels. It is used in people whose cholesterol levels are too high and when diet alone cannot lower these levels adequately.

What is high cholesterol

Cholesterol is one of several fatty substances found in the bloodstream. Your total cholesterol is made up mainly of LDL and HDL cholesterol.

LDL cholesterol is often called "bad" cholesterol because it can build up in the walls of your arteries forming plaque. Eventually this plaque buildup can lead to a narrowing of the arteries. This narrowing can slow or block blood flow to vital organs such as the heart and brain. This blocking of blood flow can result in a heart attack or stroke.

HDL cholesterol is often called "good" cholesterol because it helps keep the bad cholesterol from building up in the arteries and protects against heart disease.

How EZETIMIBE - ROSUVASTATIN SANDOZ works

EZETIMIBE - ROSUVASTATIN SANDOZ is a combination pack, which contains two different medicines. One is Ezetimibe and the other is rosuvastatin (as rosuvastatin calcium). EZETIMIBE - ROSUVASTATIN SANDOZ reduces elevated total-cholesterol, LDL ("bad") cholesterol and increases HDL ("good") cholesterol.

EZETIMIBE - ROSUVASTATIN SANDOZ works by decreasing the absorption of cholesterol in the small intestine and by reducing the amount of cholesterol made in the liver.

Your doctor may have prescribed EZETIMIBE - ROSUVASTATIN SANDOZ for another reason.

Ask your doctor if you have any questions about why EZETIMIBE - ROSUVASTATIN SANDOZ has been prescribed for you.

EZETIMIBE - ROSUVASTATIN SANDOZ is not addictive.

EZETIMIBE - ROSUVASTATIN SANDOZ is available only with a doctor's prescription.

Use in children

EZETIMIBE - ROSUVASTATIN SANDOZ is not recommended for use in children.

BEFORE YOU TAKE EZETIMIBE - ROSUVASTATIN SANDOZ

When you must not take it

Do not take EZETIMIBE - ROSUVASTATIN SANDOZ if:

- **you have an allergy to EZETIMIBE - ROSUVASTATIN SANDOZ or any of the ingredients listed at the end of this leaflet.**
Symptoms of an allergic reaction may include:
 - shortness of breath
 - wheezing or difficulty breathing
 - swelling of the face, lips, tongue, throat or other parts of the body
 - rash, itching or hives on the skin
- **you have had muscle pain, tenderness or weakness from other medicines used to treat high cholesterol or triglycerides.**
- **you are pregnant or intend to become pregnant.**
If you are a woman of childbearing age and are taking EZETIMIBE - ROSUVASTATIN SANDOZ, use a proven method of birth control to avoid pregnancy.
The medicine may affect your developing baby if you take it during pregnancy.
- **you are breast-feeding.**
This medicine may pass into breast milk and affect your baby.
- **you have active liver disease.**
- **you are taking any medicine containing fusidic acid (for bacterial infections)**
- **the expiry date on the pack has passed or if the packaging is torn or shows signs of tampering.**
If you take EZETIMIBE - ROSUVASTATIN SANDOZ after the expiry date has passed, it may not work.
Return any expired or damaged EZETIMIBE - ROSUVASTATIN SANDOZ to your pharmacist for disposal.

Do not take EZETIMIBE - ROSUVASTATIN SANDOZ together with fenofibrate if:

- **you have gall bladder disease.**

Do not use EZETIMIBE - ROSUVASTATIN SANDOZ 10mg+40mg if you:

- have low thyroid hormone levels (hypothyroidism)
- have a personal or family history of hereditary muscular disorders
- have a previous history of muscular problems from using other medicines used to treat high cholesterol or triglycerides
- have a history of very heavy alcohol consumption
- have severe kidney impairment
- are of Asian descent

- are taking another type of lipid lowering medicine called a fibrate

Before taking EZETIMIBE -

ROSUVASTATIN SANDOZ 10mg+40mg, talk to your doctor about situations that can increase rosuvastatin blood levels.

Also talk to your doctor if you are not sure whether you should start taking EZETIMIBE - ROSUVASTATIN SANDOZ.

Before you start to take it

Your doctor will ask you to have your liver function tested before you start to take EZETIMIBE - ROSUVASTATIN SANDOZ.

Tell your doctor if:

- **you have any allergies to any other medicines or any other substances, such as foods, preservatives or dyes.**
- **you have, or have had, any medical condition, including:**
 - liver problems
Your doctor will do a blood test to make sure you have no problems with your liver.
 - kidney problems
 - personal or family history of muscle problems
 - muscle pain, tenderness or weakness from other medicines used to treat high cholesterol or triglycerides
 - unexplained muscle aches or pains in your muscle
 - breathing problems
 - low thyroid hormone levels (hypothyroidism)
 - diabetes or risk factors for diabetes
- **you drink alcohol regularly. Excessive alcohol consumption may not be safe in patients taking EZETIMIBE - ROSUVASTATIN SANDOZ.**
- **you have or have had myasthenia gravis (a disease causing general muscle weakness including in some cases muscles used for breathing) or ocular myasthenia (a disease causing eye muscle weakness) as statins may lead to occurrence of myasthenia or aggravate the condition.**

If you have not told your doctor about any of the above, tell him/ her before you start taking EZETIMIBE - ROSUVASTATIN SANDOZ.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines should not be taken with EZETIMIBE - ROSUVASTATIN SANDOZ, including:

- **fusidic acid, antibiotic used to treat infection**
- **fenofibrate (a medicine to lower triglyceride levels) if you have gall bladder disease**

Some medicines may be affected by EZETIMIBE - ROSUVASTATIN SANDOZ, or may affect how well it works, or may increase the risk of side effects with EZETIMIBE - ROSUVASTATIN

SANDOZ. You may need different amounts of your medicines, or you may need to take different medicines or take your medicines at different times. Your doctor will advise you.

Tell your doctor if...

Tell your doctor if you are taking any of the following:

- bile acid sequestrants, such as cholestyramine, used to lower cholesterol levels
- other medicines to lower cholesterol or triglyceride levels, for example, gemfibrozil, other fibrates, Vitamin B3
- warfarin, fluidione or ticagrelor, used to prevent blood clots
- HIV protease inhibitors, used to treat HIV infection
- hepatitis C antiviral agents, such as elbasvir or grazoprevir, used to treat the hepatitis C virus infection
- colchicine, used to treat gout
- ciclosporin, used to suppress the immune system
- antacids, used to treat reflux or ulcers. EZETIMIBE - ROSUVASTATIN SANDOZ should be taken 2 hours before or 2 hours after taking antacid
- daptomycin, a medicine used to treat complicated skin and skin structure infections and bacteraemia.

Your doctor or pharmacist has more information on medicines to be careful with or avoid while taking EZETIMIBE - ROSUVASTATIN SANDOZ.

If you have not told your doctor about any of the above, tell them before you start taking EZETIMIBE - ROSUVASTATIN SANDOZ.

HOW TO TAKE EZETIMIBE - ROSUVASTATIN SANDOZ

Take EZETIMIBE - ROSUVASTATIN SANDOZ only when prescribed by your doctor.

Follow all directions given to you by your doctor or pharmacist carefully.

They may differ from the information contained in this leaflet.

If you do not understand these instructions, ask your doctor or pharmacist for help.

If you are currently taking a medicine that contains ezetimibe and/or rosuvastatin which are contained in EZETIMIBE - ROSUVASTATIN SANDOZ:

- stop taking your current medicine(s) that contain ezetimibe and/or rosuvastatin as this may result in you taking more medicine than you need
- take the remaining medicine(s) to your pharmacist for safe disposal

Check with your doctor if you are not sure about the medicines you are taking.

How much to take it

There are two different medicines in EZETIMIBE - ROSUVASTATIN SANDOZ. Your doctor will prescribe a strength that is appropriate for you. It is very important that you take EZETIMIBE - ROSUVASTATIN SANDOZ exactly as follows:

Take one Ezetimibe Sandoz tablet and one Rosuvastatin Sandoz (rosuvastatin calcium) tablet at the same time once a day.

If your doctor has increased the strength of EZETIMIBE - ROSUVASTATIN SANDOZ tablets you take, do not take multiple doses of any remaining supplies of lower strength EZETIMIBE - ROSUVASTATIN SANDOZ tablets instead of the new higher strength tablets as this will result in you taking more medicine (ezetimibe) than you need.

How to take it

Swallow the tablets whole with a full glass of water.

Do not chew or crush the tablets.

When to take it

EZETIMIBE - ROSUVASTATIN SANDOZ can be taken at any time of the day. However, EZETIMIBE - ROSUVASTATIN SANDOZ should be taken at about the same time each day. Taking it at the same time each day will have the best effect. It will also help you remember when to take it.

Take EZETIMIBE - ROSUVASTATIN SANDOZ with or without food.

Taking EZETIMIBE - ROSUVASTATIN SANDOZ with other cholesterol-lowering agents

Your doctor may ask you to take EZETIMIBE - ROSUVASTATIN SANDOZ with other cholesterol-lowering agents such as bile acid sequestrants.

If you are taking a bile acid sequestrant, such as cholestyramine, take EZETIMIBE - ROSUVASTATIN SANDOZ either at least two hours before or four hours after taking the bile acid sequestrant.

How long to take Rosuvastatin Sandoz

Take EZETIMIBE - ROSUVASTATIN SANDOZ every day and continue taking it for as long as your doctor tells you. EZETIMIBE - ROSUVASTATIN SANDOZ helps to lower your cholesterol levels but does not cure your condition. If you stop taking EZETIMIBE - ROSUVASTATIN SANDOZ your cholesterol levels may rise again.

If you forget to take it

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to. Otherwise, take it as soon as you remember, and then go back to taking your tablets as you would normally.

Do not take a double dose to make up for the dose that you missed.

This may increase the chance of you getting a side effect.

If you are not sure what to do, talk to your doctor or pharmacist.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

If you take too much (overdose)

Immediately telephone your doctor or the Poisons Information Centre (telephone Australia 13 11 26 or New Zealand 0800 POISON or 0800 764766) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much EZETIMIBE - ROSUVASTATIN

SANDOZ. Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

WHILE YOU ARE TAKING EZETIMIBE - ROSUVASTATIN SANDOZ

Things you must do

If you become pregnant while you are taking EZETIMIBE - ROSUVASTATIN SANDOZ, stop taking it and tell your doctor immediately.

Keep all your doctor's appointments.

Even if you are taking medicines to treat high cholesterol, it is important to have your cholesterol measured regularly. You should also know your cholesterol levels and goals.

Your doctor will ask you to have your liver function tested from time to time while you are taking EZETIMIBE - ROSUVASTATIN SANDOZ. This may help prevent unwanted side effects.

Your doctor may adjust your dose depending on the result of the test.

If you are about to be started on any new medicine, tell your doctor and pharmacist that you are taking EZETIMIBE - ROSUVASTATIN SANDOZ.

Tell any other doctors, dentists, and pharmacists who treat you that you are taking EZETIMIBE - ROSUVASTATIN SANDOZ.

Things you must not do

Do not take EZETIMIBE - ROSUVASTATIN SANDOZ to treat any other conditions unless your doctor tells you to.

Do not give EZETIMIBE - ROSUVASTATIN SANDOZ to anyone else, even if they have the same condition as you.

Things to be careful of

Avoid drinking large quantities of alcohol.

Drinking large quantities of alcohol while taking EZETIMIBE - ROSUVASTATIN SANDOZ may increase your chance of getting liver problems.

Be careful driving or operating machinery until you know how EZETIMIBE - ROSUVASTATIN SANDOZ affects you.

There have been side effects reported with EZETIMIBE - ROSUVASTATIN SANDOZ that may affect your ability to drive or operate machinery. Individual responses to EZETIMIBE - ROSUVASTATIN SANDOZ may vary.

Things that may help your condition

Lifestyle Changes -

This includes a cholesterol-lowering diet, increasing physical activity, and weight management. Ask your doctor for advice before increasing physical activity.

Medicines -

Cholesterol-lowering medicines are used together with lifestyle changes to help lower cholesterol.

SIDE EFFECTS

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking EZETIMIBE - ROSUVASTATIN SANDOZ.

EZETIMIBE - ROSUVASTATIN SANDOZ helps most people with high cholesterol levels, but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Do not be alarmed by the following lists of side effects.

You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor if...

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- diarrhoea
- muscle aches, spasms, tiredness or weakness
- constipation
- nausea
- stomach or belly pain
- heartburn, indigestion or wind
- aches and pain
- pain in arms and legs
- trouble sleeping
- dizziness
- headache
- tingling or numbness of the hands or feet
- hot flush
- shortness of breath
- unusual tiredness or weakness
- generally feeling unwell
- memory loss
- itchy skin
- decreased appetite
- dry mouth
- swelling, especially in the hands and feet
- cough

Tell your doctor if you have been diagnosed with:

- inflammation of the pancreas
- gallstones
- inflammation of the gallbladder
- elevation in some laboratory blood tests of liver or muscle function
- depression
- elevation in blood glucose levels
- diabetes
- high blood pressure
- thyroid function abnormalities

The above list includes the more common and uncommon side effects of your medicine. They are usually mild and short-lived.

Tell your doctor if you notice a significant increase in your need to urinate or if you are significantly more hungry or thirsty than usual.

Tell your doctor immediately if...

Tell your doctor immediately if you notice any of the following:

- skin rash and hives
- raised red rash, sometimes with circular-shaped lesions
- steady abdominal pain with nausea and vomiting
- joint pain

- bleeding or bruising more easily than normal
- yellowing of the skin and eyes
- dark coloured urine
- light coloured bowel motions
- problems with breathing including shortness of breath, persistent cough and fever

These may be serious side effects. You may need urgent medical attention.

Go to hospital if...

If any of the following happen, tell your doctor immediately or go to Accident and Emergency at your nearest hospital:

- swelling of the face, lips, mouth, throat or tongue which may cause difficulty in swallowing or breathing.
These may be serious side effects. If you have them, you may have had a serious allergic reaction to EZETIMIBE - ROSUVASTATIN SANDOZ. You may need urgent medical attention or hospitalisation. These side effects are very rare.
- chest pain
- unexpected muscle pain, tenderness, or weakness not caused by exercise, particularly if you also have a fever or generally feel unwell.
This may be a serious side effect. This is because on rare occasions muscle problems can be serious including muscle breakdown resulting in kidney damage. You may need urgent medical attention.
- severe skin problems.

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation.

Tell your doctor or pharmacist if you notice anything that is making you feel unwell.

Other side effects not listed above may also occur in some people.

AFTER TAKING EZETIMIBE – ROSUVASTATIN SANDOZ

Storage

Keep your tablets in a cool dry place where the temperature stays below 25°C.

Do not store EZETIMIBE - ROSUVASTATIN SANDOZ or any other medicine in the bathroom or near a sink. Do not leave it on a window sill or in the car.

Heat and dampness can destroy some medicines.

Keep it where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking EZETIMIBE - ROSUVASTATIN SANDOZ or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

PRODUCT DESCRIPTION

What it looks like

EZETIMIBE - ROSUVASTATIN SANDOZ comes in four different strengths:

- EZETIMIBE - ROSUVASTATIN SANDOZ 10 mg / 5 mg
- EZETIMIBE - ROSUVASTATIN SANDOZ 10 mg / 10 mg
- EZETIMIBE - ROSUVASTATIN SANDOZ 10 mg / 20 mg
- EZETIMIBE - ROSUVASTATIN SANDOZ 10 mg / 40 mg

Ezetimibe Sandoz 10 mg tablet is a white to almost-white, oval tablet with "10" marked on one side and "EZT" on the other side.

Rosuvastatin Sandoz 5 mg is a brown, round, film coated tablet.

Rosuvastatin Sandoz 10 mg is a brown, round, film coated tablet with "RSV 10" marked on one side.

Rosuvastatin Sandoz 20 mg is a brown, round, film coated tablet with "RSV 20" marked on one side.

Rosuvastatin Sandoz 40 mg is a brown, round, film coated tablet with "RSV 40" marked on one side.

Ezetimibe Sandoz 10mg and Rosuvastatin Sandoz (5, 10, 20, 40 mg) are supplied in 10-day calendar packs of 10 Ezetimibe and 10 Rosuvastatin tablets.

A trade pack contains 3 x 10-day calendar packs.

A starter pack contains 1 x 10-day calendar pack.

Ingredients

Ezetimibe Sandoz

Active ingredient:

- Ezetimibe 10 mg per tablet.

Inactive ingredients:

- Croscarmellose sodium
- Magnesium stearate
- Lactose
- Microcrystalline cellulose
- Hypromellose
- Sodium lauryl sulfate

Rosuvastatin Sandoz

Active ingredient:

- Rosuvastatin (as rosuvastatin calcium) 5 mg per tablet.
- Rosuvastatin (as rosuvastatin calcium) 10 mg per tablet.
- Rosuvastatin (as rosuvastatin calcium) 20 mg per tablet.
- Rosuvastatin (as rosuvastatin calcium) 40 mg per tablet.

Inactive ingredients:

- Purified talc
- Microcrystalline cellulose
- Silicon dioxide
- Colloidal anhydrous silica
- Maize starch
- Sodium stearyl fumarate
- Hypromellose
- Mannitol
- Lactose
- Macrogol 6000
- Titanium dioxide
- Iron oxide yellow
- Iron oxide red

Date of Preparation

This Consumer Medicine Information was revised in December 2023.

Ask your pharmacist about any updates.

Supplier

Sandoz Pty Ltd
100 Pacific Highway
North Sydney, NSW 2060
Australia
Tel: 1800 726 369

Australian Register Number(s)

EZETIMIBE - ROSUVASTATIN SANDOZ
10 mg / 5 mg -
Aust R 318875

EZETIMIBE - ROSUVASTATIN SANDOZ
10 mg / 10 mg -
Aust R 318873

EZETIMIBE - ROSUVASTATIN SANDOZ
10 mg / 20 mg -
Aust R 318874

EZETIMIBE - ROSUVASTATIN SANDOZ
10 mg / 40 mg -
Aust R 318872