

Blooms The Chemist Escitalopram

Contains the active ingredient, escitalopram (as escitalopram oxalate)

Consumer Medicine Information

For a copy of a large print leaflet, Ph: 1800 195 055

What is in this leaflet

This leaflet answers some common questions about this medicine. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking this medicine against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What this medicine is used for

The name of your medicine is Blooms The Chemist Escitalopram. It contains the active ingredient, escitalopram (as escitalopram oxalate).

Escitalopram is used to treat

- depression
- obsessive-compulsive disorder.

It belongs to a group of medicines called Selective Serotonin Reuptake Inhibitors (SSRIs). Escitalopram and other SSRIs are thought to help by increasing the amount of serotonin in your brain.

Depression is longer lasting or more severe than the "low moods" everyone has from time to time due to the stress of everyday life. It is thought to be caused by a chemical imbalance in parts of the brain. This imbalance affects your whole body and can cause emotional and physical symptoms such as feeling low in spirit, loss of interest in activities, being unable to enjoy life, poor appetite or overeating, disturbed sleep, often waking up early, loss of sex drive, lack of energy and feeling guilty over nothing.

Escitalopram corrects this chemical imbalance and may help relieve the symptoms of depression.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

Your doctor may have prescribed this medicine for another reason.

This medicine is available only with a doctor's prescription.

There is no evidence that escitalopram is addictive. However, if you suddenly stop taking it, you may get side effects.

Tell your doctor if you get any side effects after stopping escitalopram.

Use in children

Do not give this medicine to a child or adolescent.

There is no experience with its use in children and adolescents under 18 years of age.

Before you take this medicine

When you must not take it

Do not take this medicine if you have an allergy to:

- any medicine containing escitalopram, citalopram
- any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin

Do not take this medicine if you are pregnant or planning to become pregnant.

Medicines like escitalopram have been shown to reduce the quality of sperm in animal studies, which theoretically could affect fertility. If you are intending to start a family, ask your doctor for advice.

There have been reports that babies exposed to certain antidepressants during the third trimester of pregnancy may develop complications after birth.

Do not take this medicine whilst pregnant until you and your doctor have discussed the risks and benefits involved.

Do not breastfeed if you are taking this medicine.

It is not recommended that you breast-feed while taking this medicine because escitalopram passes into breast milk and may affect your baby.

Do not take this medicine whilst breast-feeding until you and your doctor have discussed the risks and benefits involved.

Do not take this medicine if you are taking the following other medicines:

- pimozide, used to treat disorders which affect the way you think, feel or act
- monoamine oxidase inhibitors (MAOIs), used to treat depression (phenelzine, tranylcypromine, moclobemide).
- monoamine oxidase inhibitors (MAOIs) such as linezolid which is an antibiotic and selegiline which is used in the treatment of Parkinson's Disease.

Do not take escitalopram until 14 days after stopping most MAOIs.

The exception is the MAOI, moclobemide, where you may take escitalopram one whole day after finishing taking moclobemide.

Similarly, do not take any MAOI until at least 14 days after stopping taking escitalopram.

Taking escitalopram with MAOIs may cause a serious reaction with signs such as a sudden increase in body temperature, very high blood pressure, rigid muscles, nausea/vomiting and/or fits (convulsions). Your doctor will know when it is safe to start escitalopram after the MAOI has been stopped.

Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start taking this medicine, talk to your doctor.

Before you start to take it

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you have or have had any of the following medical conditions:

- mania, hypomania, bipolar disorder or any other conditions which affect the way you think, feel or act
- epilepsy or convulsions, fits or seizures (you should avoid taking escitalopram if your epilepsy is not properly controlled; if it is properly controlled your doctor will wish to monitor you carefully if you take escitalopram)
- heart problems
- liver problems
- kidney problems
- problems with blood clotting or abnormal bleeding, i.e. a tendency to bleed or bruise easily
- thoughts or actions relating to self-harm or suicide
- diabetes
- a decreased level of sodium in your blood
- restlessness and/or a need to move often (akathisia)
- raised intraocular pressure (fluid pressure in the eye), or if you are at risk of angle-closure glaucoma.

Tell your doctor if you are receiving electroconvulsive therapy (ECT).

Tell your doctor if you are pregnant or plan to become pregnant or are breastfeeding.

Your doctor can discuss with you the risks and benefits involved.

If you have not told your doctor about any of the above, tell them before you start taking this medicine.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some combinations of medicines may increase the risk of serious side effects and are potentially life-threatening.

Therefore, some medicines MUST NOT be taken with escitalopram. These include:

- monoamine oxidase inhibitors, such as moclobemide, phenelzine, tranylcypromine, selegiline and linezolid
- pimozide.

Some other medicines may interact with escitalopram. These include:

- tryptophan, contained in some multivitamin and herbal preparations
- sumatriptan, used to treat migraines
- tramadol, a strong pain killer
- sumatriptan and similar medicines used to treat migraines and cluster headaches
- St John's Wort (Hypericum perforatum), a herbal remedy
- other medicines used to treat depression, including SSRIs, imipramine, clomipramine, nortriptyline and desipramine
- lithium, used to treat mood swings and some types of depression
- any other medicines used to treat anxiety, obsessive-compulsive disorder or premenstrual dysphoric disorder

- antipsychotics, medicines used to treat psychoses, schizophrenia and other conditions which affect the way you think, feel or act (e.g. risperidone, thioridazine and haloperidol)
- any other medicines affecting the chemicals in the brain
- prochlorperazine, used to prevent or treat severe nausea and vomiting
- bupropion, used to treat nicotine dependence
- mefloquine, an anti-malaria medicine
- fluconazole, an anti-fungal medicine
- some heart or blood pressure medications, e.g. dipyridamole, flecainide, propafenone, metoprolol
- medicines known to prolong bleeding e.g. aspirin or other non-steroidal anti-inflammatory drugs (NSAIDs) and anti-coagulants (such as warfarin and ticlopidine), which are used to prevent blood clots
- medicines used to treat reflux and ulcers, such as cimetidine, omeprazole, esomeprazole and lansoprazole
- imipramine and desipramine types of antidepressants.

These medicines may be affected by escitalopram or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking this medicine.

How to take this medicine

Follow carefully all directions given to you by your doctor.

Their instructions may be different to the information in this leaflet.

How much to take

Your doctor will tell you how much of this medicine you should take. This will depend on your condition and whether you are taking any other medicines.

The standard dose for this medicine is 10 mg per day.

Your doctor may increase your dose to 20 mg per day depending on how you respond to this medicine.

Elderly people may need smaller doses. The maximum dose for elderly people is 10 mg per day.

Patients with liver disease or with a lack of certain liver enzymes may receive a lower initial dose of 5 mg daily for the first two weeks. Your doctor may increase the dose to 10 mg daily.

How to take it

Swallow the tablets whole with a full glass of water.

Do not chew them.

When to take it

Take escitalopram as a single dose, either in the morning or in the evening.

Take this medicine at the same time each day.

Taking it at the same time each day will have the best effect and will also help you remember when to take it.

It does not matter if you take it before, with or after food.

How long to take it for

Continue taking your medicine for as long as your doctor tells you, even if it takes some time before you feel any improvement in your condition.

Make sure you have enough to last over weekends and holidays.

As with other medicines for the treatment of these conditions, it may take a few weeks before you feel any improvement.

Individuals will vary greatly in their response to escitalopram.

Your doctor will check your progress at regular intervals.

The length of treatment may vary for each individual but is usually at least 6 months.

In some cases, your doctor may decide that longer treatment is necessary.

Occasionally the symptoms of depression or other psychiatric conditions may include thoughts of harming yourself or committing suicide. It is possible that these symptoms may continue or increase until the full anti-depressant effect of your medicine becomes apparent.

You or anyone close to you or caring for you should watch for these symptoms and tell your doctor immediately or go to the nearest hospital if you have any distressing thoughts or experiences during this initial period or at any other time.

Also contact your doctor if you experience any worsening of your depression or other symptoms at any time during your treatment.

Stopping Treatment

Do not stop taking this medicine even if you begin to feel better.

Your doctor may decide that you should continue to take it for some time, even when you have overcome your problem. For best effect, this medicine must be taken regularly.

The underlying illness may persist for a long time and if you stop your treatment too soon, your symptoms may return.

Do not stop taking this medicine suddenly.

If you suddenly stop taking your medicine, you may experience mild, but usually temporary, symptoms such as dizziness, pins and needles, electric shock sensations, sleeping problems (vivid dreams, nightmares, inability to sleep), feeling anxious, restless or agitated, headaches, feeling sick (nausea), vomiting, sweating, tremor (shaking), feeling confused, feeling emotional or irritable, diarrhoea, visual disturbances, or fast or irregular heartbeats.

When you have completed your course of treatment, the dose of escitalopram is gradually reduced over a couple of weeks rather than stopped abruptly.

Your doctor will tell you how to reduce the dosage so that you help avoid getting side effects.

If you forget to take it

If you missed a dose and remember in less than 12 hours, take it straight away, and then go back to taking it as you would normally.

Otherwise, if you are more than 12 hours late, skip the dose you missed and take the next dose when you are meant to.

Do not take a double dose to make up for missed doses.

This may increase the chance of you experiencing side effects.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints to help you remember.

If you take too much (overdose)

If you think that you or anyone else may have taken too much of this medicine, immediately telephone your doctor or the Poisons Information Centre (Tel: 13 11 26 in Australia) for advice. Alternatively go to the

Accident and Emergency Department at your nearest hospital.

Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

If you take too much escitalopram, you may get symptoms of drowsiness, sleepiness, dizziness, high or low blood pressure, nausea (feeling sick), vomiting, agitation or tremor (shaking), fast or slow heart beat or change in heart rhythm, dilated pupils or, rarely, temporary paralysis or weakness of muscles, convulsions or coma.

A condition called serotonin syndrome may occur, with high fever, agitation, confusion, trembling and abrupt contraction of muscles.

While you are taking this medicine

Things you must do

People taking escitalopram may be more likely to think about killing themselves or actually trying to do so, especially when escitalopram is first started, or the dose is changed. Tell your doctor immediately if you have thoughts about killing yourself or if you are close to or care for someone using escitalopram who talks about or shows signs of killing him or herself.

All mentions of suicide or violence must be taken seriously.

Occasionally, the symptoms of depression may include thoughts of suicide or self-harm. It is possible that these symptoms continue or get worse until the full antidepressant effect of the medicine becomes apparent. This is more likely to occur if you are a young adult, i.e. 18 to 24 years of age, and you have not used antidepressant medicines before.

If you or someone you know or care for demonstrates any of the following warning signs of suicide-related behaviour while taking escitalopram, contact a doctor immediately, or even to go to the nearest hospital for treatment:

- thoughts or talk of death or suicide
- thoughts or talk of self-harm or harm to others
- any recent attempts of self-harm
- increase in aggressive behaviour, irritability or agitation
- worsening of depression.

Follow your doctor's instructions. Do not stop taking this medicine or change the dose without consulting your doctor, even if you experience increased anxiety at the beginning of treatment.

At the beginning of treatment, some patients may experience increased anxiety, which will disappear during continued treatment.

Tell your doctor immediately if you experience symptoms such as restlessness or difficulty sitting or standing still.

These symptoms can also occur during the first weeks of treatment.

Contact your doctor as soon as possible if you suddenly experience an episode of mania.

Some people with bipolar disorder (manic depression) may enter into a manic phase. Symptoms of mania include lots of rapidly changing thoughts or ideas, exaggerated gaiety, being much more physically active and much more restless.

Sometimes you may not know that you are manic, so it may be helpful to have a friend or relative watch over you for any possible signs of change in your behaviour.

Visit your doctor regularly so they can check on your progress.

Tell your doctor immediately if you become pregnant. If you are a woman of child-bearing age, you should avoid becoming pregnant while taking escitalopram.

Make sure your midwife and/or doctor know you are taking escitalopram. When taken during pregnancy, particularly in the last 3 months of pregnancy, medicines like escitalopram may increase the risk of a serious condition in babies, called persistent pulmonary hypertension of the newborn (PPHN), making the baby breathe faster and appear bluish. These symptoms usually begin during the first 24 hours after the baby is born. If this happens to your baby, you should contact your midwife and/or doctor immediately.

If used during pregnancy, escitalopram should never be stopped abruptly.

Tell your doctor if you get a headache or start to feel sick, restless, irritated, confused or fatigued or if you vomit or have fits, muscle weakness or spasms.

Some people (especially older people or those taking diuretics/water tablets) may experience low or a lack of sodium in the blood when taking this medicine.

If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking this medicine.

Tell any other doctors, dentists, and pharmacists who treat you that you are taking this medicine.

If you are going to have surgery, tell the surgeon or anaesthetist that you are taking this medicine.

It may affect other medicines used during surgery.

Tell your doctor that you are taking this medicine if you are breast-feeding or are planning to breastfeed.

If you are about to have any blood tests, tell your doctor that you are taking this medicine.

Keep all your doctor's appointments so that your progress can be checked.

Your doctor may occasionally do tests to make sure the medicine is working and to prevent side effects.

Go to your doctor regularly for a check-up.

Tell your doctor if, for any reason, you have not taken your medicine exactly as prescribed. Otherwise your doctor may think that it was not effective and change your treatment unnecessarily.

Tell your doctor if you feel this medicine is not helping your condition.

If you are being treated for depression, be sure to discuss with your doctor any problems you may have and how you feel, especially any feelings of severe sadness, thoughts of suicide, bursts of unusual energy, anger or aggression, or if you become particularly agitated or restless.

Tell your doctor immediately if you have any suicidal thoughts or other mental/mood changes.

Make sure you have enough tablets to last over weekends and holidays.

Things you must not do

Do not take this medicine to treat any other complaints unless your doctor tells you to.

Do not give your medicine to anyone else, even if they have the same condition as you.

Do not stop taking your medicine or change the dosage without checking with your doctor.

Do not let yourself run out of medicine over the weekend or on holidays.

Suddenly stopping escitalopram may cause unwanted discontinuation symptoms, such as

dizziness, sensory disturbances, sleep disturbances, agitation or anxiety, vomiting, tremor, confusion, sweating, headache, diarrhoea, palpitations, emotional instability, irritability, visual disturbances and nausea. Your doctor will tell you when and how escitalopram should be discontinued. You doctor will gradually reduce the amount you are using, usually over a period of one to two weeks, before stopping completely.

Things to be careful of

Be careful when driving or operating machinery until you know how this medicine affects you.

This medicine may cause nausea, fatigue, drowsiness, sight problems or dizziness in some people, especially early in the treatment. If you have any of these symptoms, do not drive, operate machinery, or do anything else that could be dangerous.

Avoid alcohol while you are taking this medicine.

It is best not to drink alcohol while you are being treated for depression.

You should be aware that people over 50 years of age who take antidepressants have an increased risk of having a bone fracture.

Side effects

Tell your doctor as soon as possible if you do not feel well while you are taking escitalopram or if you have any questions or concerns.

All medicines can have side effects. Sometimes they are serious, but most of the time, they are not.

Do not be alarmed by the following lists of side effects. You may not experience any of them.

Tell your doctor if you notice any of the following:

- feeling tired and weak (fatigued), hot flushes, fever, feeling unwell, shaking or tremors, migraine, headache, or giddiness
- muscle, back, bone, nerve or joint pain, stiffness, weakness or cramps, decrease or loss of touch or other senses
- increased or decreased sensitivity to outside stimuli
- feeling or being sick, reflux, diarrhoea or loose bowel motions, constipation, indigestion, stomach pain or discomfort, wind, burping, hiccups, problems swallowing, sore mouth, tongue or throat, haemorrhoids (piles)
- dry mouth, feeling thirsty increased saliva, taste disturbance
- fatigue, sleepiness or drowsiness, yawning, sleeping difficulties, strange or terrifying dreams
- teeth grinding or clenching
- increased or decreased appetite, weight loss
- excessive and/or abnormal movements
- increased muscle tension, muscle twitching
- sexual disturbances (decreased sexual drive; problems with ejaculation or erection; women may experience difficulties achieving orgasm) or prostate problems
- symptoms of hyperglycaemia (high blood sugar) – feeling hungry, thirsty and/or frequent or excessive urination
- problems with eyes or eyesight
- dizziness when you stand up suddenly, due to low blood pressure
- unable to tolerate alcohol
- menstrual irregularities, period pain, breast pain, unusual vaginal bleeding

- loss of bladder control unusual hair loss or thinning
- tingling or numbness of the hands or feet
- breast enlargement or unusual secretion of breast milk in men or women
- mild rash, or itching or prickling of the skin
- acne, eczema, dermatitis, dry skin, psoriasis or other skin problem
- pain of any type
- ringing or other persistent noise in the ears, problems hearing or earache
- increased or decreased sweating
- bruises
- osteoporosis
- tooth or jaw problems
- flu-like symptoms, runny or blocked nose, sneezing, facial pressure or pain, coughing or sore throat

Tell your doctor as soon as possible if you notice any of the following:

- becoming nervous, confused, forgetful, unable to concentrate, agitated, confused, panicky or anxious
- feeling restless or unable to sit still
- stomach pain with nausea and vomiting of blood, or blood in the bowel movements
- aggression, worsening of depression
- general swelling or swollen hands, ankles, feet or face or eye area due to fluid build-up
- problems speaking
- feelings of not being part of your body, or in a daze
- feeling sick or unwell with weak muscles or feeling confused (these symptoms may be signs of a rare condition as a result of low levels of sodium in the blood, which may be caused by antidepressants and occurs especially in elderly women) increased tendency to bleed, develop bruises or broken bones
- passing more or less urine than normal, or problems when urinating, or bladder infection
- abnormal liver function tests (increased amount of liver enzymes)
- flushing, varicose veins
- infection in any part of your body
- dizziness
- agitation, anxiety, feeling tense and restless, tired, drowsy, lack of energy, irritable, problems sleeping, headache, nausea and tingling or numbness of the hands and feet after stopping escitalopram.

The above list includes serious side effects that may require medical attention. Serious side effects are rare.

If you experience any of the following, stop taking your medicine and contact your doctor immediately or go to the Accident and Emergency department at your nearest hospital:

- seizures, tremors, movement disorders (involuntary movements of the muscles or being unco-ordinated).
- coma (unconsciousness)
- a collection of symptoms including weight gain (despite loss of appetite), feeling and being sick, muscle weakness and irritability
- severe rash, with blisters and/ or excessive peeling of skin and also possibly severe blisters and bleeding in the lips, eyes, mouth, nose and genitals

- a sudden increase in body temperature, very high blood pressure, rigid muscles, nausea/vomiting and/or fits (convulsions). These symptoms may be signs of a rare condition called Serotonin Syndrome.
- Neuroleptic Malignant Syndrome (a serious reaction to some medicines with a sudden increase in body temperature, extremely high blood pressure and severe convulsions)
- fast, slow or irregular heartbeat, high blood pressure
- palpitations, fainting or chest pain or tightness
- abnormal bleeding
- kidney pain, difficulty in passing urine, dark coloured urine or blood in the urine
- a collection of symptoms including fever, sore throat, swollen glands, mouth ulcers, unusual bleeding or bruising under the skin
- mania (mood of excitement, over-activity and uninhibited behaviour or aggression), hallucinations (hearing, seeing or feeling things that are not there)
- jaundice (yellowing of the skin and/or eyes), with or without other signs of hepatitis or liver problems (loss of appetite, tiredness, feeling or being sick, dark urine, stomach pain or swelling, confusion, unconsciousness).
- feeling paranoid, panicky, or "high" or having mood swings or feeling more depressed or in a trance
- thoughts of suicide or attempting suicide or self-harm
- sudden, severe breathing problems
- sudden weakness or numbness of the face, arms or legs, especially on one side, slurred speech
- symptoms of an allergic reaction including cough, shortness of breath, wheezing or difficulty breathing; swelling of the face, lips, tongue, throat or other parts of the body; rash, itching or hives on the skin

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation. These side effects are very rare.

Tell your doctor or pharmacist if you notice anything that is making you feel unwell.

Other side effects not listed above may also occur in some people.

Storage and disposal

Storage

Keep your medicine in its original packaging until it is time to take it.

If you take your medicine out of its original packaging, it may not keep well.

Keep your medicine in a cool dry place where the temperature will stay below 30°C.

Do not store your medicine, or any other medicine, in the bathroom, or near a sink.

Do not leave it on a windowsill or in the car. Heat and dampness can destroy some medicines.

Keep this medicine where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking this medicine, or it has passed its expiry date, your pharmacist can dispose of the remaining medicine safely.

Product description

What Blooms The Chemist Escitalopram looks like

10 mg tablets

White to off -white, oval, biconvex, film-coated tablets with "C4" debossed on one side and a notch break-line on the other side. AUST R 272630.

20 mg tablets

White to off -white, oval, biconvex, film-coated tablets with "C3" debossed on one side and a notch break-line on the other side. AUST R 272631.

Blister packs of 28 tablets.

* Not all strengths may be available.

Ingredients

Each tablet contains 10 mg or 20 mg of escitalopram (as oxalate) as the active ingredient.

It also contains the following inactive ingredients:

- microcrystalline cellulose
- colloidal anhydrous silica
- hypromellose
- magnesium stearate
- croscarmellose sodium
- purified talc
- macrogol 400
- titanium dioxide.

This medicine is gluten-free, lactose free, sucrose-free, tartrazine-free and free of other azo dyes.

Sponsor

Apotex Pty Ltd
16 Giffnock Avenue
Macquarie Park NSW 2113
Tel: (02) 8877 8333
Web: www1.apotex.com/au

This leaflet was last updated in May 2021.