

## Famciclovir Sandoz®

### for shingles

Famciclovir tablets

## Consumer Medicine Information

### WHAT IS IN THIS LEAFLET

This leaflet answers some common questions about Famciclovir Sandoz.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking this medicine against the benefits they expect it will have for you.

**If you have any concerns about taking this medicine, ask your doctor or pharmacist.**

**Keep this leaflet with the medicine.**

**You may need to read it again.**

### WHAT FAMCICLOVIR SANDOZ IS USED FOR

Famciclovir Sandoz is an antiviral medicine that is used to treat shingles in adults and adolescents.

Shingles is a condition that is caused by a herpes virus called varicella zoster, the same virus that causes chicken pox. The virus can become active again in the body, even after many years, resulting in shingles. The main symptom is a rash that can cause pain, burning and blisters.

Although Famciclovir Sandoz does not cure the viral infection, it helps to relieve the symptoms and shorten their duration.

The best results are obtained if the medicine is started as soon as possible after the first symptoms begin to appear.

Ask your doctor if you have any questions about why this medicine has been prescribed to you.

Your doctor may have prescribed it for another reason.

Famciclovir Sandoz is only available with a doctor's prescription. It is not addictive.

This medicine is not recommended for use in children under 12 years of age.

### BEFORE YOU TAKE FAMCICLOVIR SANDOZ

#### *When you must not take it*

**Do not take this medicine if you have an allergy to:**

- famciclovir, the active ingredient
- penciclovir, a related antiviral medicine.
- any of the other ingredients of Famciclovir Sandoz listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include:

- shortness of breath, wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin.

**Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.**

In that case, return it to your pharmacist.

#### *Before you start to take it*

**Tell your doctor if you have a problem with:**

- your body's immune system, which helps to fight off infections
- your kidneys
- your liver.

Your doctor may want to take extra precautions in that case.

**Tell your doctor if you are pregnant or plan to become pregnant or are breast-feeding.**

FAMCICLOVIR SANDOZ should not be used during pregnancy unless necessary. Your doctor will discuss with you the potential risks of taking FAMCICLOVIR SANDOZ during pregnancy, and will also advise you if you should take FAMCICLOVIR SANDOZ while breast-feeding, based on the benefits and risks of your personal situation.

**Tell your doctor if you are lactose intolerant.**

Famciclovir Sandoz 250 mg tablets contain lactose.

**Tell your doctor if you are allergic to any other medicines, foods, dyes or preservatives.**

**Your doctor will want to know if you are prone to allergies.**

**If you experience an allergic reaction, stop using the medicine and inform your doctor or pharmacist immediately.**

#### *Taking other medicines*

**Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.**

Some medicines and Famciclovir Sandoz may interfere with each other. These include:

- probenecid, a prescription medicine used to treat gout (a disease with painful, swollen joints caused by uric acid crystals) and to increase blood levels of penicillin-type antibiotics
- raloxifene, a medicine used to treat osteoporosis (a disease which causes bones to become less dense, gradually making them weaker, more brittle and likely to break)
- medicines that can affect your kidneys.

You may need to take different amounts of these medicines or you may need to take different medicines.

Your doctor and pharmacist have more information.

If you have not told your doctor about any of these things, tell him/her before you start taking this medicine.

### HOW TO TAKE FAMCICLOVIR SANDOZ

**Swallow the tablets whole with a full glass of water.**

The tablets may be taken with or without food. It is not necessary to chew or crush the tablets.

**Follow all directions given to you by your doctor and pharmacist carefully.**

These instructions may differ from the information contained in this leaflet.

**If you do not understand the instructions on the label, ask your doctor or pharmacist for help.**

#### *How much to take*

The usual dose is one 250 mg tablet three times each day for seven days, beginning no later than 72 hours after the rash appears.

Your doctor may have prescribed a different dose.

For people whose immune system does not work as well as it should, the dose and duration of treatment may be increased.

For people who have reduced kidney function, the dose may be reduced.

**Follow your doctor's instructions on how many Famciclovir Sandoz tablets to take.**

**Ask your doctor or pharmacist if you are unsure of the correct dose for you.**

They will tell you exactly how much to take.

**Follow the instructions that they give you.**

#### *When to take Famciclovir Sandoz*

**Unless your doctor tells you otherwise, take one tablet when you get up in the morning, one in the afternoon and one just before going to bed at night.**

**Try to take the medicine at about the same time each day.**

#### *How long to take Famciclovir*

**Continue taking Famciclovir Sandoz every day for the full course of treatment.**

To help clear up your infection, you must take the full course of treatment, even if your symptoms begin to clear up after a few days.

#### *If you forget to take it*

Take your dose as soon as you remember. Take your next tablet at the usual time, and then go back to taking it as you would normally.

**Do not take two doses within a time frame of less than one hour. In that case, skip the missed dose.**

**Do not take a double dose to make up for the one that you missed.**

This may increase the chance of you getting an unwanted side effect.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

#### *If you take too much (overdose)*

**Immediately telephone your doctor or the Poisons Information Centre (telephone Australia 13 11 26 or New Zealand 0800 POISON or 0800 764766) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much Famciclovir Sandoz. Do this even if there are no signs of discomfort or poisoning. Keep the telephone number for these places handy.**

Taking too much Famciclovir Sandoz may affect your kidneys. In people who already have kidney problems it may, rarely, lead to kidney failure if their dose is not correctly lowered.

---

## WHILE YOU ARE TAKING FAMCICLOVIR SANDOZ

---

### *Things you must do*

**If you become pregnant while taking this medicine, tell your doctor.**

Your doctor can discuss with you the risks of taking it while you are pregnant.

**If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking Famciclovir Sandoz.**

**Tell any other doctors, dentists or pharmacists who treat you that you are taking this medicine.**

### *Things you must not do*

**Do not give your medicine to anyone else even if their condition seems to be the same as yours.**

**Do not use it to treat any other complaints unless your doctor tells you to.**

**Do not stop taking your tablets or change the dosage without checking with your doctor first.**

If you stop your tablets suddenly, your condition may worsen or you may have unwanted side effects.

### *Things to be careful of*

**If you are pregnant or breast feeding, ask your doctor or pharmacist for advice before taking any medicine.**

**Be careful driving, operating machinery or doing jobs that require you to be alert until you know how Famciclovir Sandoz affects you.**

This medicine can cause dizziness, sleepiness or confusion in some people.

### *Things that may help your condition*

**Take the following precautions to avoid spreading the virus:**

- keep the areas affected by the virus as clean and dry as possible
- wear loose-fitting clothing to avoid irritating the rash
- avoid touching or scratching the sore area as you may spread the virus on your fingers.

---

## SIDE EFFECTS

---

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Famciclovir Sandoz.**

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

**Do not be alarmed by the following lists of possible side effects. You may not experience any of them.**

**Ask your doctor or pharmacist to answer any questions you may have.**

**Tell your doctor if you notice any of the following and they worry you:**

- headache
- dizziness
- nausea (feeling sick) or vomiting
- abdominal pain
- diarrhoea
- itching or an itchy rash (urticaria)
- abnormal liver function test results.

The above side effects are usually mild.

**Tell your doctor as soon as possible if you notice any of the following:**

- a rash that is separate from the shingles rash
- extreme sleepiness or confusion, usually in older people
- hallucinations (seeing or hearing things that are not really there)
- painful or swollen joints
- aching muscles or muscle tenderness or weakness that is not caused by exercise
- yellowing of the skin or eyes (sign of jaundice)
- palpitation (signs of abnormal heart beat)

The above side effects may require medical attention.

**If any of the following happen, tell your doctor immediately or go to Accident and Emergency at your nearest hospital:**

- swelling below the surface of the skin (e.g. swelling around the face, eye, eyelid or throat)
- unexplained bruising, reddish or purplish patches on the skin or bleeding more easily than usual as it may indicate that the number of platelets (a type of blood cell responsible for blood clotting) in your blood are reduced
- severe blistering of the skin or mucous membranes of the lips, eyes, mouth, nasal passages of genitals (signs of a serious skin reaction)
- purple patches, itching, burning of the skin (signs of inflamed blood vessels)
- seizure or fits
- difficulty breathing or swallowing, wheezing or cough, light headedness, changes in alertness, skin reddening, facial/throat swelling, blue discoloration of the lips, tongue or skin (signs of severe allergic reaction).

The above side effects are very rare.

**Tell your doctor if you notice anything else that is making you feel unwell.**

Other side effects not listed here or not yet known may happen in some people.

---

## AFTER USING FAMCICLOVIR SANDOZ

---

### *Storage*

- Keep your medicine in the original container until it is time to take it.
- Store in a dry place at room temperature.
- Do not store Famciclovir Sandoz or any other medicine in the bathroom or near a sink.
- Do not leave it on a window sill or in the car.

Heat and dampness can destroy some medicines. Famciclovir Sandoz tablets will keep best if they are stored cool and dry.

**Keep the medicine where children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

### *Disposal*

If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

---

## PRODUCT DESCRIPTION

---

### *What it looks like*

Famciclovir Sandoz 250mg Tablets are white, round film-coated tablets marked with "FV" on one side and "250" on the other. Each carton contains 20, 21 or 56 tablets.

Famciclovir Sandoz 500mg Tablets are white, oval film-coated tablets marked with "FV 500" on one side. Each carton contains 30 or 56 Tablets.

### *Ingredients*

Active ingredients:

- Famciclovir Sandoz 250mg - contain 250mg famciclovir per tablet
- Famciclovir Sandoz 500mg - contain 500mg famciclovir per tablet.

Inactive ingredients:

Famciclovir Sandoz 250 mg film-coated tablets contain:

- hypromellose (E 464)
- hydroxypropylcellulose (E 463)
- lactose
- macrogol
- magnesium stearate (E 572)
- sodium starch glycolate
- titanium dioxide (E171).

Famciclovir Sandoz 250 mg tablets do not contain sucrose, gluten, tartrazine or any other azo dyes.

Famciclovir Sandoz 500 mg tablets contain:

- hypromellose (E 464)
- hydroxypropylcellulose (E 463)
- macrogol
- magnesium stearate (E 572)
- sodium starch glycolate
- titanium dioxide (E171).

Famciclovir Sandoz 500 mg tablets do not contain lactose, sucrose, gluten, tartrazine or any other azo dyes.

### *Supplier*

Sandoz Pty Ltd  
ABN 60 075 449 553  
54 Waterloo Road  
Macquarie Park NSW 2113  
Australia  
Tel: 1800 634 500

This leaflet was revised in September 2016.

### **Australian Register Numbers**

Famciclovir Sandoz 250 mg tablets: AUST R 162313

Famciclovir Sandoz 500 mg tablets: AUST R 162314