

## Consumer Medicine Information

### What is in this leaflet

This leaflet answers some common questions about Exjade.

The information in this leaflet was last updated on the date listed on the final page. More recent information on the medicine may be available.

**You should ensure that you speak to your pharmacist or doctor to obtain the most up to date information on the medicine. You can also download the most up to date leaflet from [www.novartis.com.au](http://www.novartis.com.au).**

Those updates may contain important information about the medicine and its use of which you should be aware.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking this medicine against the benefits they expect it will provide.

**If you have any concerns about this medicine, ask your doctor or pharmacist.**

**Keep this leaflet with the medicine.**

You may need to read it again.

### What Exjade is used for

Exjade is used to treat a condition called iron overload, which happens when the body has too much iron. This can occur after repeated blood transfusions.

The body has no natural way to remove excess iron which comes with blood transfusions.

Exjade is also used to treat patients who have iron overload associated with their thalassemia syndromes, but who are not transfusion dependent. In patients with non-transfusion-dependent thalassemia syndromes, iron overload may develop over time due to increased absorption of dietary iron in response to low blood cell counts.

Over time, this excess iron can damage important organs such as the liver and heart.

This medicine contains an active substance called deferasirox. It attaches itself to the iron molecules to remove the excess iron from the body. This will help prevent iron-induced organ damage.

**Exjade is to be taken every day.**

This type of medicine must be taken every day to help remove the excess iron from your body.

**Ask your doctor if you have any questions about why this medicine has been prescribed for you.**

Your doctor may have prescribed it for another reason.

This medicine is not addictive.

This medicine is available only with a doctor's prescription.

### Before you take Exjade

*When you must not take it*

**Do not take Exjade if you have an allergy to deferasirox, the active ingredient, or to any of the other ingredients listed at the end of this leaflet.**

Some of the symptoms of an allergic reaction may include shortness of breath, wheezing or difficulty breathing; swelling of the face, lips, tongue or other parts of the body; rash, itching or hives on the skin.

If you develop a severe rash, or experience difficulty breathing and dizziness or swelling mainly of the face and throat (signs of severe allergic reaction) STOP taking Exjade immediately and tell your doctor straight away.

**Do not take this medicine if you have severe kidney problems.**

**Do not take this medicine if you have low platelet counts.**

**Do not take this medicine if you have an advanced stage of myelodysplastic syndrome (MDS) or advanced cancer.**

**Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.**

In that case, return it to your pharmacist.

*Before you start to take it*

**Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.**

**Tell your doctor if you have a low level of platelets in your blood test.**

**Tell your doctor if you have or have had any problems with your kidneys or liver.**

Your doctor may want to take special precautions in this case.

**Tell your doctor if you are pregnant or plan to become pregnant or are breast-feeding.**

Your doctor can discuss with you the risks and benefits involved.

**Tell your doctor if you have an intolerance to lactose.**

This medicine contains lactose.

*Taking other medicines*

**Tell your doctor or pharmacist if you are taking any other medicines, including any that you buy without a prescription from your pharmacy, supermarket or health food shop.**

Some medicines and Exjade may interfere with each other. These include:

- cyclosporin, used in transplantation to prevent organ rejection or to treat rheumatoid arthritis or atopic dermatitis
- midazolam (a medicine used to sedate)
- simvastatin (medicines used to lower cholesterol)
- hormonal contraceptive agents (birth control medicines). Their effectiveness may be reduced while taking Exjade.
- certain painkillers or anti-inflammatory medicines (e.g. aspirin, ibuprofen, corticosteroids)
- oral bisphosphonates (medicines used to treat osteoporosis)
- anticoagulant medicines (medicines used to prevent or treat blood clotting)
- antacid preparations containing aluminium, which should not be taken at the same time of day as Exjade
- a medicine for type 2 diabetes called repaglinide

- certain medicines for epilepsy or sedation (phenytoin, phenobarbitone)
- a medicine for HIV called ritonavir
- a medicine for tuberculosis called rifampicin
- a medicine for cancer called paclitaxel
- a medicine used to remove bile acids called cholestyramine
- theophylline (used to treat respiratory diseases such as asthma).
- busulfan (used as treatment prior to bone marrow transplant)

Other medicines that are processed like theophylline in the body and your doctor should know about include: clozapine, cyclobenzaprine, imipramine, haloperidol, fluvoxamine, mexiletine, naproxen, olanzapine, riluzole, tacrine, tizanidine, zileuton and zolmitriptan.

Your doctor may need to test the level of some of these medicines in your blood. Your doctor will take these tests into consideration when deciding on the most suitable dose of these medicines for you.

You may need to take different medicines. Your doctor and pharmacist have more information.

**If you have not told your doctor about any of these things, tell him/her before you start taking this medicine.**

### How to take Exjade

**Follow all directions given to you by your doctor or pharmacist carefully.**

They may differ from the information contained in this leaflet.

**If you do not understand the instructions on the label, ask your doctor or pharmacist for help.**

*How much to take*

The usual starting dose is 20 mg per kilogram body weight each day for patients receiving regular blood transfusions.

For patients NOT receiving regular blood transfusions, the usual starting dose is 10 mg per kilogram body weight.

A higher or lower starting dose may be recommended by your doctor based on your individual treatment needs.

Depending on your response, your doctor may increase the dose to a maximum 40 mg per kilogram body weight each day if you receive regular blood transfusions or 20 mg per kilogram body weight if you are NOT receiving regular blood transfusions.

*How to take it*

**Disperse the required number of tablets completely by stirring in a 100 - 200 mL glass of water or orange juice or apple juice until a suspension forms.**

When dispersing in juice, it helps to first disperse the tablets in about two tablespoons of water, before diluting with juice.

**Drink the entire contents of the glass, then add a little water or juice to what is left in the empty glass and drink that too.**

**Do not disperse the tablets in fizzy drinks or milk.**

**Do not chew, split or crush the tablets. Do not swallow them whole.**

### *When to take it*

**Take Exjade once a day, every day, at about the same time each day. Take it on an empty stomach at least 30 minutes before eating any food.**

Taking it at the same time each day, 30 minutes before eating food will have the best effect. It will also help you remember when to take it.

### *How long to take it*

**Do not stop taking Exjade unless your doctor tells you to - even if you feel well.**

This medicine helps to control your condition, but does not cure it. Therefore, you must take it every day. Continue taking the medicine as long as your doctor tells you to.

### *If you forget to take it*

**If it is almost time for your next dose, skip the dose you missed and take the next dose when you are meant to.**

**Otherwise, take it as soon as you remember, and then go back to taking it as you would normally.**

**Do not take a double dose to make up for the dose that you missed.**

This may increase the chance of you getting an unwanted side effect.

**If you have trouble remembering to take your medicine, ask your pharmacist for some hints.**

### *If you take too much (overdose)*

**Immediately telephone your doctor or the Poisons Information Centre (telephone 13 11 26) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much Exjade. Do this even if there are no signs of discomfort or poisoning. Keep the telephone numbers for these places handy.**

You may experience nausea, vomiting, headache or diarrhoea if you have taken too much Exjade.

---

### *While you are taking Exjade*

---

#### *Things you must do*

**If you notice substantially reduced urine output, tell your doctor straight away.**

This could be a sign of a problem with your kidneys.

**Elderly patients should be monitored closely by their doctor.**

Elderly patients may experience more side effects than younger patients and their doctor may adjust their dose.

**Keep all of your doctor's appointments so that your progress can be checked.**

Your doctor will do regular tests (blood, urine or Magnetic Resonance Imaging (MRI)) to make sure the medicine is working and to prevent unwanted side effects. Your doctor will also use these tests to decide when you should stop taking Exjade.

Your eyesight and hearing may be tested during treatment as a precautionary measure.

**If you become pregnant while taking this medicine, tell your doctor immediately.**

It may affect your developing baby. Your doctor will discuss the risks and benefits of continuing treatment in this case.

**If you are going to have surgery, tell the surgeon or anaesthetist that you are taking this medicine.**

It may affect other medicines used during surgery.

**If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking this medicine.**

**Tell any other doctors, dentists, and pharmacists who treat you that you are taking this medicine.**

### *Things you must not do*

**Do not give this medicine to anyone else, even if their condition seems similar to yours.**

**Do not use it to treat any other complaints unless your doctor tells you to.**

### *Things to be careful of*

**Be careful driving, operating machinery or doing jobs that require you to be alert until you know how Exjade affects you.**

This medicine may cause dizziness in some people. If you experience dizziness, do not drive, operate machinery or do anything else that could be dangerous. Children should be careful when riding bicycles or climbing trees.

---

### *Side effects*

---

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Exjade even if you do not think it is connected with the medicine.**

All medicines can have side effects. Sometimes they are serious, but most of the time they are not. You may need medical treatment if you get some of the side effects.

**Elderly patients may experience more side effects than younger patients.**

Do not be alarmed by the following lists of side effects. You may not experience any of them. Ask your doctor or pharmacist to answer any questions you may have.

**STOP taking Exjade and seek medical help immediately if you or your child experience any of the following symptoms which may be signs of an allergic reaction:**

- difficulty in breathing and swallowing
- swelling of the face, lips, tongue or throat
- severe itching of the skin, with a red rash or raised bumps

**STOP taking Exjade and tell your doctor immediately if you notice any of the following:**

- rash, red skin, blistering of the lips, eyes or mouth, skin peeling, high fever, flu-like symptoms and enlarged lymph nodes (signs of severe skin reaction)
- a substantially reduced urine output (sign of kidney problem)
- drowsiness, upper right abdominal pain, yellowing of your skin or eyes and dark urine (sign of liver problems)
- vomiting with blood and/or black stools
- frequent heartburn or abdominal pain (ulcers), particularly after eating or taking the drug
- blurred, cloudy or partial loss of vision
- hearing disturbances
- sudden back pain or pain on the right side of the abdomen (signs of gallstones)

- severe upper stomach pain (pancreatitis)
- tear in stomach or intestine wall that can be painful and cause nausea

The above list includes serious side effects which may require medical attention. These side effects do not occur frequently.

**Tell your doctor or pharmacist if you notice any of the following and they worry you:**

- nausea, diarrhoea, vomiting, pain in the abdomen, bloating, constipation or indigestion
- rash
- headache
- itching
- fever or symptoms of a cold or flu
- cough or sore throat
- dizziness
- swelling of arms or legs
- change in the colour of the skin
- anxiety
- trouble sleeping
- tiredness
- sore muscles or joints
- hair loss

The above list includes the more common side effects of your medicine. They are usually mild and short-lived.

**Tell your doctor or pharmacist if you notice anything that is making you feel unwell.**

Some people may have other side effects not yet known or mentioned in this leaflet. Some of these side effects, for example, changes in kidney and liver function, can only be found when your doctor does tests from time to time to check your progress.

---

### *After using Exjade*

---

#### *Storage*

- Keep your medicine in the original container until it is time to take it.
- Store it in a cool, dry place, below 30°C
- Do not store this medicine or any other medicine in the bathroom or near a sink
- Do not leave it in the car or on window sills.

**Keep the medicine where children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

#### *Disposal*

**If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine you have left over.**

---

### *Product description*

---

#### *What it looks like*

Exjade tablets are off-white, round, flat tablets with bevelled edge. Exjade 125 mg tablets are imprinted with "J125" on one side and "NVR" on the other. Exjade 250 mg tablets are imprinted with "J250" on one side and "NVR" on the other. Exjade 500 mg tablets are imprinted with "J500" on one side and "NVR" on the other.

Exjade comes in packs of 28 dispersible tablets in blister packs.

### *Ingredients*

Contains either 125, 250 or 500 mg of deferasirox as the active ingredient. It also contains:

- lactose
- crospovidone
- povidone
- sodium lauryl sulphate
- microcrystalline cellulose
- colloidal anhydrous silica
- magnesium stearate

### *Sponsor*

EXJADE® is supplied in Australia by:

NOVARTIS Pharmaceuticals Australia Pty Limited

ABN 18 004 244 160

54 Waterloo Road

North Ryde NSW 2113

Telephone 1 800 671 203

Web site: [www.novartis.com.au](http://www.novartis.com.au)

® = Registered Trademark

This leaflet was prepared on 17 August 2018

Australian Registration Number.

Exjade 125 mg AUST R 119230

Exjade 250 mg AUST R 119231

Exjade 500 mg AUST R 119232

Internal Document Code:

(exj170818c.doc) based on PI

(exj170818i.doc)