

# CMV Immunoglobulin-VF

*Human Cytomegalovirus Immunoglobulin, solution for intravenous infusion.*

---

## Consumer Medicine Information

---

### What is in this leaflet

---

This leaflet answers some common questions about CMV Immunoglobulin-VF. The letters CMV stand for cytomegalovirus, a type of virus. It does not contain complete information about CMV Immunoglobulin-VF. It does not take the place of talking to your doctor.

**If you have any concerns about using this medicine, ask your doctor. Follow your doctor's advice even if it is different from what this leaflet says.**

**Please read this leaflet carefully and keep it for future reference.**

The information in this leaflet is subject to change. Please check with your doctor whether there is any new information about this medicine that you should know since you were last treated.

---

### What CMV Immunoglobulin-VF is used for

---

CMV Immunoglobulin-VF is manufactured from human plasma (the liquid component of blood) collected by Australian Red Cross Lifeblood. Donations are selected on the basis that they contain high levels of specific antibody to the cytomegalovirus. CMV Immunoglobulin-VF contains protein substances called antibodies, which can provide protection against cytomegalovirus infection. CMV Immunoglobulin-VF is used to prevent cytomegalovirus infection in

specific transplant patients. It may also help in the treatment of cytomegalovirus infection.

**Ask your doctor if you have any questions why CMV Immunoglobulin-VF has been prescribed for you.**

Your doctor will have assessed the risks and benefits associated with the use of this medicine for you.

---

### Before you are given CMV Immunoglobulin-VF

---

**Make sure you tell your doctor of any reasons you know of why you should not be given this medicine. Inform your doctor of your blood group, if known.**

**You must not be given this medicine if you have:**

- a history of allergy to human immunoglobulin products (allergic reactions may include skin rash, face swelling, wheezing or breathing difficulties) or previously been told you react to any of the ingredients in CMV Immunoglobulin-VF (human immunoglobulins or maltose)
- been told you have antibodies to immunoglobulin A (IgA).

**Tell your doctor if you have allergies to any other medicines or if you have ever had an allergic reaction to an injection.**

**Tell your doctor also if you:**

- have previously been advised that you have Immunoglobulin A (IgA) deficiency
- have previously been advised that you have kidney disease

- have a history of heart, or blood vessel disease, or blood clots, have thick blood, have been immobile for some time. Also tell the doctor what medicine you are using as some medicines, such as those that contain the hormone estrogen (for example, birth control pills), may increase your risk of developing a blood clot.
- have previously been advised that you have diabetes
- have blood group A, B or AB
- are taking or using any other medicines. These include medicines bought from pharmacies, supermarkets and health food stores.
- have any other medical conditions
- are pregnant or breast-feeding
- become pregnant during your treatment
- have had any vaccination during the last two weeks.

**Vaccinations:** Please inform your doctor if you are planning to have a vaccination. CMV Immunoglobulin-VF may impair the effect of some virus vaccines such as measles, mumps, rubella and chicken pox for a period of at least 6 weeks, and up to 3 months. After receiving this medicine, a period of 3 months should be allowed before vaccination with some virus vaccines. In the case of measles vaccine, this effect may last for up to 1 year. Therefore, your vaccinating doctor should check the effectiveness of the measles vaccination.

If you want any further information, consult your doctor.

## About blood products

When products are made from human blood and injected into you, it is possible that viruses or other substances could be present in the product and cause an illness. These could be viruses such as hepatitis, human immunodeficiency virus (HIV), or human parvovirus B19 and theoretically the Creutzfeldt-Jakob Disease (CJD) agent. There could also be other infectious agents, some of which may not yet have been discovered.

To reduce the risk of this happening, extra steps are taken when manufacturing this product. Strict controls are applied when selecting blood donors and donations. The product is specially treated to remove and kill certain viruses. These special treatments are considered effective against viruses known as enveloped viruses such as HIV, hepatitis B virus and hepatitis C virus, and non-enveloped viruses, such as hepatitis A virus and human parvovirus B19. Despite these measures, the risk of viral and other agent's infectivity cannot be totally eliminated.

Vaccines are available against some of these viruses and your doctor will be able to help you decide whether it is worthwhile having any of those vaccines.

Please discuss the risks and benefits of this product with your doctor.

---

## How to use CMV Immunoglobulin-VF

---

Your doctor will determine the dose(s) of CMV Immunoglobulin-VF that you are to receive. Your doctor will give you CMV Immunoglobulin-VF as an infusion, that is, an injection given slowly into the vein.

---

## Side effects

---

Along with their intended effects, blood products occasionally cause unwanted effects, some of which can sometimes be serious.

Individuals may react differently to similar doses of the same medicine. This applies to CMV Immunoglobulin-VF.

Severe allergic reactions to immunoglobulins are rare. Should a severe allergic reaction occur, the doctor will stop the infusion.

Unwanted effects which may occur include: stomach pain, headache, chest tightness, flushed or pale face, feeling hot, shortness of breath, non-itchy skin rash, itching, faintness, dizziness, nausea or vomiting, or a hot feeling or redness at the site of injection. Should any of these effects develop during infusion, the doctor will take appropriate action.

Some patients may develop delayed unwanted effects such as nausea, vomiting, chest pain, chills or shivering, dizziness or aching legs. These effects may occur after the infusion has stopped but usually within 24 hours.

A condition called Aseptic Meningitis Syndrome (AMS) has been reported to occur infrequently in association with infusions similar to CMV Immunoglobulin-VF. It usually begins within several hours to two days following treatment. The signs include severe headache, neck stiffness, drowsiness, fever, inability to stand bright light, painful eye movements, and nausea and vomiting, or a hot feeling or redness at the site of injection. The condition reverses without ill effects when treatment is stopped.

There have been reports that the kidneys may be affected with infusions similar to CMV Immunoglobulin-VF. These occurrences are extremely rare.

**If you experience any of the mentioned effects or any other**

**abnormal signs after treatment, contact your doctor immediately.**

**Contact your doctor immediately if you experience any of these symptoms at any time:**

- fever
- loss of appetite
- extreme tiredness
- dizziness
- paleness of skin
- stomach pain
- jaundice (yellow skin and eyes)
- dark urine
- joint pains
- skin rashes
- shortness of breath
- chest pain
- pain/tenderness, swelling/discolouration of an arm or leg
- weakness or numbness on one side of the body.

## Interference with glucose estimations

The maltose present in CMV Immunoglobulin-VF may interfere with some blood glucose measurements, resulting in the overestimation of blood glucose results. If this glucose measurement is used to guide treatment, hypoglycaemia may occur. Only certain glucose tests have been implicated, so when monitoring glucose levels consult your doctor to ensure that maltose does not interfere with the blood glucose reading of the test you are using.

If you want further information, or if you are worried about any other symptoms after the infusion, consult your doctor.

## Overdose

There is no information available.

---

## Storing CMV Immunoglobulin-VF

---

Store at 2°C to 8°C (Refrigerate. Do not freeze). Protect from light.

Do not use after the expiry date shown on the label.

### *Further information*

CMV Immunoglobulin-VF can only be obtained on a doctor's prescription. This leaflet does not contain the complete information about CMV Immunoglobulin-VF. If you require further information about CMV Immunoglobulin-VF and your treatment generally, or if you have any questions or are not sure about something in this leaflet, consult your doctor.

---

## Product description

---

### *What it looks like*

CMV Immunoglobulin-VF is a clear, colourless, non-viscous (not thick) solution. It is available in glass bottles.

### *Ingredients*

In each vial of CMV Immunoglobulin-VF is a sterile solution containing 6% blood proteins of which at least 98% is immunoglobulins. It also contains 10% maltose (a sugar).

### *Manufacturer*

CMV Immunoglobulin-VF is manufactured in Australia by:

CSL Behring (Australia) Pty Ltd  
ABN 48 160 734 761  
189–209 Camp Road  
Broadmeadows VIC 3047  
Australia

### *Distributor*

Australian Red Cross Lifeblood

### *Date of revision*

June 2021

### **Australian Register Number**

AUST R 31810