

# OLMERTAN COMBI TABLET

OLMERTAN COMBI 20/12.5 Tablet

OLMERTAN COMBI 40/12.5 Tablet

OLMERTAN COMBI 40/25 Tablet

*Olmesartan medoxomil and hydrochlorothiazide*

## Consumer Medicine Information

### What is in this leaflet

This leaflet answers some common questions about OLMERTAN COMBI Tablet.

It does not contain all the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking OLMERTAN COMBI Tablet against the benefits he/she expects it will have for you.

**If you have any concerns about taking this medicine, ask your doctor or pharmacist.**

**Keep this leaflet with your OLMERTAN COMBI Tablets.**

You may need to read it again.

### What OLMERTAN COMBI Tablet is used for

OLMERTAN COMBI Tablet is used to treat high blood pressure which is sometimes called hypertension.

Everyone has blood pressure. This pressure helps push blood all around your body. Your blood pressure changes at different times of the day, depending on how busy or worried you are. You have hypertension (high blood pressure) when your blood pressure stays

higher than is needed, even when you are calm and relaxed.

There are usually no symptoms of hypertension. The only way of knowing that you have hypertension is to have your blood pressure checked on a regular basis. If high blood pressure is not treated it can lead to serious health problems, including stroke, heart disease and kidney failure.

### How OLMERTAN COMBI Tablet works

OLMERTAN COMBI Tablet belongs to a group of medicines known as angiotensin-II receptor antagonists. Angiotensin-II is a substance produced in the body which causes blood vessels to tighten. OLMERTAN COMBI Tablet blocks the action of angiotensin-II and therefore relaxes your blood vessels. This helps lower your blood pressure.

OLMERTAN COMBI Tablet also contains a diuretic which reduces the amount of fluid in the body. This also helps to lower blood pressure.

Your doctor may have prescribed OLMERTAN COMBI Tablet for another reason. Ask your doctor if you have any questions about why OLMERTAN COMBI Tablet has been prescribed for you.

OLMERTAN COMBI Tablet is not addictive.

This medicine is available only with a doctor's prescription.

### Before you take OLMERTAN COMBI Tablet

#### When you must not take it

- **Do not take OLMERTAN COMBI Tablet if you have an allergy to olmesartan, sulfonamide derived medicines (e.g. thiazide diuretics), or any of the ingredients listed at the end of this leaflet**  
Symptoms of an allergic reaction to OLMERTAN COMBI Tablet may include skin rash, itchiness, shortness of breath, swelling of the face, lips or tongue, muscle pain or tenderness or joint pain.
- **you have serious problems with your kidneys**
- **you have serious problems with your liver**
- **you have low potassium or sodium levels in the blood**
- **you have high calcium or uric acid levels in the blood.**
- **you have diabetes and are taking a medicine called aliskiren to reduce blood pressure.**

**Do not take OLMERTAN COMBI Tablet if you are pregnant or breastfeeding.**

OLMERTAN COMBI Tablet may enter your womb or it may pass into the breast milk and there is the possibility that your baby may be affected.

If pregnancy is discovered OLMERTAN COMBI Tablet should be discontinued as soon as possible.

**Do not take OLMERTAN COMBI Tablet after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.**

**If you are not sure whether you should start taking OLMERTAN COMBI Tablet, talk to your doctor.**

*Before you start to take it*

**Tell your doctor if you have allergies to:**

- any other medicines
- any other substances, such as foods, preservatives or dyes.

**Tell your doctor if you have or have had any medical conditions, especially the following:**

- kidney problems
- liver problems
- heart problems
- diabetes
- excessive vomiting or diarrhoea recently
- gout
- Systemic Lupus Erythematosus (SLE)
- high levels of potassium in your blood.
- problems with your adrenal glands (small glands above the kidneys)

**You must also tell your doctor if you:**

- are following a very low salt diet
- you are or intend to become pregnant or plan to breastfeed.
- you are taking potassium supplements, potassium-sparing agents, potassium-containing salt substitutes or other medicines that may increase serum potassium (e.g., trimethoprim-containing products)
- you have skin cancer or if you develop a new skin lesion

during treatment. Treatment with hydrochlorothiazide, particularly long-term use with high doses, may increase the risk of some types of skin and lip cancer (nonmelanoma skin cancer). Discuss with your doctor how to protect your skin from sun exposure, and avoid artificial tanning.

**If you have not told your doctor about any of the above, tell him/her before you start taking OLMERTAN COMBI Tablet.**

*Taking other medicines*

**Tell your doctor or pharmacist if you are taking any other medicines, including any that you buy without a prescription from your pharmacy, supermarket or health food shop.**

Some medicines and OLMERTAN COMBI Tablet may interfere with each other.

These include:

- other medicines used to treat high blood pressure
- other diuretics, also known as fluid or water tablets
- non-steroidal anti-inflammatory drugs (NSAIDs) or COX-2 inhibitors, medicines used to relieve pain, swelling and other symptoms of inflammation, including arthritis
- potassium supplements or potassium-containing salt substitutes
- other medicines that may increase serum potassium (e.g., trimethoprim-containing products)
- lithium or antidepressant medicines
- insulin and tablets used to treat diabetes
- antacids, medicines used to treat heartburn and indigestion
- medicines which lower your immune system, such as corticosteroids and cytotoxic medicines used to treat cancer (including radiation therapy)

- alcohol
- laxatives, medicines used to treat constipation
- medicines used to relieve pain
- medicines used to treat epilepsy
- antiarrhythmics which treat irregular heart beats
- muscle relaxants
- cholestyramine and colestipol, used to treat high cholesterol
- any medicine that contains aliskiren
- any medicines that contain colesevelam.

These medicines may be affected by OLMERTAN COMBI Tablet, or may affect how well it works. You may need different amounts of your medicine, or you may need to take different medicines.

Other medicines not listed above may also interfere with OLMERTAN COMBI Tablet. Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking this medicine.

*Use in children*

The safety and effectiveness of OLMERTAN COMBI Tablet in children has not been established.

---

## How to take OLMERTAN COMBI Tablet

---

**Follow all directions given to you by your doctor or pharmacist carefully.**

They may differ from the information contained in this leaflet.

Your doctor or pharmacist will tell you which OLMERTAN COMBI tablet you will need to take each day. This depends on your condition and whether or not you are taking any other medicines.

**If you do not understand the instructions on the box, ask your doctor or pharmacist for help.**

### *How much to take*

The dose of OLMERTAN COMBI Tablet is one tablet to be taken once a day.

### *How to take it*

**Swallow OLMERTAN COMBI Tablet whole with a full glass of water.**

Do not chew the tablets.

### *When to take it*

**Take OLMERTAN COMBI Tablet at about the same time each day, with or without food.**

Taking your tablets at the same time each day will have the best effect.

It will also help you remember when to take the tablets. It does not matter whether you take it with or without food.

### *How long to take it*

OLMERTAN COMBI Tablet helps control your condition, but does not cure it. Therefore you must take OLMERTAN COMBI Tablet every day. Continue taking your medicine for as long as your doctor tells you.

### *If you forget to take it*

**If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.**

**Otherwise, take it as soon as you remember and then go back to taking your medicine as you would normally.**

**Do not take a double dose to make up for the dose that you missed.**

This may increase the chance of you getting an unwanted side effect.

**If you are not sure what to do, ask your doctor or pharmacist.**

**If you have trouble remembering to take your medicine, ask your pharmacist for some hints.**

### *If you take too much (overdose)*

**Immediately telephone your doctor or the Australian Poisons Information Centre (telephone 13 11 26) or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much OLMERTAN COMBI Tablet. Do this even if there are no signs of discomfort or poisoning.**

**Keep telephone numbers for these places handy.**

If you take too much OLMERTAN COMBI Tablet, you may feel light-headed, dizzy or you may faint. You may also have a fast heart beat.

---

## While you are using OLMERTAN COMBI Tablet

---

### *Things you must do*

**If you are about to be started on any new medicine, tell your doctor, dentist or pharmacist that you are taking OLMERTAN COMBI Tablet.**

**Make sure you drink enough water during exercise and hot weather when you are taking OLMERTAN COMBI Tablet, especially if you sweat a lot.**

If you do not drink enough water while taking OLMERTAN COMBI Tablet, you may feel faint, light-headed or sick. This is because your blood pressure is dropping suddenly. If you continue to feel unwell, tell your doctor.

**If you have excess vomiting and/or diarrhoea, while taking OLMERTAN COMBI Tablet, tell your doctor.**

You may lose too much water and salt and your blood pressure may drop too much.

**If you feel light-headed or dizzy after taking your first dose of OLMERTAN COMBI Tablet, or**

**when your dose is increased, tell your doctor immediately.**

**If you plan to have surgery that needs a general anaesthetic, tell your doctor or dentist that you are taking OLMERTAN COMBI Tablet.**

Your blood pressure may drop suddenly.

**If you become pregnant while taking OLMERTAN COMBI Tablet, tell your doctor immediately.**

**If you are about to have any blood tests, tell your doctor that you are taking OLMERTAN COMBI Tablet.**

OLMERTAN COMBI Tablet may interfere with the results of some tests.

**Tell your doctor if photosensitivity reaction occurs during your treatment.**

**Tell your doctor immediately in case of acute onset of decreased visual acuity or ocular pain.**

**Have your blood pressure checked when your doctor says, to make sure OLMERTAN COMBI Tablet is working.**

**Go to your doctor regularly for a check-up.**

Your doctor may occasionally do a blood test to check your potassium levels and see how your kidneys are working.

### *Things you must not do*

**Do not give OLMERTAN COMBI Tablet to anyone else, even if they have the same condition as you.**

**Do not take OLMERTAN COMBI Tablet to treat any other complaints unless your doctor or pharmacist tells you to.**

**Do not stop OLMERTAN COMBI Tablet, or lower the dosage, without checking with your doctor.**

### *Things to be careful of*

#### **If you feel light-headed, dizzy or faint when getting out of bed or standing up, get up slowly.**

Standing up slowly, especially when you get up from bed or chairs, will help your body get used to the change in position and blood pressure. If this problem continues or gets worse, talk to your doctor.

#### **Be careful driving or operating machinery until you know how OLMERTAN COMBI Tablet affects you.**

As with other medicines in this class, OLMERTAN COMBI Tablet may cause dizziness, light-headedness or tiredness in some people. Make sure you know how you react OLMERTAN COMBI Tablet before you drive a car, operate machinery, or do anything else that could be dangerous if you are dizzy or light-headed. If this occurs do not drive. If you drink alcohol, dizziness or light-headedness may be worse.

### *Things that would be helpful for your blood pressure*

Some self help measures suggested below may help your condition. Talk to your doctor or pharmacist about these measures and for more information.

- Alcohol - your doctor may advise you to limit your alcohol intake.
- Weight - your doctor may suggest losing some weight to help lower your blood pressure and help lessen the amount of work your heart has to do. Some people may need a dietician's help to lose weight.
- Diet - eat a healthy diet which includes plenty of fresh vegetables, fruit, bread (preferably wholegrain), cereals and fish. Also eat less sugar and fat (especially saturated fat) which includes sausages, fatty meats, full cream dairy products, biscuits, cakes, pastries,

chocolates, chips and coconut. Monounsaturated and polyunsaturated fats from olive oil, canola oil, avocado and nuts are beneficial in small quantities.

- Salt - your doctor may advise you to watch the amount of salt in your diet. To reduce your salt intake you should avoid using salt in cooking or at the table and avoid cooked or processed foods containing high sodium (salt) levels.
- Exercise - regular exercise, maintained over the long term, helps to reduce blood pressure and helps get the heart fitter. Regular exercise also improves your blood cholesterol levels, helps reduce your weight and stress levels, and improves your sleep, mood and ability to concentrate. However, it is important not to overdo it. Walking is good exercise, but try to find a route that is reasonably flat. Before starting any exercise, ask your doctor about the best kind of programme for you.
- Smoking - your doctor may advise you to stop smoking or at least cut down. There are enormous benefits to be gained from giving up smoking. There are many professionals, organisations and strategies to help you quit. Ask your doctor or pharmacist for further information and advice.

---

## Side effects

---

#### **Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking OLMERTAN COMBI Tablet.**

OLMERTAN COMBI Tablet helps most people with high blood pressure, but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of

the time they are not. You may need medical treatment if you get some of the side effects.

If you are over 65 years of age you may have an increased chance of getting side effects.

Hydrochlorothiazide, a component of this medicine, increases sensitivity of the skin to the sun and may increase the risk of some types of skin and lip cancer (non-melanoma skin cancer).

#### **Ask your doctor or pharmacist to answer any questions you may have.**

The following is a list of possible side effects. Do not be alarmed by this list. You may not experience any of them.

#### **Tell your doctor or pharmacist if you notice any of the following and they worry you:**

- feeling light-headed, dizzy or faint
- cough
- headache
- feeling sick (nausea) or vomiting
- stomach pain
- diarrhoea or constipation
- an uncomfortable feeling in the stomach
- unusual tiredness or weakness, fatigue
- 'flu-like' symptoms
- runny or blocked nose, or sneezing
- bronchitis
- sore throat and discomfort when swallowing (pharyngitis)
- swelling of the hands, feet or ankles
- back pain
- urinary tract infection
- jaundice
- blurred vision
- skin rashes or eczema
- sleep disturbance
- depression.

These are common side effects. They are generally mild.

**Tell your doctor as soon as possible if you notice any of the following:**

- skin rash or itchiness
- aching, tender or weak muscles not caused by exercise
- painful joints
- fast heart beat
- shortness of breath or tightness in the chest
- swelling of the hands, feet or ankles
- symptoms that may indicate high potassium levels in the blood, such as nausea, diarrhoea, muscle weakness, change in heart rhythm.

These may be serious side effects. You may need medical attention. Serious side effects are rare.

**If any of the following happen, stop taking OLMERTAN COMBI Tablet and either tell your doctor immediately or go to Accident and Emergency at your nearest hospital:**

- swelling of the face, lips, mouth, tongue or throat which may cause difficulty in swallowing or breathing
- chest pain.

These are very serious side effects. You may need urgent medical attention or hospitalisation. These side effects are very rare.

**Other side effects not listed above may occur in some patients. Tell your doctor or pharmacist if you notice anything that is making you feel unwell.**

---

## After using OLMERTAN COMBI Tablet

---

### Storage

**Keep your tablets in the box until it is time to take them.**

If you take the tablets out of the box they will not keep well.

**Keep your tablets in a cool dry place where the temperature stays below 25°C.**

**Do not store OLMERTAN COMBI Tablet or any other medicine in the bathroom or near a sink.**

**Do not leave it on a windowsill or in the car on hot days.**

Heat and dampness can destroy some medicines.

**Keep it where children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

### Disposal

**If your doctor tells you to stop taking OLMERTAN COMBI Tablet or the tablets have passed their expiry date, ask your pharmacist what to do with any that are left over.**

---

## Product description

---

### What it looks like

OLMERTAN COMBI Tablet tablets come in three strengths and each has a different appearance:

OLMERTAN COMBI 20/12.5 Tablet is a reddish-yellow, round, film coated tablets debossed with "A01" on one side & plain on other side.

OLMERTAN COMBI 40/12.5 Tablet is a reddish-yellow, oval, film coated tablets debossed with "A002" on one side & plain on other side.

OLMERTAN COMBI 40/25 Tablet is a pink, oval, film coated tablets debossed with "A003" on one side & plain on other side.

The product is presented in Al/Al blister packs of 10 or 30 film-coated tablets.

### Ingredients

Active ingredients:

OLMERTAN COMBI 20/12.5 Tablet- 20mg olmesartan medoxomil and 12.5mg hydrochlorothiazide

OLMERTAN COMBI 40/12.5 Tablet - 40mg olmesartan medoxomil and 12.5mg hydrochlorothiazide

OLMERTAN COMBI 40/25 Tablet - 40mg olmesartan medoxomil and 25mg hydrochlorothiazide

Other ingredients:

- Lactose monohydrate,
- Microcrystalline cellulose,
- Hydroxypropyl cellulose,
- Magnesium stearate,
- Stearic acid
- Instacoat Universal A05G11632 yellow (ARTG No. 110035) [for 20/12.5 mg and 40/12.5 mg strengths only] that contains titanium dioxide (E171), talc (E553b), hypromellose (E464), polyethylene glycol (E1521), Iron oxide red (E172) and Iron oxide yellow (E172). [for 20mg/12.5mg and 40mg/12.5mg]
- Instacoat Universal A05G11616 orange (ARTG No. 110036) [for 40/25 mg strength only] that contains titanium dioxide (E171), talc (E553b), hypromellose (E464), polyethylene glycol (E1521), Iron oxide red (E172) and Iron oxide yellow (E172). [for 40mg/25mg]

OLMERTAN COMBI Tablet does not contain sucrose, tartrazine or any other azo dyes.

### Supplier

OLMERTAN COMBI Tablet is supplied in Australia by:

**PHARMACOR PTY. LTD  
CHATSWOOD, NSW, 2067,  
Australia**

*Australian Registration  
Numbers*

OLMERTAN COMBI Tablet  
20/12.5 Tablets blister pack –  
286101

OLMERTAN COMBI Tablet  
40/12.5 Tablets blister pack –  
286095

OLMERTAN COMBI Tablet 40/25  
Tablets blister pack - 286100

This leaflet was prepared in  
05/2021.