

FAMCICLOVIR GENERICHEALTH

For genital herpes

(famciclovir) film-coated tablets

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Famciclovir generichealth. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Famciclovir generichealth against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What Famciclovir generichealth is used for

Famciclovir generichealth is used to treat outbreaks of genital herpes and also to suppress (prevent) recurrent outbreaks of the condition.

Genital herpes is a viral infection caused by herpes simplex 1 or herpes simplex 2. It is usually transmitted through sexual contact.

Symptoms include tingling, burning or itching of the genitals, followed by blisters that may be painful.

People who have frequent episodes of genital herpes can also take Famciclovir generichealth to help prevent the attacks.

Although Famciclovir generichealth does not cure the viral infection, it

helps to relieve the symptoms and shorten their duration.

The best results are obtained if the medicine is started as soon as possible after the first symptoms begin to appear.

Taking Famciclovir generichealth does not prevent you from spreading the herpes virus to another person.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

Your doctor may have prescribed it for another reason.

This medicine is not addictive.

This medicine is available only with a doctor's prescription.

Famciclovir generichealth for treating genital herpes is not recommended for use in children under 12 years of age.

Before you take Famciclovir generichealth

When you must not take it

Do not take Famciclovir generichealth if you have an allergy to:

- any medicine containing famciclovir (active ingredient);
- penciclovir, a related antiviral medicine;
- any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include:

- shortness of breath;
- wheezing or difficulty breathing;
- swelling of the face, lips, tongue or other parts of the body;
- rash, itching or hives on the skin.

If you experience an allergic reaction, stop using the medicine and inform your doctor or pharmacist immediately.

Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start taking this medicine, talk to your doctor.

Before you start to take it

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you are prone to allergies.

Tell your doctor if you have or have had any of the following medical conditions:

- you have a problem with your body's immune system, which helps to fight off infections;
- you have a problem with your kidneys or liver.

Your doctor may want to take extra precautions in that case.

Tell your doctor if you are pregnant or plan to become pregnant or are breastfeeding.

Famciclovir generichealth should not be used during pregnancy unless

necessary. Your doctor will discuss with you the potential risks of taking Famciclovir generichealth during pregnancy, and will also advise you if you should take Famciclovir generichealth while breastfeeding, based on the benefits and risks of your particular situation.

If you have not told your doctor about any of the above, tell them before you start taking Famciclovir generichealth.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Famciclovir generichealth may interfere with each other. These include:

- probenecid, a prescription medicine used to treat gout (a disease with painful, swollen joints, caused by uric acid crystals) and to increase blood levels of penicillin-type antibiotics;
- raloxifene, a medicine used to treat osteoporosis (a disease which causes bones to become less dense, gradually making them weaker, more brittle and likely to break);
- medicines that can affect your kidneys;
- medicines that can affect your immune system.

These medicines may be affected by Famciclovir generichealth or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking this medicine.

How to take Famciclovir generichealth

Follow all directions given to you by your doctor or pharmacist carefully.

They may differ from the information contained in this leaflet.

If you do not understand the instructions on the box, ask your doctor or pharmacist for help.

Do not change the dose yourself without your doctor's advice, regardless of how well you may feel.

How much to take

Follow your doctor's instructions on how many Famciclovir generichealth tablets to take.

For people whose immune system does not work as well as it should, the dose and duration of treatment may be increased.

For people who have reduced kidney function, the dose may be reduced.

Ask your pharmacist or doctor if you are unsure of the correct dose for you.

They will tell you exactly how much to take.

How to take it

Swallow the tablets whole with a full glass of water.

They may be taken with or without food.

Try to take the tablets at about the same time each day, as directed by your doctor or pharmacist.

Taking your tablet at the same time each day will have the best effect. It will also help you remember when to take the tablets.

When to take it

1 TO TREAT AN OUTBREAK OF GENITAL HERPES

To treat an outbreak, take the tablets as soon as possible after

the first symptoms of genital herpes appear.

The tablets are best taken within 6 hours of the first symptoms of genital herpes appearing.

There are three ways to take Famciclovir generichealth to treat an outbreak of genital herpes and your doctor will tell you which regimen is best for you:

- Two 500 mg Famciclovir generichealth tablets twice daily for one day;

Or

- Two 250 mg Famciclovir generichealth tablets to start with, followed by one 250 mg tablet every 12 hours for the next 3 doses;

Or

- One 125 mg Famciclovir generichealth tablet every 12 hours for 5 days.

Take the second dose (and subsequent doses, if applicable) 12 hours after the first dose, or as close as possible to 12 hours during waking hours.

If you take the first dose in the late morning or early afternoon, you can take the next dose before going to bed, but do not take any doses less than 6 hours apart.

During normal waking hours, take any remaining doses at 12 hourly intervals.

Your doctor may have prescribed a different dose.

2 TO SUPPRESS (PREVENT) OUTBREAKS OF RECURRENT GENITAL HERPES

Start suppressive treatment to prevent outbreaks of recurrent genital herpes as soon as possible after you have your Famciclovir generichealth prescription filled.

Take one 250 mg Famciclovir generichealth tablet twice each day.

Continue to take one 250 mg Famciclovir generichealth tablet twice each day for as long as your doctor tells you to. Do this even if you do not have an outbreak.

This medicine helps to control your condition but does not cure it. Your doctor will tell you when you can stop.

Fill your next repeat prescription before using all of the tablets in your current carton. This will ensure that your treatment can be continued and give you the best results.

How long to take it

Continue taking Famciclovir generichealth every day for as long as your doctor tells you.

To help clear up your infection, you must keep taking this medicine, even if your symptoms begin to clear up after a few days.

To prevent recurrent episodes of genital herpes, you must take the tablets each day, even if you have no symptoms.

It is important to keep taking your medicine even if you feel well.

If you forget to take it

Take a dose as soon as you remember. Take your next tablet at the usual time, and then go back to taking it as you would normally.

Do not take two doses within a time frame of less than one hour. In that case, skip the missed dose.

Do not take a double dose to make up for the one that you missed.

This may increase the chance of you getting an unwanted side effect.

If you have trouble remembering when to take your medicine, ask your pharmacist for some hints.

If you take too much (overdose)

Immediately telephone your doctor or the Poisons Information Centre (telephone 13 11 26) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much Famciclovir generichealth.

Do this even if there are no signs of discomfort or poisoning. Keep the telephone numbers for these places handy.

You may need urgent medical attention.

Taking too much Famciclovir generichealth may affect the kidneys. In people who already have kidney problems it may, rarely, lead to kidney failure if their dose is not correctly lowered.

While you are taking Famciclovir generichealth

Things you must do

If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking Famciclovir generichealth.

Tell any other doctors, dentists and pharmacists who treat you that you are taking Famciclovir generichealth.

If you become pregnant while taking Famciclovir generichealth, tell your doctor.

Your doctor can discuss with you the risks of taking it while you are pregnant.

Tell your pharmacist or doctor if your symptoms do not improve within a few days, or if they become worse.

Things you must not do

Do not take Famciclovir generichealth to treat any other complaints unless your doctor tells you to.

Do not give your medicine to anyone else, even if they have the same condition as you.

Do not stop taking your tablets or change the dosage without checking with your doctor first.

If you stop your tablets suddenly, your condition may worsen or you may have unwanted side effects.

Things to be careful of

If you are pregnant or breastfeeding, ask your doctor or pharmacist for advice before taking any medicine.

Be careful driving, operating machinery or doing jobs that require you to be alert until you know how Famciclovir generichealth affects you.

This medicine can cause dizziness, sleepiness or confusion.

Practice "safer sex", including the use of condoms, when symptoms are present, even if you have started taking Famciclovir generichealth for genital herpes. This is important to prevent you passing the infection on to others.

Things that may help your condition

Take the following precautions to avoid spreading the virus:

- Use condoms between episodes to reduce the risk of infecting your partner.
- Keep the areas affected by the virus as clean and dry as possible.
- Wear loose-fitting clothing to avoid irritating the blisters.
- Avoid touching or scratching the sore area as you may spread the virus on your fingers.

This is important to prevent you passing the infection on to others.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Famciclovir generichealth.

This medicine helps most people with genital herpes, but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Do not be alarmed by the following lists of side effects. You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- headache;
- dizziness;
- tiredness, sleepiness;
- nausea (feeling sick) or vomiting;
- diarrhoea;
- stomach pain;
- itching or an itchy rash (urticaria);
- abnormal liver function test results.

The above list includes the more common side effects of your medicine. They are usually mild and short-lived.

Tell your doctor as soon as possible if you notice any of the following:

- a rash elsewhere on the body;
- extreme sleepiness or confusion, usually in older people;
- hallucinations (seeing or hearing things that are not really there);
- painful or swollen joints;
- aching muscles or muscle tenderness or weakness that is not caused by exercise;
- palpitations (signs of abnormal heart beat).

The above list includes serious side effects that may require medical attention. Serious side effects are rare.

If any of the following happen, tell your doctor immediately or go to Accident and Emergency at your nearest hospital:

- swelling below the surface of the skin (eg. swelling around the face, eye, eyelid or throat);
- bruising or bleeding more easily than usual (may indicate that the number of platelets (a type of blood cell responsible for blood

clotting) in your blood is reduced);

- widespread red, raised areas, blisters or sores of the lips, eyes, mouth, nasal passages or genitals (signs of a serious skin reaction);
- purple patches, itching, burning of the skin (signs of inflamed blood vessels);
- persistent pain in the upper right abdomen, yellowing of the skin and/or eyes, dark urine or pale bowel motions (signs of a possible liver problem);
- seizures or fits;
- difficulty breathing or swallowing, wheezing or coughing, light-headedness, changes in alertness, skin reddening, facial/throat swelling, blue discoloration of the lips, tongue or skin (signs of severe allergic reaction).

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation. These side effects are very rare.

Tell your doctor or pharmacist if you notice anything that is making you feel unwell.

Other side effects not listed above may also occur in some people.

After taking Famciclovir generichealth

Storage

Keep your tablets in the pack until it is time to take them.

If you take the tablets out of the pack they may not keep well.

Keep your tablets in a cool dry place where the temperature stays below 25°C.

Do not store Famciclovir generichealth or any other medicine in the bathroom or near a sink. Do not leave it on a window sill or in the car.

Heat and dampness can destroy some medicines.

Keep it where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

Product description

What it looks like

Famciclovir generichealth 125 mg tablets are white, round film coated tablets with "FM" on one side and "125" on the other side.

Famciclovir generichealth 250 mg tablets are white, round film coated tablets, marked with "FM" on one side and "250" on the other side.

Famciclovir generichealth 500 mg tablets are white, oval, film coated tablets with "FM" on one side and "500" on the other side.

Ingredients

Active ingredient

Famciclovir generichealth 125 contain 125 mg famciclovir per tablet.

Famciclovir generichealth 250 contain 250 mg famciclovir per tablet.

Famciclovir generichealth 500 contain 500 mg famciclovir per tablet.

Other ingredients

- microcrystalline cellulose;
- crospovidone;
- colloidal anhydrous silica;
- copovidone;
- sodium stearyl fumarate;
- Opadry White YS-22-18096.

Australian Registration Numbers

Famciclovir generichealth 125 mg:
AUST R 177010

Famciclovir generichealth 250 mg:
AUST R 177012

Famciclovir generichealth 500 mg:
AUST R 177014

Distributor

Famciclovir generichealth is
distributed in Australia by:

Generic Health Pty Ltd
Suite 2, Level 2
19-23 Prospect Street
Box Hill, VIC, 3128
Australia

Email:
ghinfo@generichealth.com.au
Telephone: +61 3 9809 7900
Website:
www.generichealth.com.au

This leaflet was prepared in March
2020.