

CLINDAMYCIN LU capsules

Clindamycin hydrochloride

Consumer Medicine Information

WHAT IS IN THIS LEAFLET

This leaflet answers some common questions about CLINDAMYCIN LU capsules. It does not contain all the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking CLINDAMYCIN LU capsules against the benefits this medicine is expected to have for you.

If you have any concerns about using this medicine, ask your doctor or pharmacist. Keep this leaflet with your medicine.

You may need to read it again.

WHAT CLINDAMYCIN LU CAPSULES ARE USED FOR

CLINDAMYCIN LU capsule is an antibiotic. It is used to treat infections in different parts of the body caused by bacteria.

CLINDAMYCIN LU capsules works by killing or stopping the growth of the bacteria causing your infection.

CLINDAMYCIN LU capsules will not work against viral infections such as colds or flu.

CLINDAMYCIN LU capsules are recommended for patients who are allergic to penicillin or patients for whom penicillin is not suitable.

Your doctor may have prescribed CLINDAMYCIN LU capsules for

another reason. Ask your doctor if you have any questions about why CLINDAMYCIN LU capsules have been prescribed for you.

This medicine is available only with a doctor's prescription. CLINDAMYCIN LU capsule is not addictive.

BEFORE TAKING CLINDAMYCIN LU CAPSULES

When you must not take CLINDAMYCIN LU capsulesDo not take CLINDAMYCIN LU capsules:

1. if you have an allergy to:
 - clindamycin or lincomycin
 - lactose or any of the other ingredients listed at the end of this leafletSymptoms of an allergic reaction may include swelling of the face, lips, tongue or other parts of the body, skin rash, itching or wheezing or difficulty in breathing.
2. if the packaging is torn or shows signs of tampering
3. after the expiry date (EXP) printed on the label.

If you take it after the expiry date, it may have no effect at all, or worse, an entirely unexpected effect.

If you are not sure whether you should start taking CLINDAMYCIN LU capsules, talk to your doctor.

Before you start to take CLINDAMYCIN LU capsules

You must tell your doctor about all of the following before you start to take CLINDAMYCIN LU capsules:

1. if you have any allergies to any other medicines or any other substances such as foods, preservatives or dyes.
2. if you are pregnant or intend to become pregnant.

Clindamycin crosses the placenta therefore CLINDAMYCIN LU capsules should only be used in pregnancy if clearly needed. Your doctor will discuss the risks and benefits of taking CLINDAMYCIN LU capsules during pregnancy.
3. if you are breastfeeding or plan to breastfeed.

CLINDAMYCIN LU capsules are not recommended during breastfeeding.
4. if you have or have ever had:
 - severe diarrhoea associated with the use of antibiotics
 - severe liver disease
 - severe kidney disease
 - bowel disease, a medicine used to treat bacterial infections
 - any gastrointestinal (stomach or gut) problems
5. if you have had any other health problems or medical conditions
6. if you are taking erythromycin, a medicine used to treat bacterial infections

If you have not told your doctor or pharmacist about any of the above, do so before you start taking CLINDAMYCIN LU capsules.

Taking other medicines

Tell your doctor if you are taking any other medicines, including medicines you buy without a prescription from a pharmacy, supermarket or health food shop.

There may be interference between CLINDAMYCIN LU capsules and some other medicines including:

- the antibiotic, erythromycin
- rifampicin, a medicine used to treat bacterial infections
- medicines used for muscle relaxation in anaesthesia

These medicines may be affected by CLINDAMYCIN LU capsules or may affect how well

CLINDAMYCIN LU capsules work. You may need different amounts of your medicine or you may need to take different medicines. Your doctor or pharmacist may have more information on medicines to be careful with or avoid while taking CLINDAMYCIN LU capsules.

HOW TO TAKE CLINDAMYCIN LU CAPSULES

Your doctor or pharmacist will tell you how to take your CLINDAMYCIN LU capsules. Follow all directions given to you by your doctor and pharmacist carefully.

Their directions may differ from the information contained in this leaflet. You may be given a different dosage depending on your condition and how you react to the medicine. The directions your doctor or pharmacist gives you should be strictly followed.

If you do not understand the instructions in this leaflet, ask your doctor or pharmacist for help.

How much to take and how long to take it

Adults: One (1) capsule (150 mg) every six hours is the usual dose. The number of capsules may increase with more serious infections. Your doctor will tell you how long to take your capsules.

Children: CLINDAMYCIN LU capsules are not recommended in children for formulation reasons.

Continue taking CLINDAMYCIN LU capsules until you finish the box or until your doctor recommends. Check with your doctor if you are not sure how long you should be taking it.

Do not stop taking CLINDAMYCIN LU capsules or change the dose because you are feeling better. If you stop taking CLINDAMYCIN LU suddenly, the infection may not clear completely or your symptoms may return.

If you do not complete the full course prescribed by your doctor, all of the bacteria causing your infection may not be killed. These bacteria may continue to grow and multiply so that your infection may not clear completely or may return.

How to take it and when to take it

CLINDAMYCIN LU capsules should be taken by mouth, with a full glass of water.

Space the doses of CLINDAMYCIN LU capsules evenly apart and take them at about the same time each day. This will allow the medicine to have its best effect. It will also help you remember when to take it.

CLINDAMYCIN LU capsules can be taken with or without food, it does not matter.

If you forget to take it

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are

meant to. Otherwise, take it as soon as you remember, then go back to taking your capsules as you would normally.

If you are not sure whether to skip the dose, talk to your doctor or pharmacist.

Do not double a dose to make up for the dose you have missed.

If you take too much (overdose)

Immediately telephone your doctor or Poisons Information Centre (telephone 13 11 26) for advice, or go to Accident and Emergency (Casualty) at your nearest hospital if you think that you or anyone else may have taken too much CLINDAMYCIN LU.

Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Keep telephone numbers for these places/services handy. Have the CLINDAMYCIN LU box or this leaflet available to give details if needed.

WHILE YOU ARE TAKING CLINDAMYCIN LU CAPSULES

Things you must do

If the symptoms of your infection do not improve within a few days, or if they become worse, tell your doctor.

If you get severe diarrhoea, tell your doctor, pharmacist or nurse immediately. Do this even if it occurs several weeks after you have stopped taking CLINDAMYCIN LU capsules. Diarrhoea may mean that you have a serious condition affecting your bowel. You may need urgent medical care.

Do not take any medicines for diarrhoea without first checking with your doctor.

If you get a sore, white mouth or tongue while taking or soon after stopping CLINDAMYCIN LU capsules, tell your doctor. Also tell your doctor if you get vaginal itching or discharge. This may mean you have a fungal/yeast infection called thrush. Sometimes the use of CLINDAMYCIN LU allows fungi/yeast to grow, and the above symptoms to occur. CLINDAMYCIN LU does not work against fungi/yeast.

If you become pregnant while taking CLINDAMYCIN LU, tell your doctor immediately.

If you are about to start taking any new medicines, tell your doctor and pharmacist that you are taking CLINDAMYCIN LU.

Tell all doctors, dentists and pharmacists who are treating you that you are taking CLINDAMYCIN LU.

If you feel that CLINDAMYCIN LU is not helping your condition, tell your doctor.

Tell your doctor if, for any reason, you have not used CLINDAMYCIN LU exactly as prescribed.

If you get a severe skin rash tell your doctor immediately. Do this even if the rash occurs after CLINDAMYCIN LU has been stopped. A severe skin rash may mean you are having an allergic reaction to CLINDAMYCIN LU. You may need urgent medical care.

If you are about to have any blood tests, tell your doctor that you are taking this medicine. It may interfere with the results of some tests.

Keep all of your doctor's appointments so that your progress can be checked. Your doctor may do some tests from time to time to make sure the medicine is working and to prevent unwanted side effects.

Things you must not do

Do not give CLINDAMYCIN LU to anyone else, even if they have the same condition as you.

Do not use CLINDAMYCIN LU to treat any other medical complaints unless your doctor tells you to.

SIDE EFFECTS

Check with your doctor or pharmacist as soon as possible if you have any problems while taking CLINDAMYCIN LU, even if you do not think the problems are connected with the medicine or are not listed in this leaflet.

Like other medicines, CLINDAMYCIN LU can cause some side effects. If they occur, most are likely to be minor and temporary. However, some may be serious and need medical attention.

Ask your doctor or pharmacist to answer any questions you may have. Tell your doctor if you notice any of the following and they worry you:

- oral thrush - white, furry, sore tongue and mouth
- vaginal thrush - sore and itchy vagina and/or discharge
- stomach cramping
- stomach discomfort
- inflammation of the food pipe; discomfort or/and pain of the food pipe
- loss or distorted sense of taste
- nausea and/or vomiting
- heartburn
- diarrhoea
- loss of appetite
- skin rash; severe irritation of the skin
- jaundice (yellowing of the skin)
- joint pain and swelling

Tell your doctor immediately if you notice any of the following side effects while you are taking CLINDAMYCIN LU capsules or

several weeks after stopping treatment:

- severe stomach cramps;
- watery and severe diarrhoea which may also be bloody;
- fever in combination with one or both of the above.

These are rare, but serious side effects. You may need urgent medical attention.

Do not take any medicine for diarrhoea without first checking with your doctor.

CLINDAMYCIN LU can cause some bacteria, which are normally present in the bowel and normally harmless to multiply and therefore cause the above symptoms. You may need urgent medical attention. However this side effect is rare.

Other side effects not listed above may also occur in some patients. Tell your doctor if you notice anything else that is making you feel unwell.

Some of these side effects (for example, abnormal blood test results and certain kidney and liver conditions) can only be found when your doctor does tests from time to time to check on your progress.

Tell your doctor immediately or go to Emergency at your nearest hospital, if you notice any of the following:

- sudden signs of allergy such as rash, itching or hives on the skin, swelling of the face, lips, tongue or other parts of the body, shortness of breath, wheezing or trouble breathing
- moderate or severe skin rash or blisters often with flu-like symptoms
- enlarged lymph glands and/or fever
- diarrhoea, usually with blood and mucus, stomach pain and fever
- yellowing of the eyes or skin, also called jaundice

- swelling of the legs, ankles, or feet, passing less urine, irregular heartbeat, confusion, shortness of breath, weakness, fatigue and nausea
- chest pain, and/or shortness of breath.

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation.

Do not be alarmed by this list of possible side effects. You may not experience any of them.

AFTER USING CLINDAMYCIN LU CAPSULES

Storage

Keep your CLINDAMYCIN LU capsules in their original container where the temperature stays below 30°C and protected from light, heat and moisture. If you take the capsules out of their original container, they may not keep as well.

Do not store your CLINDAMYCIN LU capsules, or any other medicine, in a bathroom or near a sink. Do not leave them in the car or on window sills.

Heat and dampness can destroy some medicines.

Keep your CLINDAMYCIN LU capsules where children cannot reach them. A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop using CLINDAMYCIN LU capsules or the capsules have passed their expiry date, ask your pharmacist what to do with any capsules left over.

PRODUCT DESCRIPTION

What it looks like

CLINDAMYCIN LU capsules with a strength of 150 mg are available in packs of 24 capsules, 50 capsules and 100 capsules. The No. 2 Apple Green capsules are marked with "CLD 150".

Ingredients

The active ingredient in CLINDAMYCIN LU is clindamycin hydrochloride. CLINDAMYCIN LU also contains Microcrystalline Cellulose, Lactose, Povidone, Sodium Starch Glycolate, Talc and Magnesium Stearate.

The capsule is a Hard Gelatin Capsules size 2 P Lt Green Op / Plt Green Op (proprietary ingredient number: 109543) imprinted with the text CLD 150 using the ink TekPrint SB-0007P White Ink (proprietary ingredient number: 2216).

Supplier

CLINDAMYCIN LU capsules are supplied in Australia by:

Luminarie Pty Ltd
Baulkham Hills
NSW 2153
Australia
<http://www.luminarie.com.au>

Australian Register Number(s)

150 mg Capsules
AUST R 214526

This leaflet was prepared in January 2024