

Idacio 40mg solution for injection in prefilled syringe

Adalimumab (rch)

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Idacio.

It does not contain all the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you using this medicine against the benefits they expect it will have for you.

If you have any concerns about using this medicine, ask your doctor or pharmacist.

Read this leaflet carefully before you use Idacio and keep it with the medicine.

You may need to read it again.

What Idacio is used for

The active ingredient in this medicine is adalimumab, which is a fully human monoclonal antibody. Adalimumab recognises and binds to a specific protein (tumour necrosis factor or TNF-alpha), which is present at higher levels in some inflammatory diseases.

Idacio is intended for the treatment of a number of inflammatory diseases:

- **Rheumatoid arthritis**
Rheumatoid arthritis is an inflammatory disease of the joints. Signs and symptoms of rheumatoid arthritis include joint pain, tenderness, swelling and stiffness.
Idacio is used to reduce the signs and symptoms of moderate to severely active rheumatoid arthritis, as well as to slow down and protect the joints from further damage to help them move more freely. Your doctor will decide if Idacio should be used with another medicine called methotrexate, or on its own.
Idacio can also be used to treat severe, active and progressive rheumatoid arthritis without previous methotrexate treatment.
 - **Polyarticular Juvenile Idiopathic Arthritis (pJIA)**
pJIA is an inflammatory disease of the joints. Idacio is used to reduce the signs and symptoms of moderate to severely active pJIA, in patients 2 years of age and older, when other medicines are not appropriate.
Your doctor will decide whether Idacio should be used with another medicine called methotrexate or used alone.
 - **Enthesitis-related arthritis (ERA)**
ERA is an inflammatory disease of the joints and the places where tendons join the bone. Idacio is used to treat ERA in children.
You may have already been given other medicines to treat your condition. Your doctor has prescribed Idacio for you as you haven't responded well enough to these medicines.
 - **Psoriatic arthritis (PsA)**
PsA is an inflammatory disease of the joints that is usually associated with psoriasis. Signs and symptoms include joint pain, tenderness and swelling.
Idacio is used to reduce the signs and
- symptoms, of moderate to severely active PsA, as well as to slow down and protect the joints from further damage, to help them move more freely.
You may have already been given other medicines to treat your condition. Your doctor has prescribed Idacio for you as you haven't responded well enough to these medicines.
- **Ankylosing spondylitis**
Ankylosing spondylitis is an inflammatory disease of the spine. Signs and symptoms of ankylosing spondylitis include back pain and stiffness. Idacio is used to reduce the signs and symptoms in patients with active disease.
 - **Crohn's Disease**
Crohn's disease is an inflammatory disease of the digestive tract. Idacio is used to treat moderate to severe Crohn's disease, in adults and children aged 6 years and over, to reduce the signs and symptoms of the disease and to induce and maintain periods where the symptoms are no longer present (remission).
You may have already been given other medicines to treat your condition. Your doctor has prescribed Idacio for you as you may either have not responded well enough, or you may have lost response or cannot tolerate these medicines.
 - **Ulcerative Colitis**
Ulcerative colitis is an inflammatory disease of the large intestine (bowel). Idacio is used to treat moderate to severe ulcerative colitis when other medicines are not appropriate.
 - **Psoriasis**
Psoriasis is an inflammatory disease of the skin. Plaque psoriasis, the most common form, is a skin condition that causes red, flaky, crusty patches of the skin covered with silvery scales. Plaque psoriasis can also affect nails, causing them to crumble, thicken and lift away from the nail bed which can be painful. Idacio is used to treat moderate to severe forms of the disease in adults and severe forms in adolescents and children from 4 years of age for whom topical therapy (such as creams, lotions and ointments) and phototherapy (also known as light therapy) have either not worked very well or are not suitable.
 - **Hidradenitis suppurativa (HS)**
HS (sometimes called acne inversa) is a chronic and often painful inflammatory skin disease. Symptoms may include tender nodules (lumps) and abscesses (boils) that may leak pus, which can have an unpleasant odour. It most commonly affects specific areas of the skin, such as under the breasts, the armpits, inner thighs, groin and buttocks. Scarring may also occur in affected areas.
Idacio is used for the treatment of adult and adolescents from 12 years of age with active moderate to severe HS. Idacio can reduce the number of nodules and abscesses caused by the disease, and the pain that is often associated with it.
You may have already been given other medicines to treat your condition. Your

doctor has prescribed Idacio for you as you haven't responded well enough to these medicines.

Your doctor will schedule follow-up appointments to check on your progress to determine whether you should continue treatment.

- **Uveitis**
Uveitis is an inflammatory disease affecting certain parts of the eye. This inflammation may lead to a decrease of vision and/or the presence of floaters in the eye (black dots or wispy lines that move across the field of vision). Idacio is used to treat non-infectious intermediate, posterior and pan-uveitis.
You may have already been given other medicines to treat your condition. Your doctor has prescribed Idacio for you as you may either have not responded well enough, or you have lost response or cannot tolerate these medicines.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

Your doctor may have prescribed it for another reason.

This medicine is not addictive.

This medicine is only available with a doctor's prescription.

The long-term effects of Idacio on the growth and development of children is not known.

Before you use Idacio

When you must not use it

Do not use Idacio if you have an allergy to any medicine containing adalimumab or any of the ingredients listed at the end of this leaflet.

Symptoms of an allergic reaction may include:

- chest tightness
- shortness of breath, wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- hives, itching or skin rash.

Do not use Idacio if you have a severe infection including an infection of the blood (sepsis), active tuberculosis or other severe infections that can be caused by viruses, fungi, parasites or bacteria.

Infections can occur when the body's natural defences are lowered.

Do not use Idacio if you are already using anakinra (Kineret).

Anakinra is a medicine for rheumatoid arthritis, JIA and conditions associated with a defect in a protein called cryoprin.

Do not use Idacio if you have moderate to severe heart failure.

If you are not sure whether any of the above conditions apply to you, ask your doctor.

Do not use this medicine after the expiry date printed on the label / blister / carton or if the packaging is torn or shows signs of tampering.

Return out of date or damaged medicines to your pharmacist for disposal.

Before you use it

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if:

- you have or have had an infection, including a long-term infection or an infection in one part of the body (for example, leg ulcer).
- you have had infections which keep coming back or other conditions that increase the risk of infections. If you are over 65, you may be more likely to get an infection while taking Idacio. It is important that you and your doctor pay special attention to signs of infection while you are being treated with Idacio.
- you have ever had tuberculosis, or if you have been in close contact with someone who has had tuberculosis. As cases of tuberculosis have been reported in patients treated with Idacio, your doctor will check you for signs and symptoms of tuberculosis before starting Idacio. This will include a thorough medical evaluation, including your medical history, and appropriate screening tests (for example, a chest x-ray and tuberculin test). Tuberculosis can develop during therapy even if you have received treatment for the prevention of tuberculosis. If symptoms of tuberculosis (for example a cough that doesn't go away, weight loss, lack of energy, mild fever), or any other infections appear during or after therapy, tell your doctor immediately.
- you are a carrier of the hepatitis B virus (HBV), or you have active HBV or you think you might be at risk of contracting HBV. In people who carry HBV, Idacio can cause the virus to become active again. In some rare cases, especially if you are taking other medicines that suppress the immune system, reactivation of HBV can be life threatening.
- you have or have had a fungal infection or have lived or travelled in countries where some fungal infections are common. These infections may develop or become more severe if you take Idacio.
- you have or have had uveitis, your doctor may check for signs and symptoms of neurologic disease before starting this medicine.
- you have or develop a demyelinating disease (a disease that affects the insulating layer around the nerves, such as multiple sclerosis).
- you have or have had allergic reactions such as chest tightness, wheezing, dizziness, swelling or rash.
- you have or have had a blood disorder.
- you have or have had low resistance to disease.
- you have or have had a serious heart condition.
- you have or have had cancer or autoimmune disease.
- you have a lung disease called chronic obstructive pulmonary disease (COPD).

- you have or have had kidney or liver problems.

Tell your doctor if you are scheduled for any vaccines.

Certain vaccines may cause infections and should not be given while patients are receiving Idacio.

Wherever possible, it is recommended that children be brought up to date with all immunisations according to current immunisation guidelines prior to starting Idacio therapy. Patients receiving Idacio should not receive live vaccines.

Tell your doctor if you have psoriasis and have undergone phototherapy, also known as light therapy.

Tell your doctor if you are pregnant or plan to become pregnant.

You should consider the use of adequate contraception to prevent pregnancy and continue its use for at least 5 months after the last Idacio treatment.

Idacio should only be used during pregnancy if clearly needed.

A pregnancy study found that there was no higher risk of birth defects when the mother had used Idacio during pregnancy, compared with mothers with the same disease who did not use Idacio.

If you use Idacio during pregnancy, your baby may have a higher risk of getting an infection.

It is important that you tell your baby's doctors and other healthcare professionals about your Idacio use during your pregnancy before the baby receives any vaccine.

Tell your doctor if you are breastfeeding or plan to breastfeed.

It is not known whether Idacio passes into breast milk. If you are breastfeeding, your doctor may advise you to stop breastfeeding while you are using this medicine.

If you have not told your doctor or pharmacist about any of the above, tell them before you start using Idacio.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Idacio may interfere with each other.

Tell your doctor or pharmacist if you are taking anakinra (Kineret) or abatacept (Orencia)

Taking either of these two medicines together with Idacio may increase the risk of infection.

Tell your doctor if you are taking azathioprine or 6-mercaptopurine with Idacio.

Tell your doctor if you are taking any other medicines to treat your condition.

Idacio can be taken together with other medicines such as: methotrexate and other disease-modifying anti-rheumatic agents (for example, sulfasalazine, hydroxychloroquine, leflunomide and injectable gold preparations), steroids or pain medications including non-steroidal anti-inflammatory drugs (NSAIDs) such as ibuprofen.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while using this medicine.

How to use Idacio

Idacio is given as a subcutaneous injection (under the skin). It may be injected by the patient, family member or carer.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet.

If you do not understand the instructions on the label or in this leaflet, ask your doctor or pharmacist for help.

Always use Idacio exactly as your doctor has instructed you.

You should check with your doctor or pharmacist if you are not sure.

Patients requiring a dose less than 40 mg should use the 40 mg vial presentation of Idacio.

How much to use

Rheumatoid Arthritis in Adults

The usual dose for adults with rheumatoid arthritis, is one 40mg injection every fortnight.

If you are receiving Idacio without methotrexate, your doctor may change your Idacio dose to 40 mg every week or 80 mg every fortnight, depending on your response.

Other medicines may be prescribed by your doctor to be taken while you are being treated with Idacio.

Psoriatic Arthritis & Ankylosing Spondylitis in Adults

The usual dose for patients with psoriatic arthritis and ankylosing spondylitis is one 40 mg injection every fortnight.

Other medicines may be prescribed by your doctor to be taken while you are being treated with Idacio.

Crohn's disease & Ulcerative Colitis in Adults

The usual dose for adults with Crohn's disease or ulcerative colitis is as follows:

- initial dose of 160 mg (day 1) (given as four 40 mg injections in one day OR as two 40 mg injections per day over two consecutive days) - 80 mg at day 15 (given as two 40 mg injections in one day)
- 40 mg starting at day 29, which then continues once every fortnight (maintenance dose).

Your doctor may change this ongoing (maintenance) dose to 40 mg every week or 80 mg every fortnight, depending on your response.

Psoriasis in Adults

The usual dose for adult patients with psoriasis is as follows:

- initial dose of 80 mg (day 1) (given as two 40 mg injections in one day)
- 40 mg given one week later (day 8), then
- 40 mg every fortnight from day 22 (maintenance dose).

Your doctor may change this ongoing (maintenance) dose to 40 mg every week or 80 mg every fortnight, depending on your response.

Uveitis in Adults

The usual dose for adults with uveitis is as follows:

- initial dose of 80 mg (day 1) (given as two 40 mg injections in one day),
- 40 mg one week later (day 8), then
- 40 mg every fortnight, starting at day 22, which then continues (maintenance dose).

Hidradenitis suppurativa in Adults

The usual dose for adults with HS is as follows:

- initial dose of 160 mg (day 1) (given as four 40 mg injections in one day OR as two 40 mg injections per day over two consecutive days)
- 80 mg dose (day 15) (given as two 40 mg injections in one day)
- 40 mg every week or 80 mg every fortnight starting at day 29, which then continues (maintenance dose).

Your doctor may prescribe other medicines for your condition to take with Idacio.

Juvenile Idiopathic Arthritis & Enthesitis-Related Arthritis

The usual dose for children with pJIA, or ERA depends on body weight.

For a body weight of 30 kg and above:

The usual dose is 40 mg given every fortnight.

For a body weight between 10 kg and less than 30 kg:

The usual dose is 20 mg given every fortnight.

Crohn's disease in Children

The usual dose for children with Crohn's disease depends on body weight.

For a body weight of 40 kg or above:

- initial dose of 160 mg (day 1) (given as four 40 mg injections in one day OR as two 40 mg injections per day over two consecutive days)
- 80 mg two weeks later (day 15) (given as two 40 mg injections in one day).
- 40 mg every fortnight starting at day 29, which then continues (maintenance dose).

Your doctor may change this ongoing (maintenance) dose to 40 mg every week or 80 mg every fortnight, depending on your response.

For a body weight of less than 40 kg:

- initial dose of 80 mg (day 1) (given as two 40 mg injections in one day)
- 40mg two weeks later (day 15)
- 20mg every fortnight, starting at day 29, which then continues (maintenance dose).

Your doctor may change this ongoing (maintenance) dose to 20 mg every week, depending on your response.

Treatment of Crohn's disease in children should be supported by good nutrition to allow appropriate growth.

Psoriasis in Children

The usual dose for children with psoriasis depends on body weight.

For a body weight of 40 kg and above:

The usual dose is 40mg given once every week for the first two weeks, then once every fortnight.

For a body weight of less than 40 kg:

The usual dose is 20 mg given once every week for the first two weeks, then once every fortnight.

Hidradenitis suppurativa in Adolescents

The usual dose for with HS, (from 12 years of age, weighing at least 30 kg) is as follows:

- initial dose of 80 mg (day 1) given as two 40 mg injections in one day),
- 40 mg one week later (day 8).
- 40 mg every fortnight starting at day 22, which then continues (maintenance dose).

Your doctor may change this ongoing (maintenance) dose to 40 mg every week or 80 mg every fortnight, depending on your response.

It is recommended you use an antiseptic wash daily on the affected areas.

How to use it

Idacio is injected under the skin. The injection can be self-administered or given by another person, for example a family member friend or carer, but only after proper training in injection technique.

If you are using the Idacio syringe, instructions for preparing and giving an injection of Idacio are provided in the Injecting instructions supplied with the product.

Read these instructions carefully and follow them step by step. These instructions explain how to self-inject this medicine.

Do not attempt to self-inject until you are sure that you understand how to prepare and give the injection.

Your doctor or his/her assistant will also show you best how to self-inject.

Keep Idacio out of the sight and reach of children.

How to use it

If you are using the Idacio prefilled syringe



Read Carefully these entire instructions before using your Idacio prefilled-syringe.

Only use Idacio pre-filled syringe if your healthcare professional has trained you on how to use the pre-filled syringe correctly.


Important Information:

- Idacio is a pre-filled syringe for single use only.
- Idacio pre-filled syringe has a clear needle guard that covers the needle after the injection is complete.
- Children under 12 years of age are not allowed to inject themselves and injection must be done by a trained adult.
- Keep Idacio pre-filled syringe and the sharps disposal container out of reach and sight of children.
- Do not shake. Shaking can damage the pre-filled syringe and the medicine.
- Do not use Idacio the pre-filled syringe if liquid appears cloudy or discoloured, or has particles or flakes in it. The liquid should be clear and colourless.

- Do not try to activate the clear needle guard before injecting.
- Do not insert your fingers into the opening of the clear needle guard.
- Do not use an Idacio pre-filled syringe that has been frozen or left in direct sunlight.
- Do not use the Idacio pre-filled syringe if it has been dropped or crushed, as the pre-filled syringe may be broken even if you cannot see the break. Use a new pre-filled syringe instead.

Get Familiar with your Idacio Pre-Filled Syringe:

Before Use



Step 1 Prepare for your Injection

Each box of Idacio pre-filled syringe comes with two syringes.

1.1 Prepare a clean flat surface, such as a table or countertop, in a well-lit area.

1.2 You will also need (Figure A):



Figure A

- an alcohol swab (included in the box)
- a cotton ball or gauze, and
- a sharps disposal container.

Open your sharps disposal container so it is ready to use.

1.3 Remove the box from the refrigerator (Figure B).




Figure B

1.4 Check the expiration date on the side of the box (Figure B).

Warning: Do not use if expiration date has passed.

1.5 Caution: Do not pick up the syringe by the plunger or the needle cap. Doing so could damage the syringe or activate the clear needle guard.

Take a syringe out of the original box:

- place two fingers on middle of the clear needle guard
- pull the syringe straight up and out of the packaging (Figure C)



Figure C

Put it on a clean flat surface.

1.6 Place the remaining syringe in its original box back in the refrigerator (Figure D).




Figure D

Refer to Storage information for how to store your unused syringe.

1.7 Leave the syringe at room temperature for 30 minutes to allow the medicine to warm up. Injecting cold medicine can be painful (Figure E).



Figure E

Warning: Do not warm the syringe any other way, such as in a microwave, hot water, or direct sunlight.

Warning: Do not remove the needle cap while allowing syringe to reach room temperature.

Step 2 Wash your hands

2.1 Wash your hands well with soap and water (Figure F) and dry them.

Warning: Gloves will not replace the need for washing hands.



Figure F

Step 3 Check the Syringe

3.1 Check the syringe to make sure that:

- The syringe, the clear needle guard, and the needle cap are not cracked or damaged (Figure G).




Figure G

- The needle cap is securely attached (Figure H).



Figure H

- The needle guard spring is not extended (Figure I).



Figure I

Warning: Do not use the syringe if it shows any sign of damage.

If so, throw away the syringe in a sharps disposal container and contact your healthcare professional or pharmacist.

3.2 Check the liquid to make sure that:

- The liquid is clear, colourless, and free of particles (Figure J).



Figure J

Warning: Do not use the syringe if liquid contains particles, or is cloudy or if it is coloured or has flakes in it.

3.3 Check the label to make sure that:

- The name on the syringe says Idacio (Figure K).
- The expiration date on syringe has not passed (Figure K).



Figure K

Warning: Do not use the syringe if:

- The name on the syringe is not Idacio.
- The expiration date on the syringe has passed.

If so, throw away the syringe in a sharps disposal container and contact your healthcare professional or pharmacist.

Step 4 Choose the Injection Site

4.1 Choose an injection site (Figure L) on:

- Top of the thighs.
- Abdomen (inject at least 2 inches away from the belly button).



Figure L

4.2 Choose a different site (at least one inch away from the previous injection site) each time to reduce redness, irritation or other skin problems.

Warning: Do not inject into an area that is sore (tender), bruised, red, hard, scarred or where you have stretch marks.

Warning: If you have psoriasis, do not inject into any lesions or red, thick, raised or scaly patches.

Step 5 Clean the Injection Site

5.1 Wipe the skin of your injection site with an alcohol swab to clean it. (Figure M).

Warning: Do not blow on or touch the injection site after cleaning.



Figure M

Step 6 Give your Injection

6.1 Remove the needle cap

- Always hold the syringe by the clear needle guard.
- Hold the syringe upward and pull the needle cap straight off (Figure N).

You may see drops of liquid at the needle tip.



Figure N

Warning: Do not touch the needle.

6.2 Pinch the skin

- Hold the syringe like a pencil.
- With your other hand gently pinch skin (without squeezing) to avoid injecting into a muscle (Figure O).



Figure O

6.3 Insert the needle

- With a quick, short motion, push the needle all the way into the skin at an angle between 45° and 90° (Figure P).



Figure P

- After the needle is inserted, release the pinched skin.

6.4 Inject

- Use your thumb to gently push plunger all the way down (Figure Q).



Figure Q

- Give plunger a final push to ensure the full dose has been injected (Figure R).
- Hold the syringe firmly without moving it, at the same angle (Figure R).



Figure R

Do not remove the needle from the skin when the plunger reaches the end.

- Slowly release your thumb up. This will allow the needle to move up into the clear needle guard and cover the entire needle (Figure S).



Figure S

Warning: Call your healthcare professional or pharmacist if:

- You did not inject the full dose or
- The clear needle guard does not activate after injecting.

Warning: Do not reuse a syringe in case of partial injection.

Do not try to recap needle as it could lead to needle stick injury.

6.5 If there is blood or liquid on the injection site, gently press a cotton ball or gauze on the skin (Figure T).



Figure T

Step 7 Throw away your Syringe

7.1 Throw away your used syringe in a sharps disposal container right away after use (Figure U).



Figure U

Warning: Keep your sharps disposal container out of the reach of children.

Warning: Do not throw away the syringe in your household trash.

If you do not have a sharps disposal container, you may use a household container that is:

- Made of a heavy-duty plastic;
- Can be closed with a tight-fitting, puncture-resistant lid; that will keep sharps from coming out,
- Upright and stable during use,
- Leak-resistant and
- Properly labelled to warn of hazardous waste inside the container.

7.2 When your sharps disposal container is almost full, you will need to follow your local guidelines for the right way to dispose of your sharps disposal container.

Do not recycle your used sharps disposal container.

Step 8 Record your Injection

8.1 To help you remember when and where to do your next injection, you should keep a record of the dates and injection sites used for your injections (Figure V).



Figure V

How long to use it

Keep using Idacio for as long as your doctor tells you.

Idacio will not cure your condition but should help your symptoms.

Ask your doctor if you are not sure how long to take the medicine for.

If you forget to use it

If you forget to give yourself an injection, you should inject the next dose of Idacio as soon as you remember. Then inject your next dose as you would have on your originally scheduled day.

Do not try to make up for missed doses by injecting more than one dose at a time.

If you use too much (overdose)

If you accidentally inject Idacio more frequently than prescribed by your doctor, immediately telephone your doctor or the Poisons Information Centre (Australia Telephone 13 11 26) or go to Accident and Emergency at your nearest hospital. Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention. Always take the outer carton of the medicine with you.

While you are using Idacio

Things you must do

Check with your doctor before you receive any vaccines.

Wherever possible, it is recommended that children be brought up to date with all immunisations in according to current immunisation guidelines prior to starting Idacio therapy.

Patients receiving Idacio should not receive live vaccines (for example, BCG or oral polio vaccine).

If you become pregnant while using Idacio, tell your doctor immediately.

If you are about to be started on any new medicine, tell your doctor you are using Idacio.

Tell all doctors, dentists, and pharmacists who are treating you that you are using Idacio.

If you are going to have surgery, tell all doctors that you are using Idacio. Your doctor may recommend you discontinue using Idacio temporarily.

Keep all of your doctor's appointments so that your progress can be checked.

Things you must not do

Do not give Idacio to anyone else, even if they have the same condition as you.

Do not use Idacio to treat any other complaints unless your doctor tells you to.

Do not stop taking Idacio, without checking with your doctor.

Do not take Idacio and anakinra (Kineret) or Idacio and abatacept (Orencia) together.

Taking either of these two medicines with Idacio may lead to an increased risk of developing a serious infection.

Things to be careful of

It is important to tell your doctor if you get symptoms of an infection, such as a fever, skin sores, feeling tired or any problems with your teeth and gums.

You might get infections more easily while you are receiving Idacio treatment. These infections may be serious and include tuberculosis, infections caused by viruses, fungi, parasites or bacteria, or other infections. Sepsis, an infection of the blood, may, in rare cases, be life-threatening.

Your doctor may recommend you discontinue Idacio if you develop an infection.

Be careful driving or operating machinery until you know how Idacio affects you.

The effects on your ability to drive and use machines whilst taking this medicine are not known.

Side Effects

Tell your doctor as soon as possible if you do not feel well while using Idacio or you have any problems using it.

Do this even if you do not think the problems are connected with the medicine or are not listed in this leaflet.

All medicines can have unwanted side effects. Sometimes they are serious, but most of the time they are not. You may need medical attention if you get some side-effects.

Do not be alarmed by this list of possible side effects. You may not experience any of them.

Ask your doctor or pharmacist any questions you may have.

Tell your doctor immediately or go to Accident and Emergency at your nearest hospital if you experience any of the following:

- Signs of an allergic reaction such as:
 - Chest tightness
 - Shortness of breath, wheezing or difficulty breathing
 - Swelling of the face, lips, tongue or other parts of the body
 - Hives, itching or skin rash.
- Signs and symptoms suggestive of heart failure such as shortness of breath with exertion or upon lying down or swelling of the feet
- Signs and symptoms suggestive of a blood disorder such as persistent fever, bruising, bleeding very easily, paleness.

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation.

Tell your doctor as soon as possible if you notice any of the following:

- Signs of tuberculosis such as persistent cough, weight loss, listlessness, fever
- Signs of infection such as fever, lack of energy, skin sores, problems with your teeth or gums, burning when you pass urine
You might get infections more easily while you are receiving Idacio treatment.
- Signs of nervous system disorders such as numbness or tingling throughout your body, arm or leg weakness, double vision
- Signs of soft tissue infection, such as a bump or open sore that doesn't heal.

The above list includes serious side effects. You may need urgent medical attention. Serious side effects are rare.

Tell your doctor if you notice any of the following and they worry you:

- Pain, swelling, redness or itching at the site of injection
- Cold, runny nose, sinus infection, sore throat, cough, congestion on the chest, asthma or worsening of asthma symptoms

- Lower respiratory tract infections (such as bronchitis, pneumonia)
- Pain in the ear which could suggest an ear infection
- Pain or inflammation of the eye or eye lid or changes to your vision
- Mouth ulcers, pain or excessive bleeding from the gums
- Burning or pain when passing urine, or blood in the urine
- Skin bumps or sores that don't heal
- Headache or migraine, dizziness, vertigo
- Muscle weakness or numbness, difficulty balancing
- Fever, flushing, increased sweating
- Nausea, vomiting, abdominal pain
- Reflux or heartburn
- Chest pain
- Rash, itching, redness or scaly patches
- Problems with your finger or toe nails
- Hair loss
- Fatigue, tiredness, lack of energy
- Muscle, joint or bone pain
- Bleeding or bruising more easily than usual
- Feeling overwhelmed or sad, or lacking motivation (depression)
- Feeling anxious, especially fearful or worried (anxiety)
- Increased heart rate
- Viral infections (including the flu, cold sore blisters, chicken pox and shingles)
- Bacterial infections (including urinary tract infection)
- Fungal Infections.

The above list includes the more common side effects of Idacio. They are usually mild and short-lived.

There have been cases of certain kinds of cancer in patients using Idacio or similar medicines. People with more serious rheumatoid arthritis that have had the disease for a long time may have a higher chance of getting a kind of cancer that affects the lymph system, called lymphoma, or that affects the blood, called leukaemia. If you take Idacio your risk may increase.

On rare occasions, a specific and severe type of lymphoma has been observed in patients taking Idacio.

Tell your doctor if new skin lesions (skin spots or sores) appear, or if existing lesions change appearance during or after Idacio treatment.

Very rare cases of skin cancer have been observed in patients taking Idacio.

If you have COPD, or are a heavy smoker, you should discuss with your doctor whether treatment with a TNF blocker is right for you.

There have been cases of cancers other than lymphoma in patients with a specific type of lung disease called Chronic Obstructive Pulmonary Disease (COPD) treated with another TNF blocker.

Laboratory results

Some side effects observed with Idacio may not have symptoms and may only be discovered through blood tests. These include, most commonly, increased lipids, elevated liver enzymes and low levels of

white blood cells, and red blood cells in the blood.

Tell your doctor or pharmacist if you notice anything else that is making you feel unwell.

Other side effects not listed above may occur in some people.

After using Idacio

Keep your pre-filled syringe in the pack until it is time to use it.

Keep Idacio in a refrigerator (2°C to 8°C). Do not freeze.

Keep Idacio in the refrigerator in away children cannot get to it.

Keep the medicine at the right temperature when you travel. This is important when travelling by car, bus, train, plane or any other form of transport.

Store a pre-filled syringe at room temperature (below 25°C) for a maximum period of 14 days, protected from light.

Once removed from the refrigerator and stored at room temperature, the syringe must be used within 14 days or discarded, even if it is returned to the refrigerator.

Write down the date you first remove the syringe from the refrigerator on the label, so you can check how long it has been.

Disposal

After injecting Idacio, immediately throw away the used pre-filled syringe in a special 'sharps' container as instructed by your doctor, nurse or pharmacist. If your doctor tells you to stop using Idacio or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

Product Description

What it looks like

Idacio is a clear, colourless solution.

Ingredients

Adalimumab

Monobasic sodium phosphate dihydrate

Dibasic sodium phosphate dihydrate

Mannitol

Sodium chloride

Citric acid monohydrate

Sodium citrate dihydrate

Polysorbate 80

Sodium hydroxide (for pH adjustment)

Water for injections

Supplier

Idacio is supplied in Australia by:

Fresenius Kabi Australia Pty Limited
Level 2, 2 Woodland Way Mount Kuring-
gai, NSW, 2080 Australia
Telephone: (02) 9391 5555

Idacio is supplied in New Zealand by:

Fresenius Kabi New Zealand Limited
60 Pavilion Drive
Airport Oaks, Auckland 2022
New Zealand
Freecall: 0800 144 892

Australian Registration Number

Aust R 320241

® = Registered Trademark

This leaflet was prepared on 16 June 2020.