

## FOSAMAX® Once Weekly

Alendronate sodium

### Consumer Medicine Information

#### What is in this leaflet

This leaflet answers some common questions about FOSAMAX Once Weekly. It is particularly important that you read the sections "When to take it" and "How to take it" before you take this medicine. This leaflet does not contain all the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking FOSAMAX Once Weekly against the benefits they expect it will have for you.

**If you have any concerns about taking this medicine, ask your doctor or pharmacist.**

**Keep this leaflet with the medicine.**

You may need to read it again.

#### What FOSAMAX Once Weekly is used for

FOSAMAX Once Weekly is used to treat osteoporosis.

This condition is caused by changes in the way bone is normally maintained.

#### Understanding bone

Bone is a living, growing tissue. Throughout life, our bodies are breaking down old bone and rebuilding new bone in a continuous cycle. Until our late 20s, while bones are still developing, we gain bone by building more than we lose. From then until about age 35 the process is usually in balance, so that the amount of bone lost is about equal to the amount that is replaced. This balanced process keeps your skeleton healthy and strong. After about age 35 this balance is disturbed, with bone loss occurring at a slightly faster rate than it can be replaced. In women, after menopause, hormonal changes cause bone loss at an even faster rate. When bone loss is excessive, bones can become thinner and weaker, and therefore are more likely to break.

#### Osteoporosis

"Osteo" means bone, and "porosis" means something that has holes in it, like a sponge. Therefore, osteoporosis is a disease which causes bones to become more porous, gradually making them weaker, more brittle and likely to break. Osteoporosis is common in postmenopausal women. The menopause occurs when the ovaries virtually stop producing the female hormone, oestrogen, or are removed (which may occur, for example, at the time of a hysterectomy). At this time, bone is removed faster than it is formed, so bone loss occurs and bones become weaker. The earlier a woman reaches the menopause, the greater the risk of osteoporosis.

Osteoporosis also occurs in men but is less common than in women.

Osteoporosis can also occur in people receiving corticosteroid medicines. If taken in high doses or for a long period of time, corticosteroid medicines can cause bone to be removed faster than it is formed. This causes loss of bone and therefore, bones become weaker and are more likely to break.

Maintaining bone mass and preventing further bone loss are important to keep your skeleton healthy.

Early on, osteoporosis usually has no symptoms. However, if left untreated it can result in broken bones, also called fractures. Although fractures

usually cause pain, fractures of the bones of the spine may go unnoticed until they cause height loss. Fractures may occur during normal, everyday activity, such as lifting, or from minor injury that would not ordinarily fracture normal bone. Fractures usually occur at the hip, spine, or wrist and can lead not only to pain, but also to considerable deformity and disability, such as stooped posture from curvature of the spine, and loss of mobility.

#### How does FOSAMAX Once Weekly work?

FOSAMAX Once Weekly works by slowing down the process of old bone being removed, which allows the bone-forming cells time to rebuild normal bone. FOSAMAX Once Weekly not only helps prevent the loss of bone but actually helps to rebuild bone and makes bone less likely to fracture. Thus, FOSAMAX Once Weekly reverses the progression of osteoporosis. FOSAMAX Once Weekly starts working on the bone cells immediately, but measurable effects on bone mass may not be seen for several months or more.

FOSAMAX Once Weekly belongs to a group of non-hormonal medicines called bisphosphonates.

#### Before you take FOSAMAX Once Weekly

You should know that in some people, FOSAMAX Once Weekly can irritate or burn the food pipe (also called oesophagus). The chances of this happening should be reduced when you follow the instructions for 'How to take FOSAMAX Once Weekly' section in this leaflet.

#### When you must not take it

**Do not take FOSAMAX Once Weekly if:**

- you have an allergy to FOSAMAX Once Weekly or any of the ingredients listed at the end of this leaflet
- you have certain disorders of the food pipe (oesophagus) including those that cause difficulty in swallowing
- you are unable to stand or sit upright for at least 30 minutes
- your doctor has told you that you currently have low blood calcium
- your dentist advises you to consult your doctor first

**Do not take FOSAMAX Once Weekly if you are pregnant or breast-feeding.**

FOSAMAX Once Weekly has not been studied in pregnant or breast-feeding women.

**Do not take FOSAMAX Once Weekly if:**

- the packaging is torn or shows signs of tampering
- the expiry date on the pack has passed.

If you take this medicine after the expiry date has passed, it may not work.

**If you are not sure whether you should start taking FOSAMAX Once Weekly, talk to your doctor.**

**Do not give FOSAMAX Once Weekly to a child.**

FOSAMAX Once Weekly has not been studied in children.

#### Before you start to take it

**Tell your doctor if:**

1. you plan to become pregnant or breast-feed.

2. you have any medical conditions, especially the following:
  - kidney disease
  - swallowing or digestive problems, such as ulcers
3. you have any allergies to any other medicines or any other substances, such as foods, preservatives or dyes.
4. you have dental or jaw-bone problems or are planning to have a course of dental surgery.
5. you currently smoke or have been a smoker in the past.

**If you have not told your doctor about any of the above, tell them before you take any FOSAMAX Once Weekly.**

#### Taking other medicines

**Tell your doctor if you are taking any other medicines, including medicines that you buy without a prescription from your pharmacy, supermarket or health food shop.**

Some medicines may affect the way other medicines work.

**Some medicines are likely to interfere with the absorption of FOSAMAX Once Weekly if taken at the same time. These include:**

- antacids, medicines used to treat indigestion e.g. Gaviscon, Mylanta
- calcium supplements
- vitamins

**Therefore, take FOSAMAX Once Weekly at least 30 minutes before taking any of these or other medicines to make sure there is no problem with absorption. Check with your doctor or pharmacist if you are not sure whether you are taking any of these medicines.**

You can take aspirin while you are being treated with FOSAMAX Once Weekly. However, both aspirin and FOSAMAX Once Weekly may increase the chance of stomach upsets.

Your doctor or pharmacist has more information on medicines to be careful with or avoid while taking FOSAMAX Once Weekly.

#### How to take FOSAMAX Once Weekly

##### How much to take

**Take FOSAMAX Once Weekly only when prescribed by your doctor.**

The usual dose of FOSAMAX Once Weekly is 70 mg tablet once a week.

**Choose the day of the week that best fits your schedule. Every week, take one tablet of FOSAMAX Once Weekly on your chosen day.**

**Follow all directions given to you by your doctor carefully.**

They may differ from the information contained in this leaflet.

**If you do not understand the instructions on the box, ask your doctor or pharmacist for help.**

##### When and how to take it

**Take FOSAMAX Once Weekly after getting up for the day and before taking your first food, beverage or other medication. Do not take it at bedtime.**

**Swallow one tablet whole with a full glass of plain water [not mineral water, not coffee or tea, not juice].**

Do not take any food, medicines or drinks other than plain tap water with your FOSAMAX Once Weekly. It is important to take FOSAMAX Once Weekly with plain water only, not mineral water. Food, other drugs, mineral water and other drinks, including fruit juices, coffee and tea, will reduce the effect of FOSAMAX Once Weekly by interfering with the absorption into the body.

**Stay upright for at least 30 minutes after swallowing FOSAMAX Once Weekly and do not take any food, medicines or drinks other than plain tap water during this time. Do not lie down immediately after swallowing it.**

It is important to stay upright (sitting, standing or walking around) for at least 30 minutes after swallowing your tablet.

**It is also very important to stay upright until after you have eaten your first food of the day.**

These actions will help make sure your dose of FOSAMAX Once Weekly reaches your stomach quickly and help reduce the potential for irritation to your food pipe (oesophagus).

FOSAMAX Once Weekly is effective only if taken when your stomach is empty. Food, drinks other than plain water, and other medicines will lessen the effect of FOSAMAX Once Weekly by interfering with its absorption into the body.

**Do not chew or suck on a tablet of FOSAMAX Once Weekly.**

Mouth ulcers may occur if the tablet is chewed or dissolved in the mouth.

#### **How long to take it**

It is important that you take FOSAMAX Once Weekly for as long as your doctor prescribes it. FOSAMAX Once Weekly can only treat your osteoporosis, by helping prevent further loss of bone and continuing to rebuild bone, if you take it every week. Since it is not known how long you should take FOSAMAX Once Weekly, you should discuss the need to stay on this medication with your doctor periodically to determine if FOSAMAX Once Weekly is still right for you.

#### **If you forget to take it**

**If you miss a tablet, take one tablet on the morning after you remember.**

**Do not take two tablets on the same day. Return to taking one tablet once a week, as originally scheduled on your chosen day.**

**If you are not sure about what to do, talk to your doctor or pharmacist.**

**If you have trouble remembering to take your FOSAMAX Once Weekly, ask your pharmacist for some hints.**

#### **If you take too much (overdose)**

**Immediately telephone your doctor or Poisons Information Centre (telephone 13 11 26), or go to accident and emergency at your nearest hospital, if you think that you or anyone else may have taken too much FOSAMAX Once Weekly. Do this even if there are no signs of discomfort or poisoning.**

**If you take too many tablets at one time, drink a full glass of milk. Do not induce vomiting. Do not lie down.**

#### **While you are using FOSAMAX Once Weekly**

##### **Things you must do**

**If you develop difficulty or pain upon swallowing, chest pain, or new or worsening heartburn, stop taking FOSAMAX Once Weekly and call your doctor.**

**If you become pregnant while taking FOSAMAX Once Weekly, stop taking FOSAMAX Once Weekly and tell your doctor.**

**If you are about to be started on any new medicine, tell your doctor and pharmacist that you are taking FOSAMAX Once Weekly.**

**If you develop a toothache or require a dental procedure, tell your dentist that you are taking FOSAMAX Once Weekly.**

**If you develop new or unusual pain in your hip or thigh, tell your doctor.**

Rarely, patients have experienced fracture in a specific part of the thigh bone.

**Make sure you have an adequate intake of calcium in your diet.**

Your doctor, dietician or pharmacist can tell you what foods you should eat.

#### **Things you must not do**

**Do not give FOSAMAX Once Weekly to anyone else, even if they have the same condition as you.**

#### **Things to be careful of**

**There have been side effects reported with FOSAMAX Once Weekly that may affect your ability to drive or operate machinery.**

**Individual responses to FOSAMAX Once Weekly may vary. (See Side Effects).**

#### **Things that would be helpful for your osteoporosis**

Some self help measures suggested below may help your osteoporosis. Talk to your doctor or pharmacist about these measures and for more information.

- Exercise - can be helpful in building and maintaining strong bones. Regular exercise such as a brisk walk is a good idea. Talk to your doctor before you begin any exercise program.
- Diet - eat a balanced diet. You may need to increase the amount of calcium in your diet by eating calcium-rich foods or taking a calcium supplement. Your doctor will advise you.
- Smoking - appears to increase the rate at which you lose bone and, therefore, may increase your risk of fracture. Your doctor may ask you to stop smoking or at least cut down.
- Alcohol - your doctor may advise you to cut down the amount of alcohol you drink. If you drink excessively on a regular basis, you may increase your risk of developing osteoporosis.

#### **Side Effects**

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking FOSAMAX Once Weekly.**

FOSAMAX Once Weekly helps most people with osteoporosis, but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

**Ask your doctor or pharmacist to answer any questions you may have.**

**Tell your doctor if you notice any of the following and they worry you:**

- stomach pain, gas in the stomach or bowel, wind
- an uncomfortable feeling in the stomach or belching after eating, also called dyspepsia, or heartburn
- feeling sick (nausea), vomiting

- constipation, diarrhoea
- headache
- aching muscles, joints and/or bones, which rarely can be severe.
- flu-like symptoms typically at the start of treatment, such as aching muscles, generally feeling unwell and rarely fever.
- swelling of joints
- dizziness or spinning sensation
- unusual tiredness or weakness
- swelling of hands, ankles or feet
- hair loss
- changed sense of taste

Most of these are the more common side effects of FOSAMAX Once Weekly. For the most part, these have been mild.

**Tell your doctor immediately if you notice any of the following:**

- skin rash or redness of the skin, sometimes made worse by sunlight, itchiness
- mouth ulcers
- blurred vision, pain or redness in the eye
- ear pain
- symptoms of low blood calcium levels including muscle cramps or spasms or tingling sensation in the fingers or around the mouth
- new or unusual pain in your hip or thigh

These side effects are rare, and very rarely, may be serious.

**Tell your dentist and doctor immediately if you notice any of the following:**

- Jaw-bone or dental problems (including toothache). Jaw-bone problems may include infection, and delayed healing after a tooth extraction or other work that involves drilling into the jaw-bone.

These side effects are rare and may be serious.

**If any of the following happen, stop taking FOSAMAX Once Weekly and tell your doctor immediately:**

- difficulty or pain upon swallowing
- chest pain
- new or worsening heartburn

These side effects may be due to irritation or ulceration of the food pipe. They may worsen if you continue taking FOSAMAX Once Weekly. Rarely, these side effects may be serious.

**If any of the following happen, stop taking FOSAMAX Once Weekly and tell your doctor immediately or go to accident and emergency at your nearest hospital:**

- swelling of the face, lips, mouth, throat or tongue which may cause difficulty in breathing or swallowing
- pinkish, itchy swellings on the skin, also called hives or nettlerash
- severe skin reactions
- black tar-like and/or bloody stools

These may be serious side effects. You may need urgent medical attention. These side effects are rare.

If you have the swelling described above, you may be having a serious allergic reaction to FOSAMAX Once Weekly.

Rarely, stomach or duodenal ulcers (some severe) have occurred, but it is not known whether these were caused by FOSAMAX Once Weekly.

**Other side effects not listed above may also occur in some patients. Tell your doctor if you notice any other effects.**

Do not be alarmed by this list of possible side effects. You may not experience any of them.  
FOSAMAX Once Weekly is not addictive.

---

## After using FOSAMAX Once Weekly

---

### **Storage**

**Keep your tablets in the blister pack until it is time to take them.**

If you take the tablets out of the blister pack they may not keep well.

**Keep FOSAMAX Once Weekly in a cool dry place where the temperature stays below 30°C. Do not freeze the product. Do not store it or any other medicine in the bathroom or near a sink.**

**Do not leave it in the car or on window sills.**

Heat and dampness can destroy some medicines.

**Keep it where children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

### **Disposal**

**If your doctor tells you to stop taking FOSAMAX Once Weekly, or the tablets have passed their expiry date, ask your pharmacist what to do with any that are left over.**

---

## Product Description

---

### **What it looks like**

FOSAMAX Once Weekly comes as a white oval tablet, with 31 marked on one side and an outline of a bone image on the other.

A box of tablets contains 4 tablets.

### **Ingredients**

Active ingredient:

- alendronate sodium equivalent to 70 mg alendronic acid per tablet

Inactive ingredients:

- microcrystalline cellulose
- anhydrous lactose
- croscarmellose sodium
- magnesium stearate

FOSAMAX Once Weekly does not contain sucrose, gluten, tartrazine or any other azo dyes.

### **Supplier**

FOSAMAX Once Weekly is supplied in Australia by:

Merck Sharp & Dohme (Australia) Pty Limited  
Level 1, Building A,  
26 Talavera Road,  
Macquarie Park NSW 2113

This leaflet was prepared in September 2015.

Australian Register Number:

70 mg Tablets - AUST R 73772

WPPI-MK0217-OW-MF-052015