

CARBETOCIN WOCKHARDT Injection

Carbetocin

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about **CARBETOCIN WOCKHARDT**.

It does not contain all the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you having **CARBETOCIN WOCKHARDT** against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor.

Keep this leaflet.

You may need to read it again.

What CARBETOCIN WOCKHARDT is used for

CARBETOCIN WOCKHARDT contains carbetocin, a substance which is an oxytocic agent and causes the womb (uterus) to contract.

CARBETOCIN WOCKHARDT is used to minimise post-delivery bleeding in women who have delivered their baby by elective caesarean section under local anaesthetic around the spinal cord (epidural).

How it works

In some women, following delivery by caesarean section, the womb fails to contract quickly enough which increases the risk of bleeding. Drugs that act directly on the womb to make it contract are used to minimise post-delivery bleeding and are sometimes called oxytocic agents because they act in a similar manner to oxytocin. Ask your doctor if you have any questions about why this medicine has been prescribed for you.

Your doctor may have prescribed it for another reason.

CARBETOCIN WOCKHARDT is available only with a doctor's prescription. It is not addictive.

Before you are given CARBETOCIN WOCKHARDT

When you must not be given it

CARBETOCIN WOCKHARDT must not be given if you have an allergy to:

- carbetocin, the active ingredient, or to any of the ingredients listed at the end of this leaflet
- any medicines containing oxytocin.

Symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin.

You must not be given CARBETOCIN WOCKHARDT during pregnancy or any time before delivery of the baby. should only be given after delivery of your baby.

You must not be given CARBETOCIN WOCKHARDT if you are suffering from serious heart disease.

CARBETOCIN WOCKHARDT should not be given to children

Safety and effectiveness in children have not been established.

CARBETOCIN WOCKHARDT must not be given after the expiry date printed on the syringes and outer packaging.

CARBETOCIN WOCKHARDT must not be given if packaging is torn or shows signs of tampering.

If you not sure whether you should be given CARBETOCIN WOCKHARDT, talk to your doctor.

Before you are given it

Tell your doctor if you have high blood pressure or any heart problems.

Your doctor may want to take extra precautions.

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you have or have had any of the following medical conditions:

- severe heart disease
- high blood pressure
- liver or kidney problems
- problems with blood clotting
- disease of a gland
- gestational diabetes (diabetes that starts during pregnancy)
- epilepsy
- migraine
- asthma.

If you have not told your doctor about any of the above, tell him/ her before you have CARBETOCIN WOCKHARDT.

Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

How CARBETOCIN WOCKHARDT is given

CARBETOCIN WOCKHARDT is a medicine that is used only in hospital and should only be administered by qualified staff.

CARBETOCIN WOCKHARDT will be given to you by injection into one of your veins immediately after the delivery of your baby.

How much you will be given

CARBETOCIN WOCKHARDT will be given as a single dose of 100 micrograms (1 mL) into a vein by injection, slowly over one minute.

Side effects

Tell your doctor or nurse immediately if you do not feel well after you have been given CARBETOCIN WOCKHARDT.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Tell your doctor or nurse if you notice any of the following and they worry you:

- nausea (feeling sick) or vomiting
- stomach pain
- itching
- flushing or feeling of warmth
- feeling light headed, dizzy or faint (low blood pressure)
- headache
- tremor.

These are the more common side effects of **CARBETOCIN WOCKHARDT**. Mostly these are mild and short-lived.

Tell your doctor or nurse immediately if you notice any of the following:

- back pain
- dizziness
- metallic taste
- signs of anaemia, such as tiredness, being short of breath and looking pale
- sweating
- chills
- nervousness, feeling anxious. All of these side effects are rare.
- **Tell your doctor immediately or go to Accident and Emergency at your nearest hospital if you notice the following:**
- chest pain
- fast or abnormal heart beat
- shortness of breath
- swelling of the face, lips, tongue or other parts of the body
- cold clammy skin.
- fainting or palpitations
- The above list includes very serious side effects. You may need urgent medical attention or hospitalisation.

Other side effects not listed above may also occur in some patients. Tell your doctor or nurse if you notice anything that is making you feel unwell.

Do not be alarmed by this list of possible side effects.

You may not experience any of them.

Ask your doctor to answer any questions you may have.

After using CARBETOCIN WOCKHARDT

Storage

CARBETOCIN WOCKHARDT is usually stored in the doctor's surgery or clinic, or at the pharmacy. However, if you need to store **CARBETOCIN WOCKHARDT**:

Keep it where children cannot reach it.

Keep CARBETOCIN WOCKHARDT in the original container until it is time for it to be given.

Keep syringe in the refrigerator between 2°C to 8°C. Do NOT Freeze.

If you take your medicine out of the original container, it will not keep well.

Product description

What it looks like

CARBETOCIN WOCKHARDT is a clear colourless solution in a 1 mL clear glass syringe

Ingredients

CARBETOCIN WOCKHARDT contains 100 micrograms of carbetocin as the active ingredient.

It also contains:

- sodium chloride
- water for injections.
- glacial acetic acid

This medicine does not contain lactose, sucrose, gluten, tartrazine or any other azo dyes.

Sponsor

CARBETOCIN WOCKHARDT is supplied in Australia by:

Wockhardt Bio Pty Ltd

Suite 103, 39 East

Esplanade

MANLY NSW 2095

AUSTRALIA

Australian Registration Number

AUST R 305583–

CARBETOCIN WOCKHARDT

carbetocin 100

microgram/1 mL

syringe in packs of 5

This leaflet was prepared in May 2020