

CEPHALEXIN GENERICHEALTH

(cefalexin monohydrate) capsule

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Cephalexin generichealth. It does not contain all the available information. It does not take the place of talking with your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Cephalexin generichealth against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, talk to your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What Cephalexin generichealth is used for

Cephalexin generichealth contains cefalexin monohydrate as the active ingredient.

It is used to treat infections caused by bacteria in different parts of the body, including infections of the:

- respiratory tract (throat, tonsils, chest and lungs);
- nose (sinusitis);
- ears (otitis media);
- skin and soft tissue;
- kidneys and bladder (genitourinary tract).

This medicine belongs to a group of medicines called cephalosporins. These antibiotics work by killing the bacteria that are causing your infection.

Ask your doctor or pharmacist if you have any questions about why this medicine has been prescribed for you.

Your doctor may have prescribed it for another reason.

This medicine is available only with a doctor's prescription.

There is no evidence that it is addictive.

Before you take Cephalexin generichealth

When you must not take it

Do not take Cephalexin generichealth if you have allergic reaction to:

- any medicine containing cefalexin monohydrate;
- other cephalosporins;
- any of the ingredients listed at the end of this leaflet.

Do not take this medicine if you have had a serious allergic reaction to penicillin.

Some of the symptoms of an allergic reaction may include shortness of breath, wheezing or difficulty breathing; swelling of the face, lips, tongue or other parts of the body; rash, itching or hives on the skin.

Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start taking this medicine, talk to your doctor.

Before you start to take it

Tell your doctor if you are allergic to any other medicines, or any foods, preservatives or dyes.

Tell your doctor if you have or have had any of the following medical conditions:

- liver disease;
- kidney disease;
- bowel disease.

Tell your doctor if you are pregnant or plan to become pregnant or are breastfeeding.

Your doctor can discuss with you the risks and benefits involved.

If you have not told your doctor about any of the above, tell them before you start taking Cephalexin generichealth.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and cefalexin may interfere with each other. These include:

- probenecid, a medicine commonly used to treat gout;
- metformin, a medicine used in the treatment of diabetes.

These medicines may be affected by Cephalexin generichealth or may affect how well it works. You may need different amounts of your medicines or you may need to take different medicines. Your doctor or pharmacist has more information on medicines to be careful with or avoid while taking Cephalexin generichealth.

How to take Cephalexin generichealth

Follow all directions given to you by your doctor or pharmacist.

They may differ from the information contained in this leaflet.

If you do not understand the instructions on the carton, ask your doctor or pharmacist for help.

How much to take

Your doctor will tell you how much and how often you should take Cephalexin generichealth. This will depend on the type of infection. The dose varies from patient to patient.

Adults: The usual adult dose is 250 mg every 6 hours.

Children: Your child's doctor will tell you how much cefalexin your child should take. This will depend on your child's age, weight and the type of infection.

The 250 mg strength is available in other brands.

Oral suspension is not available in Cephalexin generichealth. It is available in other brands. For more information refer to the relevant brand product information.

How to take it

Swallow the capsules whole with a full glass of water.

When to take it

It does not matter if you take this medicine with or without food.

How long to take it

Continue taking your medicine for as long as your doctor tells you.

It is important to complete the full course prescribed by your doctor, even if you begin to feel better after a few days. If you do not, the bacteria causing your infection may not clear completely or your symptoms may return.

If you forget to take it

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Otherwise, take it as soon as you remember, and then go back to taking your medicine as you would normally.

Do not take a double dose to make up for the dose that you missed.

If you are not sure what to do, ask your doctor or pharmacist.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

If you take too much (overdose)

Immediately telephone your doctor or Poisons Information Centre (13 11 26) for advice, or go to Accident and Emergency at the nearest hospital, if you think you or anyone else may have taken too much Cephalexin generichealth. Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

Symptoms of an overdose may include feeling sick in the stomach, diarrhoea, blood in the urine, unusually increased reflexes, convulsions or hallucinations.

While you are taking Cephalexin generichealth

Things you must do

Tell your doctor if the symptoms of your infection do not improve within a few days or if they become worse.

Tell any other doctors, dentists and pharmacists who are treating you that you are taking Cephalexin generichealth, especially if you are about to be started on any new medicines.

Tell your doctor if you become pregnant while taking this medicine.

If you are about to have any blood tests, tell your doctor that you are taking Cephalexin generichealth.

It may interfere with the results of some tests.

If you are diabetic, check with your doctor or pharmacist before using urine sugar tests.

Cefalexin may affect the results of some of these tests.

Keep all of your doctor's appointments so that your progress can be checked.

Your doctor may do some tests from time to time to make sure the medicine is working and to prevent unwanted side effects.

Things you must not do

Do not take Cephalixin generichealth to treat any other complaints unless your doctor tells you to.

Do not give your medicine to anyone else, even if they have the same condition as you.

Do not stop taking your medicine or change the dosage without first checking with your doctor.

If you stop taking it, your condition may worsen.

Things to be careful of

Be careful driving or operating machinery until you know how Cephalixin generichealth affects you.

It may cause dizziness or drowsiness in some people.

Children should be careful when riding bicycles or climbing trees.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Cephalixin generichealth.

This medicine helps most people with infection but it may have unwanted side effects in a few people.

All medicines have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor if you notice any of the following and they worry you:

- sore or white mouth or tongue (oral thrush);
- sore and itchy vagina and/or discharge (another form of thrush);
- diarrhoea;
- nausea;
- vomiting;
- dizziness;
- unusual tiredness/weakness;
- abdominal pain;
- headache.

The above list includes the more common side effects of your medicine.

Tell your doctor as soon as possible if you notice any of the following and they worry you:

- severe body rash;
- swelling of face, lips, mouth, tongue or throat which may cause difficulty in swallowing or breathing;
- fever;
- swelling of the joints;
- yellowing of the skin or eyes;
- joint pain;
- confusion and hallucinations;
- seizure
- encephalopathy (such as reduced ability to think clearly or concentrate, memory loss, drowsiness, seizures, muscle twitches and personality change)
- myoclonus
- bleeding or bruising more easily than normal.

The above list includes serious side effects which may require medical attention. Serious side effects are rare.

Tell your doctor immediately if you notice any of the following, particularly if they occur several weeks after stopping treatment with Cephalixin generichealth:

- severe stomach cramps or pain;
- severe, watery or bloody diarrhoea;
- fever, in combination with one of the above.

Do not take any diarrhoea medicine without first checking with your doctor.

You may have a serious condition affecting your bowel requiring urgent medical attention.

Tell your doctor immediately or go to Accident and Emergency at your nearest hospital if you notice any of the following:

- sudden signs of allergy such as rash, itching or hives on the skin with swelling of the face, lips, tongue or other parts of the body, shortness of breath, wheezing or trouble breathing.

This is a very serious side effect. You may need urgent medical attention or hospitalisation. This side effect is very rare.

Tell your doctor or pharmacist if you notice anything else that is making you feel unwell.

Other side effects not listed above may also occur in some people. Some of these side effects can only be found when your doctor does tests from time to time to check your progress.

Do not be alarmed by this list of possible side effects. You may not experience any of them.

After taking Cephalixin generichealth

Storage

Keep your capsules in the blister pack until it is time to take them.

If you take your capsules out of the blister pack, they may not keep as well.

Store below 30°C.

Do not store Cephalixin generichealth or any other medicine in the bathroom or near a sink. Do not leave it on a window sill or in the car.

Heat and dampness can destroy some medicines.

Keep your medicine where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

Product description

What it looks like

Cephalixin generichealth 500 mg are size '0' capsules with dark green cap imprinted with "500" in black ink and light green body without imprint containing white to yellowish-white granular powder.

Available in packs of 20.

Ingredients

Active ingredient

Cephalixin generichealth capsules contain 500 mg of cefalexin (as monohydrate) as the active ingredient.

Other ingredients

- microcrystalline cellulose;
- magnesium stearate;
- gelatin;
- purified water;
- sodium lauryl sulfate;
- sunset yellow FCF;
- quinoline yellow;
- titanium dioxide;
- patent blue V;
- TekPrint SW-9808 Black Ink (PI 2328) or Marking Solution Black BKI004 Black Ink ACG PAM BKI004 (PI 139674).

Cephalixin generichealth 500 mg capsules do not contain lactose, sucrose, gluten, tartrazine or any other azo dyes.

Australian Registration Numbers

Cephalixin generichealth 500 mg: AUST R 154381.

Sponsor

Generic Health Pty Ltd
Suite 2, Level 2
19-23 Prospect Street
Box Hill, VIC, 3128
Australia

E-mail: ghinfo@generichealth.com.au

Telephone: +61 3 9809 7900

Website: www.generichealth.com.au

Where to go for further information

Pharmaceutical companies are not in a position to give people an individual diagnosis or medical advice. Your doctor or pharmacist is the best person to give you advice on the treatment of your condition.

This leaflet was prepared in April 2024.