

# APOHEALTH Sinus & Pain Relief Tablets

## Consumer Medicine Information

### What is in this leaflet

This leaflet answers some common questions about APOHEALTH Sinus & Pain Relief.

It does not contain all the available information. It does not take the place of talking to your pharmacist or doctor.

All medicines have risks and benefits. Your pharmacist or doctor has weighed the risks of you taking **APOHEALTH Sinus & Pain Relief** against the benefits they expect it will have for you.

**If you have any concerns about taking this medicine, ask your pharmacist or doctor.**

**Keep this leaflet with the medicine.**

You may need to read it again.

### What APOHEALTH Sinus & Pain Relief is used for

**APOHEALTH Sinus & Pain Relief** contains paracetamol and pseudoephedrine as active ingredients and is used to provide effective temporary relief without drowsiness from sinusitis symptoms:

- Headache
- Nasal congestion
- Sinus congestion
- Sinus Pain

Paracetamol works to stop the pain messages from getting through to

the brain. It also acts in the brain to reduce fever.

Pseudoephedrine belongs to a group of medicines called sympathomimetic decongestants. It works by reducing congestion in the upper respiratory tract, including the nose, nasal passages and sinuses, and making it easier to breathe.

**Ask your pharmacist or doctor if you have any questions about this medicine.**

Your pharmacist or doctor may have given it for another reason.

APOHEALTH Sinus & Pain Relief is only available from your pharmacist.

### Before you take/give APOHEALTH Sinus & Pain Relief Tablets

*When you must not take it*

**Do not take this medicine if you have an allergy to:**

- any medicine containing paracetamol or pseudoephedrine
- any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin

**Do not take this medicine if you have:**

- very high blood pressure
- severe coronary artery disease (heart disease caused by poor blood flow or narrowing of the blood vessels of the heart)
- taken monoamine oxidase inhibitors, medicines used to treat depression, in the last 14 days

**Do not take this medicine if you are pregnant or plan to become pregnant.**

It may affect your developing baby if you take it during pregnancy. Your pharmacist or doctor will discuss the benefits and possible risks of taking the medicine during pregnancy.

**Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.**

If it has expired or is damaged, return it to your pharmacist for disposal.

**If you are not sure whether you should start taking this medicine, talk to your pharmacist or doctor.**

*Before you start to take it*

**Tell your pharmacist or doctor if you have allergies to any other medicines, foods, preservatives or dyes.**

**Tell your pharmacist or doctor if you have or have had any of the following medical conditions:**

- liver or kidney disease
- high blood pressure
- overactive thyroid gland
- diabetes

- heart disease and poor blood flow in the blood vessels of the heart
- glaucoma (high pressure in the eyes)
- prostate problems

**Ask your pharmacist or doctor about taking this medicine if you are breastfeeding.**

Your pharmacist or doctor will discuss the potential benefits and risks of taking the medicine if you are breastfeeding.

**If you have not told your pharmacist or doctor about any of the above, tell him/her before you start taking APOHEALTH Sinus & Pain Relief.**

*Taking other medicines*

**Tell your pharmacist or doctor if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.**

Some medicines may interfere with each other. These include:

- warfarin, a medicine used to prevent blood clots
- metoclopramide, a medicine used to control nausea and vomiting
- medicines used to treat epilepsy or fits
- chloramphenicol, an antibiotic used to treat ear and eye infections
- alcohol
- medicines used to treat depression
- medicines used to treat heart conditions
- medicines used to treat high blood pressure
- medicines used to treat urinary tract infections and bladder problems
- medicines used to treat behavioural disorders
- phenylephrine, a medicine used to treat congestion
- appetite suppressants

These medicines may be affected by APOHEALTH Sinus & Pain Relief or may affect how well it works. You may need different amounts of your medicine, or you may need to take different medicines.

Your doctor and pharmacist will have more information on these and other medicines to be careful with or avoid while taking this medicine.

---

## How to take APOHEALTH Sinus & Pain Relief Tablets

---

**Follow all directions given to you by your pharmacist or doctor carefully.**

They may differ from the information contained in this leaflet.

**If you do not understand the instructions on the box/bottle, ask your pharmacist or doctor for help.**

*How much to take*

**Adults and children over 12 years:**

2 tablets (maximum 8 tablets in 24 hours).

**Children 7 – 12 years:**

1 tablet (maximum 4 tablets in 24 hours).

Keep to the recommended dose.

Not recommended to children under 7 years of age.

**Do not take more than the recommended dose.**

*How to take it*

Tablets should be swallowed with a glass of water. Tablets can be taken with or without food.

*When to take it*

Every 6 hours as necessary.

*How long to take it:*

**Adults: Only take paracetamol for a few days at a time unless your**

**doctor tells you to take it for longer.**

**Children: Only give paracetamol to children for up to 48 hours unless a doctor has told you to give it longer.**

*If you forget to take it*

**If it is less than 1 hour before your next dose, skip the dose you missed and take your next dose when you are meant to.**

**Otherwise, take it as soon as you remember, and then go back to taking your medicine as you would normally.**

**Do not take a double dose to make up for the dose that you missed.**

This may increase the chance of you getting an unwanted side effect.

**If you are not sure what to do, ask your pharmacist or doctor.**

**If you have trouble remembering to take your medicine, ask your pharmacist for some hints.**

*If you take too much (overdose)*

**Immediately telephone your doctor or the Poisons Information Centre (telephone 13 11 26) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much APOHEALTH Sinus & Pain Relief tablets. Do this even if there are no signs of discomfort or poisoning.**

You may need urgent medical attention because of the risk of delayed, serious liver damage.

---

## While you are using APOHEALTH Sinus & Pain Relief Tablets

---

*Things you must do*

**Talk to your pharmacist or doctor if your symptoms do not improve.**

Your pharmacist or doctor will assess your condition and decide if you should continue to take the medicine.

### *Things you must not do*

**Do not take APOHEALTH Sinus & Pain Relief to treat any other complaints unless your pharmacist or doctor tells you to.**

**Do not give your medicine to anyone else, even if they have the same condition as you.**

**Do not take more than the recommended dose unless your pharmacist or doctor tells you to.**

### *Things to be careful of*

**Be careful driving or operating machinery until you know how APOHEALTH Sinus & Pain Relief affects you.**

This medicine may cause dizziness in some people. If this happens, do not drive or use operate machinery.

### *Things you must not do*

**Children: Do not give paracetamol for more than 48 hours unless a doctor has told you to.**

**Adults: Do not take for more than a few days at a time unless your doctor tells you to.**

**Do not take more than the recommended dose unless your doctor tells you to.**

**Do not take APOHEALTH Sinus & Pain Relief to treat any other complaints unless your pharmacist or doctor tells you to.**

**Do not give your medicine to anyone else, even if they have the same condition as you.**

### *Things to be careful of*

**Only drink small quantities of alcohol (beer, wine or spirits) while taking APOHEALTH Sinus & Pain Relief.**

Drinking large quantities of alcohol while taking this medication may increase the risk of liver side effects.

---

## Side effects

---

**Tell your pharmacist or doctor as soon as possible if you do not feel well while you are taking APOHEALTH Sinus & Pain Relief.**

This medicine helps most people with the relief of sinus pain and congestion, but it may have unwanted side effects in a few people.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

**Do not be alarmed by the following lists of side effects. You may not experience any of them.**

**Ask your pharmacist or doctor to answer any questions you may have.**

Tell your pharmacist or doctor if you notice any of the following and they worry you:

- nausea or dyspepsia
- drowsiness
- difficulty sleeping
- nervousness
- excitability
- restlessness
- dizziness
- fear or anxiety
- rapid heart beat
- tremor
- hallucinations

The above list includes the more common side effects of your medicine. They are usually mild and short-lived.

Children and people over 65 years of age may have an increased chance of getting side effects.

**If any of the following happen, tell your doctor immediately or go to Accident and Emergency at your nearest hospital:**

- urine retention
- shortness of breath
- wheezing or difficulty breathing

- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin
- sudden abdominal pain, rectal bleeding

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation.

**These side effects are very rare.**

**Tell your pharmacist or doctor if you notice anything that is making you feel unwell.**

Other side effects not listed above may also occur in some people.

---

## After using APOHEALTH Sinus & Pain Relief Tablets

---

### *Storage*

**Keep your medicine in the original pack until it is time to take.**

**Keep your tablets in a cool dry place where the temperature stays below 30°C.**

Do not store APOHEALTH Sinus & Pain Relief or any other medicine in the bathroom or near a sink.

**Do not leave it on a window sill or in the car.**

Heat and dampness can destroy some medicines.

**Keep it where children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

### *Disposal*

**Ask your pharmacist what to do with any medicine that is left over, or if the expiry date has passed.**

---

## Product description

---

### *What it looks like*

**APOHEALTH Sinus & Pain Relief** are capsule shaped, pink, uncoated tablets.

### *Ingredients*

APOHEALTH Sinus & Pain Relief contains 500 mg of paracetamol and 30 mg of pseudoephedrine hydrochloride as the active ingredients.

It also contains the following inactive ingredients:

- Microcrystalline cellulose
- Maize starch
- Colloidal anhydrous silica
- Povidone
- Croscarmellose sodium
- Stearic acid
- Magnesium stearate
- Sodium starch glycollate
- Self-emulsifying glyceryl monostearate
- Erythrosine ci45430 (127)

**This medicine does not contain lactose, sucrose, gluten and tartrazine.**

### *Supplier*

Arrotex Pharmaceuticals Pty Ltd  
15 – 17 Chapel Street  
Cremorne Victoria 3121

APOHEALTH Sinus & Pain Relief  
AUST R 353481

This leaflet was updated in March 2024.