

ARGIPRESSIN MEDSURGE

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I using ARGIPRESSIN MEDSURGE?

ARGIPRESSIN MEDSURGE contains the active ingredient argipressin. ARGIPRESSIN MEDSURGE is used to treat deficiency of argipressin in diabetes insipidus. Argipressin can also be to prevent or relieve stomach bloating after you have had surgery or a radiation procedure of the abdomen.

For more information, see Section [1. Why am I using ARGIPRESSIN MEDSURGE?](#) in the full CMI.

2. What should I know before I use ARGIPRESSIN MEDSURGE?

Do not use if you have had an allergic reaction to argipressin or any of the ingredients listed at the end of the CMI. Do not use if you have chronic kidney problems which has caused accumulation of nitrogen in the blood.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.

For more information, see Section [2. What should I know before I use ARGIPRESSIN MEDSURGE?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with ARGIPRESSIN MEDSURGE and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I use ARGIPRESSIN MEDSURGE?

- ARGIPRESSIN MEDSURGE is for injection into a muscle or under the skin, or through the nostrils using a spray, cotton wad (pledgets) or a dropper
- The usual dose for diabetes insipidus is 0.25 mL to 0.5 mL repeated two or three times a day. Your doctor will tell you how much ARGIPRESSIN MEDSURGE to use

More instructions can be found in Section [4. How do I use ARGIPRESSIN MEDSURGE?](#) in the full CMI.

5. What should I know while using ARGIPRESSIN MEDSURGE?

Things you should do	<ul style="list-style-type: none">• If you become pregnant while using ARGIPRESSIN MEDSURGE, tell your doctor immediately• Remind any doctor, dentist, pharmacist, surgeon or anaesthetist you visit that you are using ARGIPRESSIN MEDSURGE
Things you should not do	<ul style="list-style-type: none">• Do not stop using this medicine suddenly or change the dosage without checking with your doctor
Driving or using machines	<ul style="list-style-type: none">• Be careful before you drive or use any machines until you know how ARGIPRESSIN MEDSURGE affects you
Drinking alcohol	<ul style="list-style-type: none">• Be careful when drinking alcohol while taking ARGIPRESSIN MEDSURGE. Alcohol may decrease the effects of ARGIPRESSIN MEDSURGE
Looking after your medicine	<ul style="list-style-type: none">• Keep ARGIPRESSIN MEDSURGE in the refrigerator (2-8°C), in a dark place, protected from light, where children cannot reach it• Do not leave it in a car, on window sills, near a sink or in the bathroom.• Keep ARGIPRESSIN MEDSURGE in its sealed vial until it is time to use it

For more information, see Section [5. What should I know while using ARGIPRESSIN MEDSURGE?](#) in the full CMI.

6. Are there any side effects?

If you get any of the following side effects, tell your doctor immediately or go to Accident and Emergency at your nearest hospital: chest pain, confusion, difficulty breathing or wheezing, difficulty passing urine, drowsiness, listlessness, pain/redness/irritation or swelling at the injection site, rapid weight gain, seizures (convulsions), severe or continuous headaches, skin rash or itching (hives).

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

ARGIPRESSIN MEDSURGE (Pi-TRES-sin)

Active ingredient: *argipressin*

Consumer Medicine Information (CMI)

This leaflet provides important information about using ARGIPRESSIN MEDSURGE. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using ARGIPRESSIN MEDSURGE.**

All medicines have risks and benefits. Your doctor has weighed the possible risks of taking ARGIPRESSIN MEDSURGE against the benefits they expect it will have for you.

Keep this leaflet with the medication. You may want to read it again.

Where to find information in this leaflet:

[1. Why am I using ARGIPRESSIN MEDSURGE?](#)

[2. What should I know before I use ARGIPRESSIN MEDSURGE?](#)

[3. What if I am taking other medicines?](#)

[4. How do I use ARGIPRESSIN MEDSURGE?](#)

[5. What should I know while using ARGIPRESSIN MEDSURGE?](#)

[6. Are there any side effects?](#)

[7. Product details](#)

1. Why am I using ARGIPRESSIN MEDSURGE?

ARGIPRESSIN MEDSURGE contains the active ingredient argipressin or antidiuretic hormone. Argipressin is normally produced by the pituitary gland.

Argipressin acts on blood vessels and the kidney and reduces the amount of fluids you pass in the urine. Argipressin also causes muscles called "smooth muscles" to contract.

If your body does not produce enough argipressin you can get a condition called "diabetes insipidus".

ARGIPRESSIN MEDSURGE is used to treat deficiency of argipressin in diabetes insipidus. Symptoms of diabetes insipidus include passing large volumes of pale urine and feeling extremely thirsty. Diabetes insipidus is not the same as diabetes mellitus where sugar is passed in the urine.

Argipressin can also be used to prevent or relieve stomach bloating after you have had surgery or a radiation procedure of the abdomen.

Ask your doctor if you have any questions why this medicine has been prescribed for you.

Your doctor may have prescribed it for another reason.

2. What should I know before I use ARGIPRESSIN MEDSURGE?

Warnings

Do not use ARGIPRESSIN MEDSURGE if:

- you have chronic kidney problems which has caused accumulation of nitrogen in the blood
- you have an allergy to argipressin, or any of the ingredients listed at the end of this leaflet
- Always check the ingredients to make sure you can use this medicine.

Check with your doctor if you:

- **have allergies to any other medicines, foods, preservatives or dyes.**
- **have or have had any of the following medical conditions:**
 - coronary artery disease. ARGIPRESSIN MEDSURGE may bring on angina pain or heart problems
 - asthma
 - blood vessel (vascular) disease
 - swelling of thyroid glands with heart problems
 - heart disease
 - kidney disease associated with high blood pressure
 - migraine
 - epilepsy or seizures (convulsions)
 - complications of pregnancy such as pre-eclampsia (high blood pressure)

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

If you become pregnant while using ARGIPRESSIN MEDSURGE, tell your doctor immediately.

It is not known whether ARGIPRESSIN MEDSURGE will harm your baby.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

It is not known whether ARGIPRESSIN MEDSURGE will affect your baby.

3. What if I am taking other medicines?

Tell your doctor, nurse or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and ARGIPRESSIN MEDSURGE may interfere with each other. These include:

- carbamazepine, a medicine used to treat epilepsy or convulsions
- tricyclic antidepressants such as amitriptyline, clomipramine, dosulepin, doxepin, imipramine and nortriptyline
- heparin, a medicine used to thin the blood
- fludrocortisone, a medicine used to treat Addison's disease or conditions of insufficiency of body salt.
- noradrenaline, a medicine used to treat very low blood pressure
- urea, a medicine used to treat fluid retention (oedema)
- demeclocycline, a medicine used to treat certain infections
- lithium, a medicine used to treat certain mood disorders
- H₂ blockers that are used to treat stomach ulcers, such as famotidine and nizatidine

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect ARGIPRESSIN MEDSURGE.

4. How do I use ARGIPRESSIN MEDSURGE?

How much to use

- The usual dose for diabetes insipidus is 0.25 mL to 0.5 mL repeated two or three times a day as instructed by your doctor
- **Do not use more or less than the doctor has prescribed**
- The above doses are for the use of ARGIPRESSIN MEDSURGE in diabetes insipidus only. If you are given ARGIPRESSIN MEDSURGE for another condition your doctor or nurse will give you this medicine in a hospital or clinic.

How to use ARGIPRESSIN MEDSURGE

- ARGIPRESSIN MEDSURGE is for injection into a muscle (intramuscular) or under the skin (subcutaneous), or through the nostrils using a spray, cotton wad (pledgets) or a dropper.
- If you have been instructed to give yourself ARGIPRESSIN MEDSURGE, make sure your doctor or pharmacist shows you how to use this medicine.
- Always use exactly as directed by your doctor.

Continue to use ARGIPRESSIN MEDSURGE for as long as your doctor tells you. This medicine helps you control your condition, but does not cure it. It is important to keep using your medicine even if you feel well.

If you forget to use ARGIPRESSIN MEDSURGE

If you miss your dose at the usual time, and **if it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.**

Otherwise use it as soon as you remember, and then go back to using your medicine as you would normally.

Do not take a double dose to make up for the dose you missed.

If you are not sure what to do, ask your doctor or pharmacist.

If you use too much ARGIPRESSIN MEDSURGE

If you think that you or anyone else may have used too much ARGIPRESSIN MEDSURGE, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre (by calling 13 11 26 or New Zealand 0800 POISON or 0800 764 766) for advice, or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital

You should do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

5. What should I know while using ARGIPRESSIN MEDSURGE?

Things you should do

Remind any doctor, dentist or pharmacist you visit that you are using ARGIPRESSIN MEDSURGE.

Tell your doctor immediately if you:

- Become pregnant while using ARGIPRESSIN MEDSURGE

Things you should not do

- Do not stop taking ARGIPRESSIN MEDSURGE, or lower the dosage, even if you are feeling better, without checking with your doctor

Driving or using machines

Be careful before you drive or use any machines or tools until you know how ARGIPRESSIN MEDSURGE affects you.

Drinking alcohol

Tell your doctor if you drink alcohol.

Be careful when drinking alcohol while taking ARGIPRESSIN MEDSURGE. Alcohol may decrease the effects of ARGIPRESSIN MEDSURGE.

Looking after your medicine

Keep ARGIPRESSIN MEDSURGE:

- in its sealed vial until it is time to use it

- in the refrigerator (2-8°C)

Follow the instructions in the carton on how to take care of your medicine properly.

Store it in a cool dry place away from moisture, heat or sunlight; for example, do not store it:

- In the bathroom or near a sink, or
- In the car or on window sills.

Keep it where young children cannot reach it.

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Less serious side effects

Less serious side effects	What to do
<p>Gut and digestion-related:</p> <ul style="list-style-type: none"> • belching, stomach gas • nausea • vomiting • stomach pain or cramps <p>Heart related:</p> <ul style="list-style-type: none"> • palpitations <p>Breathing and Allergy related:</p> <ul style="list-style-type: none"> • difficulty sleeping (insomnia) • dry mouth, nose or throat • itching or skin rash • injection site redness or pain <p>Other:</p> <ul style="list-style-type: none"> • dizziness • sweating • trembling • facial flushing with a pale area around the mouth • headache 	<p>Speak to your doctor if you have any of these less serious side effects and they worry you.</p>

Serious side effects

Serious side effects	What to do
<p>Signs of an allergic reaction such as:</p> <ul style="list-style-type: none"> • difficulty breathing, wheezing • pain, redness irritation or swelling at the injection site • skin rash, itching (hives) 	<p>Call your doctor straight away, or go straight to the Emergency Department at your nearest</p>

<p>Signs of fluid overload:</p> <ul style="list-style-type: none"> • drowsiness • severe or continuous headaches • listlessness Seizures (convulsions) <p>Other</p> <ul style="list-style-type: none"> • chest pain • difficulty passing urine • rapid weight gain • confusion 	<p>hospital if you notice any of these serious side effects.</p>
---	---

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems (in Australia) or <https://nzphvc.otago.ac.nz/consumer-reporting/> (in New Zealand). By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What ARGIPRESSIN MEDSURGE contains

Active ingredient (main ingredient)	argipressin
Other ingredients (inactive ingredients)	sodium acetate hydrochloric acid sodium Hydroxide water for Injections

Do not take this medicine if you are allergic to any of these ingredients.

The argipressin in ARGIPRESSIN MEDSURGE is synthetic and not of animal origin.

What ARGIPRESSIN MEDSURGE looks like

ARGIPRESSIN MEDSURGE is a clear colourless solution sealed in 1 mL glass vials. (AUST R 449846).

Who distributes ARGIPRESSIN MEDSURGE

Medsurge Healthcare Pty Limited
Unit 1& 2, 6-7 Gilda Court
MULGRAVE VICTORIA 3170
Tel: 1300 788 261
www.medsurgehc.com

This leaflet was prepared in Aug 2025.