

# ACTAZE Tablets 15 mg, 30 mg & 45 mg

*pioglitazone (as hydrochloride) Tablets*

---

## Consumer Medicine Information

---

### What is in this leaflet

---

This leaflet answers some common questions about ACTAZE Tablets. It does not contain all the available information and does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits.

Your doctor has weighed the risks of you taking ACTAZE Tablets against the benefits they expect it will have for you.

**If you have any concerns about taking this medicine, ask your doctor or pharmacist.**

**Keep this leaflet with the medicine.**

You may need to read it again.

---

### What ACTAZE Tablets are used for

---

ACTAZE Tablets are used to improve the action of the body's naturally produced insulin.

ACTAZE Tablets are used in the management of type 2 diabetes not controlled by diet.

ACTAZE Tablets help to control the level of glucose in your blood when you have type 2 diabetes. This is the 'adult onset' type of diabetes and is controlled by diet, certain oral medications and occasionally insulin.

This medicine is also called pioglitazone hydrochloride and belongs to a group of medicines

called glitazones. Glitazones decrease insulin resistance.

**Ask your doctor if you have any questions about why this medicine has been prescribed for you.**

Your doctor may have prescribed it for another reason.

ACTAZE Tablets can be used alone (when diet and exercise is not enough to treat your diabetes) or together with other anti-diabetic medicines.

This medicine is available only with a doctor's prescription.

The use of ACTAZE Tablets has not been studied in children.

---

### Before you take ACTAZE Tablets

---

*When you must not take it*

**Do not take ACTAZE Tablets if:**

- You have heart failure requiring treatment. Talk to your doctor if you have heart failure
- you have type 1 diabetes or diabetic ketoacidosis (often caused by very high blood glucose levels)
- You have an allergy to any medicine containing pioglitazone hydrochloride or any of the ingredients listed at the end of this leaflet (see 'Product Description').

Some of the symptoms of an allergic reaction may include shortness of breath, wheezing or difficulty breathing; swelling of the

face, lips, tongue or other parts of the body; rash, itching or hives on the skin.

**Do not take this medicine after the date printed on the pack or if the packaging is torn or shows signs of tampering.**

If it has expired or is damaged, return it to your pharmacist for disposal.

**If you are not sure whether you should start taking this medicine, talk to your doctor.**

*Before you start to take it*

**Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.**

**Tell your doctor if you have or have had any of the following medical conditions:**

- Heart disease with shortness of breath after minimal physical activity
- Heart disease with severe symptoms at rest
- Swelling of hands, ankles or feet.
- Bladder cancer or symptoms associated with bladder cancer such as blood in the urine (hematuria) often accompanied by pain and burning or sudden urges to urinate.
- Problems with your liver.
- Problems with your kidneys that require dialysis. ACTAZE Tablets are not recommended for use if you are on dialysis.
- Some women who do not have monthly periods and have not been through menopause may

restart their periods when taking ACTAZE Tablets. These women may be at increased risk of pregnancy.

- Bone fractures, usually in the hand, upper arm or foot, have been seen in some women when taking ACTAZE Tablets. Talk to your doctor for advice on how to keep your bones healthy.

**Tell your doctor if you are pregnant or plan to become pregnant or are breast-feeding.**

Like most medicines, ACTAZE Tablets are not recommended for use during pregnancy. If there is a need to consider ACTAZE Tablets during your pregnancy, your doctor will discuss with you the benefits and risks of taking ACTAZE Tablets. It is recommended that you do not breast-feed while taking ACTAZE Tablets, as it is not known whether ACTAZE Tablets passes into breast milk.

**Tell your doctor if you are using another medicine for diabetes. ACTAZE Tablets can enhance the action of other medicines. You may be at risk of low blood sugar (hypoglycaemia).**

If this happens, your doctor may need to adjust the dose of your other medicines.

**Tell your doctor if you suffer from lactose intolerance (because ACTAZE Tablets contain lactose).**

**If you have not told your doctor about any of the above, tell him/her before you start taking ACTAZE Tablets.**

*Taking other medicines*

**Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.**

Some medicines and ACTAZE Tablets may interfere with each other. These include:

- chlorpropamide
- gemfibrozil

- glibenclamide
- gliclazide
- insulin
- oral contraceptives
- rifampicin
- tolbutamide

These medicines may be affected by ACTAZE Tablets or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines.

Your doctor or pharmacist has more information on medicines to be careful with or to avoid while taking this medicine.

---

## How to take ACTAZE Tablets

---

**Follow all directions given to you by your doctor or pharmacist carefully.**

They may differ from the information contained in this leaflet.

**If you do not understand the instructions on the carton, ask your doctor or pharmacist for help.**

*How much to take*

Your doctor will tell you how many ACTAZE Tablets you should take. The dose your doctor will prescribe for you will usually be in the range of 15 mg to 45 mg per day.

**ACTAZE Tablets should be taken once a day as advised by your doctor.**

Your doctor may increase or decrease your dose in order to find the appropriate dose for your condition.

*How to take it*

**ACTAZE Tablets should be swallowed whole with a glass of water.**

*When to take it*

**Take your medicine at about the same time each day.**

Taking it at the same time each day will have the best effect. It will also help you remember when to take it.

It does not matter if you take this medicine before or after food.

*How long to take it*

**Continue taking your medicine for as long as your doctor tells you.**

This medicine helps to control your condition, but does not cure it. It is important to keep taking your medicine even if you feel well.

*If you forget to take it*

**If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.**

**Otherwise, take it as soon as you remember, and then go back to taking your medicine as you would normally.**

**Do not take a double dose to make up for the dose that you missed.**

**If you are not sure what to do, ask your doctor or pharmacist.**

**If you have trouble remembering to take your medicine, ask your pharmacist for some hints.**

---

## If you take too much (overdose)

---

Immediately telephone the Australian Poisons Information Centre (telephone 13 11 26) or the New Zealand National Poisons Information Centre (0800 POISON or 0800 764 766) or go to Accident and Emergency at your nearest hospital, if you think that you or anyone else may have taken too much ACTAZE Tablets. Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

---

## While you are taking ACTAZE Tablets

---

### *Things you must do*

It is important that you remember to take ACTAZE Tablets daily and at the dose prescribed by your doctor.

Tell any other doctors, dentists and pharmacists who treat you that you are taking ACTAZE Tablets

If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking ACTAZE Tablets.

If you are going to have surgery, tell the surgeon or anaesthetist that you are taking this medicine.

It may affect other medicines used during surgery.

If you become pregnant while taking this medicine, tell your doctor immediately.

If you are about to have any blood tests, tell your doctor that you are taking this medicine.

It may interfere with the results of some tests.

Keep all of your doctor's appointments so that your progress can be checked.

Your doctor may do some tests from time to time to make sure the

medicine is working and to prevent unwanted side effects.

**Tell your doctor if you have gained weight since taking ACTAZE Tablets.**

Weight gain can be associated with improved blood sugar control however; it may also be a symptom of heart failure.

### *Things you must not do*

**Do not take ACTAZE Tablets to treat any other complaints unless your doctor tells you to.**

**Do not give your medicine to anyone else, even if their symptoms seem similar to yours or they have the same condition as you.**

**Do not stop taking your medicine or lower the dosage without checking with your doctor.**

If you stop taking it suddenly, your condition may worsen or you may have unwanted side effects.

### *Things to be careful of*

**ACTAZE Tablets alone are unlikely to affect your ability to drive or operate machinery. However, be careful to avoid hypoglycaemia whilst driving or operating machinery if using ACTAZE Tablets in combination with other diabetes medicines.**

---

## Side effects

---

**Tell your doctor or pharmacist as soon as possible if you experience any undesirable effect or feel unwell while you are taking ACTAZE Tablets.**

This medicine helps most people with type 2 diabetes not controlled by diet, but it may have unwanted side effects in some people.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you experience some of the side effects.

Some side effects may be related to the dose of ACTAZE Tablets. Accordingly, it is important that you tell your doctor as soon as possible about any unwanted effects. Your doctor may then decide to adjust the dose of ACTAZE Tablets you are taking.

**Do not be alarmed by the following list of side effects. You may not experience any of them.**

**Ask your doctor or pharmacist to answer any questions you may have.**

A few patients have experienced the following side effects whilst taking ACTAZE Tablet:

- a small increase in weight
- low blood sugar levels (hypoglycaemia). This occurs more often when ACTAZE Tablets are taken with a sulfonylurea or insulin
- heart failure which may show as localised swelling of the ankles, feet and hands (oedema) and/or fluid in the lungs (pulmonary oedema). This has been reported in clinical trials mainly in patients who are taking ACTAZE Tablets in combination with insulin
- Increased risk of fracture in women
- macular oedema (an eye disorder that can affect vision)
- altered or impaired liver function.

**Tell your doctor if you notice any of the following side effects:**

- Weight gain.
- Signs of hypoglycaemia which may include weakness, trembling or shaking, sweating, light-headedness, headache, dizziness, lack of concentration, tearfulness or crying, irritability, hunger, numbness around the lips and fingers.
- Eye problems including blurred or double vision.

The above list includes the more common side effects of your medicine.

**Tell your doctor as soon as possible if you notice any of the following:**

- Dark urine or pale stools, yellowing of the skin or eyes, severe cramps of the stomach, nausea or vomiting, loss of weight, tiredness.
- Shortness of breath when at rest or after minimal physical activity with swelling of legs, feet and hands, rapid increase in weight.
- Blood in the urine often accompanied by pain and burning, these can be symptoms of bladder cancer.

The above list includes serious side effects, which may require medical attention. Serious side effects are rare.

**Tell your doctor or pharmacist if you notice anything that is making you feel unwell.**

Other side effects not listed above may also occur in some people.

---

## After using ACTAZE Tablets

---

### *Storage*

**Keep your tablets in the aluminium blister pack until it is time to take them.**

**Keep your tablets in a cool, dry place below 25°C.**

**Do not store ACTAZE Tablets or any other medicine in the bathroom or near a sink. Do not leave it on a window sill or in the car.**

Heat and dampness can destroy some medicines.

**Keep it where children cannot reach it.**

A locked cupboard at least 1½ meters above the ground is a good place to store medicines.

### *Disposal*

**If your doctor tells you to stop taking ACTAZE Tablets or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.**

---

## Product description

---

### *What it looks like*

ACTAZE Tablets are available in three strengths.

#### **ACTAZE Tablets 15 mg**

AUST R 254272

White to off white, round, biconvex, uncoated tablets debossed with 'P' on one side and '15' on other side.

#### **ACTAZE Tablets 30 mg**

AUST R 254292

White to off white, round, flat, uncoated tablets with bevelled edges debossed with 'PIO' on one side and '30' on the other side.

#### **ACTAZE Tablets 45 mg**

AUST R 254312

White to off white, flat, round uncoated tablets with bevelled edges debossed with 'PIO' on one side and '45' on the other side.

ACTAZE Tablets are presented in aluminium blister packs containing 7 (sample pack) and 28 tablets.

### *Ingredients*

ACTAZE Tablets contain 15 mg, 30 mg and 45 mg of pioglitazone as the active ingredient.

It also contains:

- lactose monohydrate
- carmellose calcium
- hypromellose
- magnesium stearate

This medicine does not contain sucrose, gluten, tartrazine or any other azo dyes. This medicine contains sugars as lactose.

### *Name and Address of the Sponsor*

Arrow Pharma Pty Ltd  
15-17 Chapel Street  
Cremorne VIC 3121

### *Date of first inclusion in the ARTG*

30 September 2015

This leaflet was revised in April 2022.