

ATTORA LA

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

WARNING: Important safety information is provided in a boxed warning in the [full CMI](#). Read before using this medicine.

1. Why am I taking ATTORA LA?

ATTORA LA contains the active ingredient methylphenidate hydrochloride. ATTORA LA capsules are used to treat Attention Deficit Hyperactivity Disorder (ADHD). For more information, see Section [1. Why am I taking ATTORA LA?](#) in the full CMI.

2. What should I know before I take ATTORA LA?

Do not take if you are allergic to methylphenidate or to any of the other ingredients listed at the end of the CMI. Do not take ATTORA LA if you are taking a medicine called a monoamine oxidase inhibitor (MAOI) or have been taking it within the past 14 days.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding. For more information, see Section [2. What should I know before I take ATTORA LA?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with ATTORA LA and affect how it works. A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I take ATTORA LA?

- You can take ATTORA LA with or without food. Swallow the capsules whole as a single dose in the morning with a full glass of water. Do not crush or chew the capsules.
- Follow all directions given to you by your doctor carefully.

More instructions can be found in Section [4. How do I take ATTORA LA?](#) in the full CMI.

5. What should I know while taking ATTORA LA?

Things you should do	<ul style="list-style-type: none">• Remind any doctor, dentist or pharmacist you visit that you are taking ATTORA LA.• Take ATTORA LA exactly as your doctor has prescribed.• Keep all of your doctor's appointments so that your or your child's progress can be checked.• Tell your doctor if you are pregnant or breast-feeding.
Things you should not do	<ul style="list-style-type: none">• Do not stop your treatment without first checking with your doctor.• Do not change the dose without talking to the doctor.• Do not take this medicine to treat any other complaints unless your doctor tells you to.• Do not give it to anyone else, even if their condition seems similar to yours.
Driving or using machines	<ul style="list-style-type: none">• Be careful while driving, operating machinery or doing jobs that require you to be alert while you are taking ATTORA LA until you know how it affects you.
Drinking alcohol	<ul style="list-style-type: none">• Do not drink alcohol whilst you are taking ATTORA LA.
Looking after your medicine	<ul style="list-style-type: none">• Keep your medicine in the original container until it is time to take a dose.• Store it in a cool, dry place.• Do not store it or any medicine in the bathroom or near a sink. Do not leave it on a windowsill or in the car.

For more information, see Section [5. What should I know while taking ATTORA LA?](#) In the full CMI.

6. Are there any side effects?

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects. For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

WARNING: DRUG DEPENDENCE: ATTORA LA should be used cautiously by patients with a history of drug dependence or alcoholism. Ongoing overuse of ATTORA LA can lead to tolerance and can affect mental wellbeing in various degrees of severity. Withdrawal of ATTORA LA overuse should only take place under careful supervision. If ATTORA LA is being withdrawn after being used for its intended purposes, underlying symptoms of the disorder might be noticed and may require follow up by your doctor.

ATTORA LA

Active ingredient(s): *methylphenidate hydrochloride*

Consumer Medicine Information (CMI)

This leaflet provides important information about taking ATTORA LA. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about taking ATTORA LA.**

Where to find information in this leaflet:

- [1. Why am I taking ATTORA LA?](#)
- [2. What should I know before I take ATTORA LA?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I take ATTORA LA?](#)
- [5. What should I know while taking ATTORA LA?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am I taking ATTORA LA?

ATTORA LA contains the active ingredient methylphenidate hydrochloride. Methylphenidate hydrochloride is a central nervous system stimulant.

ATTORA LA (long-acting) capsules are used to treat Attention Deficit Hyperactivity Disorder (ADHD).

ATTORA LA is a stimulant that increases attention and decreases impulsiveness and hyperactivity in patients with ADHD.

ATTORA LA is thought to work by regulating specific chemicals in the brain that affect behaviour. It helps to focus attention, shut out distraction and allows impulsive people to think before they act. If successful, it will enhance an inattentive person's natural ability.

ATTORA LA should be given as part of a total treatment program for ADHD that may include other measures (psychological, educational and social). It is not intended for use in patients who have symptoms due to environmental factors and/or other primary psychiatric disorders, including psychosis.

Ask your doctor if you have any questions about why this medicine has been prescribed for you or your child. Your doctor may have prescribed it for another purpose.

ATTORA LA, like all medicines containing central nervous system stimulants, will be given to you only under close medical supervision and after diagnosis.

2. What should I know before I take ATTORA LA?

Warnings

Do not take ATTORA LA if:

- you are allergic to methylphenidate, or any of the ingredients listed at the end of this leaflet.
- Always check the ingredients to make sure you can use this medicine.

Some of the symptoms of an allergic reaction may include shortness of breath, wheezing or difficulty breathing; swelling of the face, lips, tongue or other parts of the body; rash, itching or hives on the skin.
- you have periods of severe anxiety, tension or agitation
- you have Tourette's syndrome (a condition with uncontrolled speech and body movements or tics) or you have a family history of this disorder
- you have tics (muscle twitching which is usually in the face or shoulders) or if your brothers or sisters have tics
- you have increased pressure in the eye (glaucoma)
- you have an overactive thyroid (hyperthyroidism) or other thyroid problems
- you have heart problems such as heart attack, irregular heartbeat, chest pain (angina), heart failure, heart disease or if you were born with a heart problem
- you have very high blood pressure (hypertension) or narrowing of the blood vessels (arterial occlusive disease, which can cause pain in the arms and legs)
- you have severe depression or other mental illness
- you have a tumour of the adrenal gland, which sits near the kidney (pheochromocytoma)
- you have a history of alcohol or drug abuse or dependence

Ask your doctor or pharmacist if you are not sure whether any of the above medical conditions apply to you.

Do not take ATTORA LA if you are taking a medicine called a monoamine oxidase inhibitor (MAOI) or have been taking it within the past 14 days.

MAOI medicines include moclobemide, phenelzine, tranylcypromine and linezolid.

Taking ATTORA LA together with MAOI medicines may cause a serious reaction with a sudden increase in body temperature, extremely high blood pressure and severe convulsions

Ask your doctor or pharmacist if you are not sure if you have been taking one of these medicines.

Check with your doctor if you:

- are allergic to any other medicines, foods, dyes or preservatives. Your doctor will want to know if you are prone to allergies.
- have any of the following medical conditions or behaviours:
 - any heart defects (e.g. structural cardiac abnormality)
 - a family history of sudden death and /or irregular heart beat
 - hardening of the arteries
 - any other current or previous heart problems
 - any disorders of the blood vessels in the brain, e.g. weakening of the blood vessel (aneurysm), stroke, or inflammation of blood vessels (vasculitis)
 - severe depression, bipolar disorder or other mental illness
 - epilepsy (seizures, convulsions, or fits)
 - high blood pressure
 - history of alcohol or drug abuse or dependence
 - acute mental disorders that cause abnormal thinking and perceptions (psychosis) or feeling unusually excited, over-active and un-inhibited (acute mania) -your doctor will have told you if you have this
 - psychotic symptoms such as seeing or feeling things that are not really there (hallucinations)
 - aggressive behaviour
 - suicidal thoughts or behaviour
 - fingers and toes feeling numb, tingling and changing colour (from white to blue, then red) when cold ('Raynaud's phenomenon')
 - have or develop eye problems, including increased pressure in your eye, glaucoma or problems with your close-up vision (farsightedness).

Your doctor may want to take special precautions if you have any of the above conditions.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

Ask your doctor about the risks and benefits of taking ATTORA LA during pregnancy. ATTORA LA is not to be used during pregnancy unless specifically prescribed by your doctor.

This medicine may affect your developing baby if you take it while you are pregnant.

Do not breast-feed during treatment with ATTORA LA.

The active ingredient in ATTORA LA can pass into the breast milk.

Children

There is not enough information to recommend ATTORA LA use in children under 6 years old.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines may interfere with ATTORA LA and affects how it works. These include:

- medicines that increase blood pressure
- alpha 2 agonists like clonidine (used to treat high blood pressure)
- medicines used to treat depression, such as tricyclic antidepressants (imipramine, desipramine) and monoamine oxidase inhibitors (MAO inhibitors)
- some anticonvulsants (medicines used to treat epilepsy or fits e.g. phenobarbitone, primidone, phenytoin)
- oral anticoagulants or warfarin (medicines used to prevent blood clots)
- phenylbutazone (used to treat pain or fever)
- guanethidine
- anaesthetics
- medicines that influence the level of dopamine in the body (dopaminergic medicines used to treat Parkinson's disease or psychosis)
- medicines that raise the level of serotonin in the body (serotonergic medicines, for example those used to treat depression like sertraline and venlafaxine).

You or your child may need to take different amounts of your medicines, or you or your child may need to take different medicines.

If you have not told your doctor about any of these things, tell him/ her before you or your child start taking this medicine.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect ATTORA LA.

4. How do I take ATTORA LA?

How much to take

- ATTORA LA is available in long-acting capsules containing 10 mg, 20 mg, 30 mg, 40 mg or 60 mg of methylphenidate hydrochloride as the active ingredient.
- Your doctor will usually start with a low dose and, if necessary, increase it gradually at weekly intervals up to 20 mg to 40 mg each day.

- Do not exceed the maximum recommended dose
- In children, the maximum recommended daily dose is 60 mg. In adults, the maximum recommended daily dose is 80 mg.

When to take ATTORA LA

- **Take your doses at around the same time in the morning every day.**
If you have trouble remembering when to take your medicine, ask your pharmacist for some hints.

How to take ATTORA LA

- You can take ATTORA LA with or without food. Swallow the capsules whole as a single dose in the morning with a full glass of water. Do not crush or chew the capsules.
- If you have trouble swallowing the capsules, you can carefully open them and sprinkle the contents (little beads) over a small amount of soft food (e.g. applesauce).
The soft food should not be warm as this could affect the special properties of the beads.
- Immediately eat all of the soft food and ATTORA LA mixture. Do not store any of the soft food that has been mixed with ATTORA LA for future use.
- If the capsules upset your stomach, you can take them with food, but always take them in the same way (e.g. always with food or always without food). That way the effect will always be the same.
- Follow all directions given to you by your doctor and pharmacist carefully.
They may differ from the information contained in this leaflet.
If you do not understand the instructions on the label, ask your doctor or pharmacist for help

How long to take ATTORA LA

- Continue taking ATTORA LA for as long as your doctor tells you.
This medicine helps to control your symptoms but does not cure your condition. Your doctor will check your progress to make sure the medicine is working and will discuss with you how long your treatment should continue.
- Treatment for ADHD varies in length from patient to patient. During treatment for ADHD, your doctor may stop ATTORA LA every so often (e.g. over weekends or school holidays) to see whether it is still needed.
Breaks from treatment also help to prevent a slow-down in growth that sometimes happens when children take this medicine for a long time.

If you forget to take ATTORA LA

If you forget to take a dose of ATTORA LA capsules and you remember before mid-day, take the dose as soon as you remember. Then go back to your usual schedule on the following day.

If you do not remember before mid-day, miss your dose of ATTORA LA for that day and wait until the following morning to take your next dose.

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Do not take a double dose to make up for the dose you missed. Your chance of an unwanted side effect may be increased if you do.

If you take too much ATTORA LA

If you think that you or anyone else have taken too much ATTORA LA, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre (**by calling 13 11 26**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

Some of the symptoms of an overdose may include vomiting, agitation, headache, tremors, muscle twitching, irregular heart beat, flushing, fever, sweating, dilated pupils, breathing problems, confusion, seizures and muscle spasms accompanied by fever and red-brown urine.

5. What should I know while taking ATTORA LA?

Things you should do

Take ATTORA LA exactly as your doctor has prescribed.

Like all stimulants, this medicine may become habit-forming and can be abused by some people. If you take it correctly as instructed by your doctor, abuse or dependence should not be a problem, either now or later in life.

Be sure to keep all of your doctor's appointments so that your or your child's progress can be checked.

Your doctor will want to check your or your child's blood pressure, height, weight and do blood tests from time to time to prevent unwanted side effects from happening.

If you become pregnant while taking ATTORA LA, tell your doctor. Your doctor can discuss with you the risks and benefits of taking it while you are pregnant.

Tell any doctor, dentist or pharmacist who treats you that you are taking ATTORA LA.

If your child is not growing or gaining height or weight as expected, treatment with ATTORA LA may need to be interrupted.

If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking ATTORA LA.

Things you should not do

- Do not stop your treatment without first checking with your doctor.
If you suddenly stop taking this medicine, your condition may reappear or you may get unwanted effects such as depression. To prevent this, your

doctor may want to gradually reduce the amount of medicine you take each day before stopping it completely.

- You will need medical supervision after having interrupted the treatment.
- Do not change the dose without talking to the doctor. If you have the impression that the effect of ATTORA LA is too strong or too weak, talk to the doctor.
- Do not take this medicine to treat any other complaints unless your doctor tells you to.
- Do not give it to anyone else, even if their condition seems similar to yours.

Things to be careful of

- ATTORA LA may give a false positive result when testing for drug use. This includes testing used in sport.
- Some children taking ATTORA LA for a long time may have slower than normal growth, but they usually catch up once the treatment is stopped.
- In some patients ATTORA LA may cause stomach upset, loss of appetite and difficulty sleeping, especially at the start of treatment. Your doctor can usually help to reduce these symptoms by lowering the dose of ATTORA LA or changing the times when the capsules are taken.
- **Tell your doctor if you are going to have an operation.**
There is a chance of a sudden rise in blood pressure during the operation if an anaesthetic is used. Your doctor will advise if you/your child should take ATTORA LA on the day of the operation.
- If you experience abnormally sustained or frequent and painful erections of the penis on ATTORA LA treatment or after treatment discontinuation, you may need urgent medical treatment. This can occur in any age group.
- **If this occurs, tell your doctor immediately.**
If taking ATTORA LA with medicines that raise the level of serotonin in the body (serotonergic medicines, e.g. sertraline and venlafaxine used to treat depression) and you experience a combination of the following symptoms: restlessness, tremor, sudden muscle contractions, abnormal high temperature, nausea and vomiting, stop treatment with ATTORA LA and these medicines and tell your doctor immediately.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how ATTORA LA affects you.

This medicine may cause hallucinations, dizziness, drowsiness, blurred vision or other central nervous system side effects which can affect concentration in some people. If you have any of these symptoms, do not drive, use machines or do anything else that needs quick reactions or could be dangerous.

Drinking alcohol

Do not drink alcohol whilst you are taking ATTORA LA. Remember that some foods and medicines contain alcohol.

Drinking alcohol is not recommended because it can worsen some of the unwanted effects of this medicine, such as dizziness and drowsiness

Looking after your medicine

Keep your medicine in the original container until it is time to take a dose.

Store it in a cool dry place away from moisture, heat or sunlight; for example, do not store it:

- in the bathroom or near a sink, or
- in the car or on window sills.

Keep it where young children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Getting rid of any unwanted medicine

If your doctor tells you to stop taking this medicine or the expiry date has passed, take it to any pharmacy for safe disposal.

Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Less serious side effects

Less serious side effects	What to do
Mental health-related <ul style="list-style-type: none">• excessive emotional distress or excitement• nervousness• feeling anxious, agitated• feeling depressed• feeling aggressive• unusually active, depressed mood• troubled sleep or restlessness,• difficulty falling asleep, sleep disturbance, sleepiness• excessive teeth grinding	Speak to your doctor if you have any of these less serious side effects and they worry you.
Gastrointestinal-related <ul style="list-style-type: none">• nausea (feeling sick)• dry mouth• vomiting, stomach pain, upset	

<p>stomach, indigestion, toothache</p> <ul style="list-style-type: none"> • loss of appetite, decreased weight • loss of weight and slower growth in children <p>Eye-related</p> <ul style="list-style-type: none"> • blurred vision or problems focusing your eyes <p>Nervous system-related</p> <ul style="list-style-type: none"> • feeling jittery • uncontrolled speech and body movements (Tourette's syndrome) • involuntary shaking of the body (sign of tremor) • headache • dizziness • stuttering <p>Skin-related</p> <ul style="list-style-type: none"> • excessive sweating • skin rash, itchy rash and hives • fever • hair loss <p>Infection-related</p> <ul style="list-style-type: none"> • sore throat and runny nose • cough <p>Heart-related</p> <ul style="list-style-type: none"> • abnormal heart rhythm • palpitations <p>Muscle-related</p> <ul style="list-style-type: none"> • joint pain • spasm of the jaw muscles that makes it difficult to open the mouth • muscle cramps <p>Urinary-related</p> <ul style="list-style-type: none"> • bedwetting in children during the night <p>Other possible side effects with unknown frequency:</p> <p>Immune system-related</p> <ul style="list-style-type: none"> • swelling of the ears (a symptom of allergic reaction) <p>Mental health-related</p> <ul style="list-style-type: none"> • stuttering, feeling irritated, mood changes, abnormal behaviour or thinking anger, excessive awareness of surroundings, feeling over-active and uninhibited (mania), feeling disorientated, changes in sex drive, lack of feeling or emotion, doing things over and over again, being obsessed with one thing, confusion, addiction 	
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

<p>Nervous system-related</p> <ul style="list-style-type: none"> • temporary muscle weakness, loss of skin sensation or other functions of the body due to a temporary lack of blood supply to the brain (reversible ischemic neurological deficit), migraine <p>Eye-related</p> <ul style="list-style-type: none"> • double vision, dilated pupils, trouble seeing • increased eye pressure (increased intraocular pressure) <p>Heart-related</p> <ul style="list-style-type: none"> • stopped heartbeat, heart attack • tiredness <p>Respiratory-related</p> <ul style="list-style-type: none"> • sore throat, shortness of breath • nose bleed <p>Gastrointestinal-related</p> <ul style="list-style-type: none"> • diarrhoea, constipation <p>Skin-related</p> <ul style="list-style-type: none"> • swelling of face and throat, redness of the skin, large red blotches on the skin appearing within a few hours of taking the medicine. <p>Muscle-related</p> <ul style="list-style-type: none"> • muscle pain, muscle twitching <p>Reproductive system-related</p> <ul style="list-style-type: none"> • inability to develop or maintain an erection <p>Breast-related</p> <ul style="list-style-type: none"> • swelling of the breasts in men <p>Additional side effects that occurred with other medicines containing same drug substance of ATTORA LA (frequency unknown):</p> <p>Urinary-related</p> <ul style="list-style-type: none"> • blood in the urine <p>Heart-related</p> <ul style="list-style-type: none"> • abnormal sounds from heart • chest pain, sudden death of heart origin 	
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

Serious side effects

Serious side effects	What to do
<p>Severe allergic reaction</p> <ul style="list-style-type: none"> • swelling of the face, lips or tongue (severe allergic reaction) <p>Nervous system-related</p> <ul style="list-style-type: none"> • sudden high fever, severe convulsions 	<p>Call your doctor straight away or go straight to the Emergency Department at your nearest</p>

<ul style="list-style-type: none"> • severe headache or confusion, weakness or paralysis of limbs or face, difficulty speaking • uncontrollable writing movements of the limb, face and/or trunk • uncontrollable twitching, jerking • fits (seizures, convulsions, epilepsy) • muscle twitching or tics <p>Heart related</p> <ul style="list-style-type: none"> • fast heartbeat, chest pain <p>Infection-related</p> <ul style="list-style-type: none"> • a sore throat and fever or chills <p>Mental health-related</p> <ul style="list-style-type: none"> • seeing or feeling things that are not really there (hallucination) • thoughts or attempts of killing yourself (suicidal ideation or attempt (including completed suicide)) <p>Skin-related</p> <ul style="list-style-type: none"> • skin blisters or itching • red blotches on the skin • bruising <p>Reproductive system-related</p> <ul style="list-style-type: none"> • prolonged erection, causing discomfort of the penis (sign of priapism) <p>Blood related</p> <ul style="list-style-type: none"> • abnormal decrease in the levels of all type of blood cells. This is associated with anaemia, decreased platelets, and decreased white blood cells. This can lead to fatigue, increased risk of bleeding, and/or increased risk of infection. This may be serious or life threatening (pancytopenia). <p>Blood vessel-related</p> <ul style="list-style-type: none"> • fingers and toes feeling numb, feeling cold, tingling and changing colour (from white to blue, then red) when cold (Raynaud’s phenomenon, peripheral coldness) 	<p>hospital if you notice any of these serious side effects.</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at

www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription and your doctor has special permission to prescribe it.

What ATTORA LA contains

Active ingredient (main ingredient)	ATTORA LA capsules contain 10 mg, 20 mg, 30 mg, 40 mg or 60 mg of methylphenidate hydrochloride as the active ingredient.
Other ingredients (inactive ingredients)	<ul style="list-style-type: none"> • microcrystalline cellulose • hypromellose • methacrylic acid copolymer • purified talc • ethylcellulose • hypromellose • triethyl citrate • gelatin • titanium dioxide • iron oxide yellow (only for 10 mg, 30 mg, 40 mg and 60 mg capsules) • Capsugel Ink 10A2 Black (ARTG PI No. 109521)
Potential allergens	N/A

ATTORA LA does not contain gluten.

Do not take this medicine if you are allergic to any of these ingredients.

What ATTORA LA looks like

ATTORA LA 10 mg capsules are opaque hard gelatin capsule with rich yellow cap and white body imprinted “10” with black ink filled with white to off-white spherical pellets; bottles of 30 capsules.

ATTORA LA 20 mg capsules are opaque hard gelatin capsule with white cap and white body imprinted “20” with black ink filled with white to off-white spherical pellets; bottles of 30 capsules.

ATTORA LA 30 mg capsules are opaque hard gelatin capsule with ivory cap and ivory body imprinted “30” with black ink filled with white to off-white spherical pellets; bottles of 30 capsules.

ATTORA LA 40 mg capsules are opaque hard gelatin capsule with rich yellow cap and rich yellow body imprinted “40” with black ink filled with white to off-white spherical pellets; bottles of 30 capsules.

ATTORA LA 60 mg capsules are opaque hard gelatin capsule with ivory cap and white body imprinted "60" with black ink filled with white to off-white spherical pellets; bottles of 30 capsules.

Who sponsors ATTORA LA

Arrotex Pharmaceuticals Pty Ltd
15 – 17 Chapel Street
Cremorne VIC 3121
Australia
www.arrotex.com.au

This leaflet was prepared in January 2026.

Australian Registration Number

ATTORA LA 10 mg: AUST R 408176

ATTORA LA 20 mg: AUST R 408177

ATTORA LA 30 mg: AUST R 408175

ATTORA LA 40 mg: AUST R 408178

ATTORA LA 60 mg: AUST R 408174