

Xylocaine 2% Jelly

lidocaine hydrochloride

Consumer Medicine Information

What is in this leaflet

This leaflet answers some of the common questions people ask about Xylocaine 2% Jelly.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor will have weighed the risks of you using Xylocaine 2% Jelly against the benefits they expect it will have for you.

If you have any concerns about using this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What Xylocaine 2% Jelly is for

Xylocaine 2% Jelly is used to prevent pain and discomfort during medical tests and procedures. Your doctor will explain fully the tests which will be carried out and the reason for them.

Xylocaine 2% Jelly belongs to a group of medicines called local anaesthetics. It works by making the nerves unable to pass messages to the brain. The jelly also acts as a lubricant.

Your doctor will have explained why you are being treated with Xylocaine 2% Jelly and told you what dose you will be given.

Follow all directions given to you by your doctor carefully.

They may differ from the information contained in this leaflet.

Your doctor may prescribe this medicine for another use. Ask your doctor if you want more information.

Xylocaine 2% Jelly is not addictive.

Before you use it

When you must not use it

Do not use Xylocaine 2% Jelly if you are pregnant or breastfeeding unless your doctor says to do so. Ask your doctor about the risks and benefits involved.

There is no evidence available at the moment to show that being given Xylocaine 2% Jelly under your doctor's instructions is harmful to unborn babies.

Your baby can take in very small amounts of Xylocaine 2% Jelly from breast milk if you are breastfeeding, but it is unlikely that the amount available to the baby will do any harm.

Do not use after the use by (expiry) date printed on the pack. It may have no effect at all, or worse, an entirely unexpected effect if you take it after the expiry date.

Do not use Xylocaine 2% Jelly if the packaging is torn or shows signs of tampering.

Do not use it to treat any other complaints unless your doctor tells you to.

Do not give this medicine to anyone else.

Before you start to use it

You must tell your doctor if:

- 1. you have any allergies to**

- ingredients listed at the end of this leaflet
- other local anaesthetics
- any other substances.

If you have an allergic reaction, you may get a skin rash, hayfever, asthma or feel faint.

2. you have any of these medical conditions

- epilepsy
- heart problems
- liver problems
- kidney problems
- open wounds or infection where the jelly will be used.

It may not be safe for you to use Xylocaine 2% Jelly if you have any of these conditions.

Taking other medicines

Tell your doctor if you are taking any other medicines, including medicines that you buy at the chemist, supermarket or health food shop.

These medicines may affect the way Xylocaine 2% Jelly works.

Your doctor or pharmacist can tell you what to do if you are taking any of these medicines.

If you have not told your doctor about any of these things, tell them before you use any Xylocaine 2% Jelly.

Using Xylocaine 2% Jelly

How to use it

Xylocaine 2% Jelly will be given to you by your doctor or nurse. Your doctor or pharmacist may also

suggest that you buy it without prescription.

Xylocaine 2% Jelly is given before a procedure to prevent discomfort, or applied to the end of medical instruments for lubrication as well.

The maximum dose of jelly is 30 mL (600 mg) in a 12 hour period.

Overdose

Telephone your doctor or go to casualty at your nearest hospital immediately if you think that you or anyone else may have used too much Xylocaine 2% Jelly even if there are no signs of discomfort or poisoning.

Xylocaine 2% Jelly can be absorbed into the blood, but very large amounts of it are needed to give an overdose.

Signs of overdose include dizziness, blurred vision, tremor or nervousness.

While you are using it

Things to be careful of

Be careful driving or operating machinery after you have used or have been given Xylocaine 2% Jelly.

You may be drowsy and your reflexes may be slow.

Do not eat or drink anything for at least 1 hour after using Xylocaine 2% Jelly in the mouth or throat area.

You may swallow your food the wrong way or burn or bite your mouth.

Please talk to your doctor or pharmacist about these possibilities if you think they may bother you.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel

well while you are using Xylocaine 2% Jelly.

Xylocaine 2% Jelly will help to relieve pain and discomfort in most people, but it may have unwanted side-effects in a few people.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor if you notice any of the following and they worry you:

- sore throat following use in the mouth or throat
- skin rash or irritation
- drowsiness.

These are all mild side effects of Xylocaine 2% Jelly.

Tell your doctor immediately or go to casualty at your nearest hospital if you notice any of the following:

- wheezing or difficulty breathing
- chest pain
- severe rash or itching
- increased sweating
- dizziness
- confusion
- numbness
- blurred vision
- vomiting
- drowsiness
- sensations of heat or cold.

These are all serious side effects. You may need urgent medical attention.

Serious side effects are rare.

Xylocaine 2% Jelly can pass into the bloodstream, although this is extremely unlikely to happen in high enough amounts from Xylocaine 2% Jelly to cause any problems. In high doses serious side effects occur. These may include:

- fits
- unconsciousness

- breathing problems
- low blood pressure
- slow heart beat
- collapse.

These are very serious side effects. You may need urgent medical attention or hospitalisation.

These serious side effects are very rare.

Tell your doctor if you notice anything else that is making you feel unwell.

Some people may get other side effects while using Xylocaine 2% Jelly.

Storage

Keep your Xylocaine 2% Jelly in the pack until it is time to use it.

If you take it out of the pack it will not keep well.

Keep it in a cool dry place where the temperature stays below 25°C.

Do not store it or any other medicine in the bathroom or near a sink.

Heat and dampness can destroy some medicines.

Keep it where young children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Do not leave it in the car on hot days.

Disposal

Ask your pharmacist what to do with any medicine you have left over if your doctor tells you to stop using it, or you find that the expiry date has passed.

Product description

Xylocaine 2% Jelly is a clear to almost clear, slightly coloured, highly viscous liquid, packed in a 30 g aluminium tube. The tube is enclosed in a clear blister and

includes one single use disposable nozzle.

It is supplied for single use only.

Xylocaine 2% Jelly is sterile, until opened.

Each gram of Xylocaine 2% Jelly contains lidocaine hydrochloride 21.3 mg (equivalent to lidocaine hydrochloride anhydrous 20 mg) as the active ingredient.

Excipients include:

- hypromellose
- methyl hydroxybenzoate
- propyl hydroxybenzoate
- sodium hydroxide
- purified water
- hydrochloric acid (for pH adjustment).

Sponsor

Aspen Pharmacare Australia Pty
Ltd
34-36 Chandos St
St Leonards NSW 2065
Australia

Australian Registration Number:
AUST R 12026

This leaflet was revised in July
2024.