VARENICLINE VIATRIS

Active ingredient(s): [varenicline (as tartrate)]

Consumer Medicine Information

This leaflet provides important information about using VARENICLINE VIATRIS. You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using VARENICLINE VIATRIS.

Where to find information in this leaflet:

- 1. Why am I using VARENICLINE VIATRIS?
- 2. What should I know before I use VARENICLINE VIATRIS?
- 3. What if I am using other medicines?
- 4. How do I use VARENICLINE VIATRIS?
- 5. What should I know while using VARENICLINE VIATRIS?
- 6. Are there any side effects?
- 7. Product details

concentrating, restlessness, decreased heart rate, increased appetite or weight gain. Not everybody is affected by symptoms of nicotine withdrawal.

Although it is not recommended that you smoke after your quit date, VARENICLINE VIATRIS will reduce your enjoyment of cigarettes if you do smoke while on treatment.

1. Why am I using VARENICLINE VIATRIS?

VARENICLINE VIATRIS contains the active ingredient varenicline (as tartrate).

VARENICLINE VIATRIS is used to help adults stop smoking.

It can help to reduce craving and withdrawal symptoms that may happen while you stop smoking.

People giving up smoking are often affected by nicotine withdrawal. Symptoms of nicotine withdrawal can include craving for tobacco, irritability, frustration, feeling angry, sleep problems, depressed mood, feeling anxious, difficulty