

---

## Consumer Medicine Information

---

### What is in this leaflet

This leaflet answers some common questions about SOTALOL-WGR. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you using SOTALOL-WGR against the benefits they expect it will have for you.

**If you have any concerns about using this medicine, ask your doctor or pharmacist.**

**Keep this leaflet with the medicine.**

You may need to read it again.

---

### What SOTALOL-WGR is used for

SOTALOL-WGR contains sotalol hydrochloride as the active ingredient which belongs to the family of drugs known as beta-blockers.

SOTALOL-WGR is used to treat “arrhythmias”, which is a problem when the heart beats too quickly or with the wrong rhythm.

SOTALOL-WGR slows down and steadies the heart beat, reducing the effort the heart has to put into pumping blood.

**Ask your doctor if you have any questions about why SOTALOL-WGR has been prescribed for you.**

Your doctor may have prescribed it for another reason.

There is no evidence that this medicine is addictive.

SOTALOL-WGR is only available with a doctor's prescription.

---

### Before using it

---

#### *When you must not use it*

**Do not use SOTALOL-WGR if you have an allergy to:**

- sotalol hydrochloride
- other beta blockers
- any of the ingredients listed at the end of this leaflet

Some of the symptoms of an allergic reaction may include:

- shortness of breath, wheezing or difficulty breathing
- skin rash, itching or hives
- swelling of the face, lips, tongue or any part of the body.

**Do not take SOTALOL-WGR if you have asthma.**

**You should not take this medicine if you have allergies, or problems with your kidneys or thyroid gland, unless you have discussed it with your doctor.**

**You should not take this medicine if you are pregnant or are breast feeding, or if you intend to breast feed.**

**If you have any problems with your heart or circulation or amount of magnesium in your blood, discuss them with your doctor.**

**You should not take SOTALOL-WGR with any other medicines your doctor does not know about, particularly if they are to control high blood pressure, heart**

**conditions, depression, hayfever, allergies, infections or diabetes.**

**Do not take it if the expiry (EXP) date printed on the pack has passed**

**Do not take it if the packaging is torn or shows signs of tampering.**

**Talk to you doctor if you are unsure whether you should start taking this medicine.**

#### *Before you start to take it*

- **Make sure your doctor knows what other medicines you are already taking including ones you have bought yourself at the chemist or supermarket.**
- **In particular remind your doctor if you have asthma, bronchitis or any allergies such as hay fever, food allergies or are allergic to bee or wasp stings.**
- **Make sure your doctor is aware of any kind of heart disease, diabetes, pheochromocytoma, kidney disease or thyroid disease that you have or have had, or if you have been told that your pulse is slow or irregular.**
- **Tell your doctor if you have ever had trouble with the levels of salts like potassium or magnesium in your blood.**
- **Remind your doctor if you are going to have surgery involving a general anaesthetic even if it is only minor.**
- **Tell your doctor if you are pregnant, if you are planning to become pregnant, or if you are breast feeding.**
- **Tell your doctor if you have been given SOTALOL-WGR (or any other beta-blocker) before and if you had any problems.**
- **Remind your doctor if you have hardening of the arteries (cold fingers and toes or pain in the back of your legs when you walk).**

**Be careful driving or operating machinery until you know how SOTALOL-WGR affects you.**

As with other medicines it may cause dizziness, light-headedness or drowsiness in some people. If this occurs do not drive or operate machinery or undertake any other activity that could be dangerous if you are dizzy, light-headed or drowsy.

#### *Taking other medicines*

Some medicines can affect the way SOTALOL-WGR works.

**You should always tell your doctor about any other medicines you take, even those bought without a doctor's prescription. It is especially important that you tell your doctor if you are taking the following:**

- medicines which lower blood pressure (including other betablockers)
- floctafenine (medicine used for the short term treatment of mild to moderate pain)
- for the treatment of certain infections (e.g. erythromycin IV, amphotericin B, pentamidine, halofantrine)
- steroids
- laxatives

- clonidine (sometimes used to treat hot flushes or headaches)
- any other medicines used to treat irregular heart rhythm or beat
- digoxin, a medicine used for heart failure
- medicines used to treat angina or other heart conditions
- antidepressants (medicines used to treat depression)
- insulin or other drugs used to control diabetes
- Medicines used to control or prevent asthma (inhalers or tablets) or to control allergies or which are used for other lung problems
- Antihistamine medicine including terfenadine and astemizole that may be used to treat hayfever, allergies or to relieve symptoms of cold and flu
- quinolone antibiotics (medicines used to treat infections)
- diuretics (water tablets)
- neuromuscular blocking agents like tubocurarine
- some medicine used during surgery or emergency situations, such as anaesthetics.

**If you are not sure whether you are taking any of these medicines, check with your doctor or pharmacist.**

---

### How to take it

**Follow all directions given to you by your doctor or pharmacist carefully.**

They may differ from the information contained in this leaflet.

**If you do not understand the instructions on the box, ask your doctor or pharmacist for help.**

#### *How much to take*

Your doctor will decide on the dose of SOTALOL-WGR tablets for you, and may need to change the dose a few times to get the best level for you.

The usual dose is 80mg to 160mg twice a day. Your doctor may need to increase this as a very few patients may need up to three to four 160mg tablets spread over a day. The dosage may need to be different if you have a kidney problem.

#### *How to take it*

You should take your tablets with water one to two hours before meals. Do not take SOTALOL-WGR with milk or meals.

#### *How long to take it for*

If you have been prescribed SOTALOL-WGR you must be sure to follow your doctors instructions carefully.

**Do not stop taking SOTALOL-WGR tablets suddenly.**

The dose needs to be reduced gradually over 7 to 14 days.

#### *If you forget to take it*

**It is important not to miss a dose but if you do, take your next dose at the normal time and with the normal amount.**

*If you take too much (overdose)*

**Immediately telephone your doctor or the Poisons Information Centre (telephone 13 11 26) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much SOTALOL-WGR. Do this even if there are no signs of discomfort or poisoning.**

You may need medical attention.

Too much of this medicine will cause your blood pressure and heart rate to drop to dangerous levels. Serious heart problems may develop and this could be fatal.

---

## Side Effects

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking SOTALOL-WGR.**

SOTALOL-WGR helps most people with arrhythmia, but it may have unwanted side effects in some people. Rarely, serious heart problems can develop while you are taking normal doses but you must remember that you are taking this medicine because your heart already has a serious problem. It is very important that your doctor keeps a check on your progress.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

Ask your doctor or pharmacist to answer any questions you may have.

**Tell your doctor if you notice any of the following and they worry you:**

- dizziness, lightheadedness or fainting, especially when you get up from a sitting or lying position. Getting up slowly may help.
- tiredness, lack of energy, weakness
- headache, fever
- cramps
- irritated eyes, blurred vision, worsening of eyesight, increased sensitivity of the eyes to sunlight, eye infections
- feeling sick, vomiting, stomach upset, diarrhoea, wind
- change in taste sensation
- anxiety, depression, mood changes
- confusion hallucinations
- problems with sexual function
- sleep problems, unusual dreams
- worsening of psoriasis, thickening of skin, itching, alopecia (hair loss)
- hearing disturbances
- tingling or numbness in the hands or feet, cold limbs
- muscle spasms
- excessive sweating
- dry mouth

**Tell your doctor immediately or go to casualty at the nearest hospital if you notice any of the following:**

- chest tightness, wheezing, shortness of breath
- very slow heart beat
- fast, irregular heart beat, palpitations
- chest pain
- any type of p rash, itching
- shortness of breath (sometimes with tiredness, weakness and reduced ability

to exercise), which may occur together with swelling of the feet or legs due to fluid build up

- fainting

This is not a complete list of all possible side effects. Others may occur in some people and there may be some side effects not yet known.

**Tell your doctor if you notice anything that is making you feel unwell while you are taking, or soon after you have finished taking SOTOCOR, even if it is not on this list.**

---

## While you are taking it

### *Things you must do*

**Tell your doctor if you become pregnant while taking SOTALOL-WGR.**

You doctor can discuss with the risks of taking it while you are pregnant.

**If you are about to be started on any new medicine, remind your doctor or pharmacist that you are taking SOTALOL-WGR.**

**Tell any other doctor, dentist, or pharmacist who treats you that you are taking this medicine.**

### *Things you must not do*

**Do not give your medicine to anyone else, even if they have the same condition as you.**

**Do not take SOTALOL-WGR to treat any other complaints unless your doctor tells you to.**

**Do not let yourself run out of tablets over weekends or on holidays.**

---

## After taking it

### *Storage*

**Store your tablets safely in a cool dry place. Protect from light and moisture.**

**Do not store SOTALOL-WGR or any other medicine in the bathroom or near a sink.**

**Do not leave it in the car on hot days or on window sills.**

Heat and dampness can destroy some medicines.

**Keep your tablets in the original packaging they are provided in until it is time to take them.**

If you take the tablets out of the blister they will not keep well.

**Keep all medicines out of the reach of children.**

A locked cupboard at least one-and-a-half meters above the ground is a good place to store medicines.

---

## Product description

### *What it looks like*

SOTALOL-WGR 80mg tablets are light blue, biconvex, capsule shaped tablets, plain on one side and engraved 'S', '80' and a break bar on the other side.

SOTALOL-WGR 160mg are light blue biconvex, capsule shaped tablets, plain on one side and engraved 'S', '160' and a break bar on the other side.

### *Active ingredients*

**SOTALOL-WGR 80mg tablets contain 80mg sotalol hydrochloride per tablets**

**SOTALOL-WGR 160mg tablets contain 160mg sotalol hydrochloride per tablet**

The tablets also contain:

- microcrystalline cellulose
- magnesium stearate
- maize starch
- lactose (anhydrous)
- stearic acid
- colloidal anhydrous silica
- indigo carmine (CI 73015)

This medicine contains sugars as lactose.

---

## Supplier

Wagner Pharmaceuticals Pty Ltd  
6 Albert Street  
Preston VIC 3072  
Tel: 1800 936 140

Australia Registration Numbers:

**SOTALOL-WGR** Tablets 80mg AUST R 333533 in blister pack of 60 tablets

**SOTALOL-WGR** Tablets 160mg AUST R 333534 in blister pack of 60 tablets

This leaflet was revised in August 2024.