

# NEXOLE Tablets

*Esomeprazole magnesium*

---

## Consumer Medicine Information

---

### What is in this leaflet

---

This leaflet answers some common questions about NEXOLE.

It does not contain all the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking NEXOLE against the benefits it is expected to have for you.

**If you have any concerns about taking this medicine, ask your doctor or pharmacist.**

**Keep this leaflet with the medicine.**

You may need to read it again.

---

### What NEXOLE is used for

---

#### *Reflux oesophagitis*

NEXOLE is taken to treat reflux oesophagitis. This can be caused by "washing back" (reflux) of food and acid from the stomach into the food pipe (oesophagus).

Reflux can cause a burning sensation in the chest rising up to the throat, also known as heartburn.

NEXOLE is also taken to help stop reflux oesophagitis coming back or relapsing.

#### *Upper gastrointestinal symptoms associated with Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) therapy*

NEXOLE is taken to treat the symptoms of pain or discomfort in the stomach caused by NSAIDs, a type of medicine for pain or inflammation.

NEXOLE is also taken to help heal and prevent ulcers caused by NSAIDs.

#### *Peptic ulcers associated with Helicobacter pylori Infection*

Most people who have a peptic (gastric and duodenal) ulcer also have a bacterium called *Helicobacter pylori* in their stomach. Depending on the position of the ulcer it is called a gastric or duodenal ulcer. A gastric ulcer occurs in the stomach. A duodenal ulcer occurs in the duodenum which is the tube leading out from the stomach.

If you have a peptic ulcer, your doctor will prescribe NEXOLE with antibiotics. When NEXOLE and antibiotics are taken together, they work to kill the bacterium and let your ulcer heal. You may need further treatment with antibiotics.

#### *Zollinger-Ellison syndrome*

NEXOLE is also used to treat a rare condition called Zollinger-Ellison syndrome, where the stomach produces large amounts of acid, much more than in ulcers or reflux disease.

#### *Prevention of rebleeding of gastric or duodenal ulcers*

NEXOLE is also used as prolonged treatment after prevention of rebleeding of ulcers of stomach or gut with intravenous esomeprazole.

When peptic ulcers become severe enough, they start to bleed. You may receive treatment injected into your veins initially. This treatment may be followed with NEXOLE for a longer period of time. This is to help your ulcer/s to heal.

#### *How NEXOLE works*

NEXOLE is a type of medicine called a proton-pump inhibitor. It works by decreasing the amount of acid made by the stomach to give relief of symptoms and allow healing to take place. This does not stop food being digested in the normal way.

**Follow all directions given to you by your doctor carefully.**

They may differ from the information contained in this leaflet.

**Ask your doctor if you want more information.**

Your doctor may have prescribed NEXOLE for another reason.

There is no evidence that NEXOLE is addictive.

---

### Before you take NEXOLE

---

#### *When you must not take NEXOLE*

**Do not take NEXOLE if you have allergies to:**

- Esomeprazole or any ingredient listed at the end of this leaflet
- Any medicines containing a proton-pump inhibitor.

Some of the symptoms of an allergic reaction may include shortness of breath, wheezing or difficulty breathing; swelling of the face lips, tongue or other parts of the body; rash, itching or hives on the skin.

**Do not take NEXOLE if you are taking atazanavir or cilostazol.**

Please check with your doctor or pharmacist if you are taking these medicines. These medicines will be affected by NEXOLE.

**NEXOLE is not approved for use in children younger than 1 year of age.**

There is no specific information about use in children younger than 1 year of age, so NEXOLE is not recommended in these patients.

**Do not take NEXOLE after the use by (expiry) date printed on the pack or if the packaging is torn or shows signs of tampering.**

If it has expired or is damaged, return it to your pharmacist for disposal.

*Before you start to take it*

**You must tell your doctor if you have:**

- allergies to any other medicines, foods, dyes or preservatives
- any problems with your liver
- experienced any significant weight loss, difficulty in swallowing or dark tarry stools
- severe kidney problems
- any other medical conditions
- been diagnosed with osteoporosis
- if you have ever had a skin reaction after treatment with a medicine similar to NEXOLE that reduces stomach acid

**Do not take NEXOLE if you are pregnant or breast-feeding unless your doctor says so. Ask your doctor about the risks and benefits involved.**

It is not known if it is safe for you to take NEXOLE while you are pregnant. It may affect your baby.

It is not known if your baby can take in NEXOLE from breast milk if you are breastfeeding.

*Taking other medicines*

**Do not take NEXOLE if you are taking the following medicines:**

- atazanavir and nelfinavir (used to treat HIV)
- cilostazol (used to reduce leg pain when walking caused by blood circulation disease)

**Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from your pharmacy, supermarket or health food shop.**

Some medicines may interfere with NEXOLE. These include:

- medicines used to treat fungal infections such as ketoconazole, itraconazole and voriconazole
- medicines used to treat bacterial infections such as clarithromycin and rifampicin
- cisapride
- diazepam, a medicine used to treat anxiety and some other conditions
- phenytoin, a medicine used to treat epilepsy
- medicines used to treat depression such as citalopram, clomipramine or imipramine
- warfarin and clopidogrel - medicines used to prevent blood clots
- digoxin, a medicine used to treat certain heart problems
- medicines for HIV treatment such as nelfinavir
- St. John's Wort, a herbal remedy used to treat mood disorders
- methotrexate, a medicine used to treat arthritis and some types of cancer
- tacrolimus – a medicine used to assist in organ transplants

- erlotinib or related medicines used to treat cancer

These medicines may be affected by NEXOLE or may affect how well it works. You may need different amounts of your medicine or you may need to take different medicines.

Your doctor can tell you what to do if you are taking any other medicines.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking NEXOLE.

**If you have not told your doctor about any of these things, tell them before you take NEXOLE.**

---

## How to take NEXOLE

---

**Follow all directions given to you by your doctor and pharmacist carefully.**

These instructions may differ from the information contained in this leaflet.

**If you do not understand the directions on the label, ask your doctor or pharmacist for help.**

*How to take it*

**Take one NEXOLE tablet each day, unless your doctor has told you otherwise.**

The dose of NEXOLE tablets is usually 20 mg or 40 mg a day depending on what condition you are being treated for and how severe it is.

**Swallow NEXOLE tablets whole with a glass of water. Do not crush or chew the tablets.**

If the tablets are chewed or crushed they will not work properly.

**If you have difficulty swallowing the tablets:**

1. Place the tablet in half a glass of non-carbonated water. Mineral water or other liquids are not suitable.

- Gently mix the tablet and water by stirring, taking care not to crush the tablet.
- Stir until the tablet dissolves into little pellets.
- Drink the liquid with the pellets immediately, or within 30 minutes. Do not chew the pellets.
- Rinse the glass with half a glass of water and drink.

**If you cannot swallow at all, follow steps 1-3 above to disperse the tablets and administer the liquid and pellets through a gastric tube.**

**Take NEXOLE at about the same time each day.**

Keeping a regular time for taking NEXOLE will help to remind you to take it.

**Keep taking NEXOLE for as long as your doctor recommends.**

In most patients, NEXOLE relieves symptoms rapidly and healing is usually complete within 4 weeks. Continue taking NEXOLE for as long as your doctor tells you to.

NEXOLE can be taken with food or on an empty stomach.

*If you forget to take it*

**If you forget to take a dose, take it as soon as you remember, and then go back to taking it as you would normally.**

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

**Do not take a double dose to make up for the dose that you missed.**

If you have trouble remembering when to take your medicine, ask your pharmacist for some hints.

*Taking too much (overdose)*

**Telephone your doctor or the Poisons Information Centre (13 11 26) or go to Accident and Emergency at your nearest hospital immediately if you think that you or anyone else may have**

**taken too many NEXOLE tablets. Do this even if there are no signs of discomfort or poisoning.**

---

## While you are taking NEXOLE

---

*Things you must do*

**Take NEXOLE exactly as your doctor has prescribed.**

**If you are about to be started on any new medicine, tell your doctor, dentist or pharmacist that you are taking NEXOLE.**

**Tell all doctors, dentists and pharmacists who are treating you that you are taking NEXOLE.**

**Tell your doctor if you become pregnant while you are taking NEXOLE.**

**Tell your doctor if your symptoms return.**

Although NEXOLE can heal ulcers successfully, it may not prevent them recurring at a later date.

**If you need to have any medical tests while you are taking NEXOLE, tell your doctor.**

It may affect the results of some tests.

*Things you must not do*

**Do not take it to treat any other complaints unless your doctor tells you to.**

**Do not give this medicine to anyone else, even if their symptoms seem similar to yours.**

---

## Side Effects

---

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking NEXOLE.**

NEXOLE helps most people with peptic ulcers or reflux disease, but it may have unwanted side effects in a few people.

All medicines can have side effects. Sometimes they are serious, most of

the time they are not. You may need medical treatment if you get some of the side effects.

**Ask your doctor or pharmacist to answer any questions you may have.**

**Tell your doctor if you notice any of the following and they worry you:**

- headache
- stomach pain
- constipation
- diarrhoea
- wind
- nausea
- vomiting
- itching
- skin rash
- dizziness
- dry mouth
- swelling of hands, ankles or feet

These side effects are usually mild.

**Tell your doctor immediately if you notice any of the following:**

- disturbances in vision (blurred vision)
- feeling depressed
- mood changes, agitation, aggression, hallucination
- confusion
- muscle pain or weakness
- joint pain
- increase in breast size (males)
- increased sweating
- changes in sleep patterns
- fever
- increased bruising
- tremor
- blood in the urine
- hair loss
- liver problems (often seen as yellowing of your skin or the whites of your eyes, feeling generally unwell, nausea, vomiting, loss of appetite)
- difficulty in breathing (bronchospasm)
- tingling, pricking or numbness of the hands and feet (“pins and needles”)

- taste disturbance, sore mouth, thrush
- increased sensitivity to sunlight
- frequent infections such as fever, severe chills, sore throat or mouth ulcers

These are all rare or very rare side effects. These side effects may require medical attention.

**If you notice any of the following happen, tell your doctor immediately or go to Accident and Emergency at the nearest hospital:**

- shortness of breath, wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- severe skin reaction which may include rash, itching, redness, blistering or peeling of the skin.

These are very serious side effects. You may need urgent medical treatment or hospitalisation. These side effects are rare.

Occasionally, NEXOLE may be associated with changes in your liver or blood, which may require your doctor to do certain blood tests.

**Tell your doctor if you think you have any of these effects or notice anything else that is making you feel unwell.**

Some people may get other side effects while taking NEXOLE.

Other problems are more likely to arise from the condition itself rather than the treatment.

For this reason, contact your doctor immediately if you notice any of the following:

- pain or indigestion during treatment with NEXOLE
- you begin to vomit blood or food
- you pass black (blood-stained) motions

Tell your doctor if your reflux symptoms return after you stop taking NEXOLE.

---

## After taking NEXOLE

---

### Storage

**Keep your NEXOLE in the blister pack until it is time to take them.**

If you take NEXOLE out of the blister pack it will not keep well.

**Keep it in a cool, dry place where the temperature stays below 25°C.**

**Do not store it or any other medicine in the bathroom or near a sink.**

Do not leave it in the car or on a window sill. Heat and dampness can destroy some medicines.

**Keep it where young children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

### Disposal

**If your doctor tells you to stop taking NEXOLE or the tablets have passed their expiry date, ask your pharmacist what to do with any tablets you have left over.**

---

## Product description

---

### What NEXOLE looks like

NEXOLE 20 mg tablets are light brick red to brown, oval, biconvex, enteric-coated tablets with 'E5' debossed on one side and plain on other side.

NEXOLE 40 mg tablets are light brick red to brown, oval, biconvex, enteric-coated tablets with 'E6' debossed on one side and plain on other side.

### Ingredients

#### Active ingredient:

NEXOLE 20 mg tablets – esomeprazole magnesium equivalent to 20 mg esomeprazole

NEXOLE 40 mg tablets – esomeprazole magnesium equivalent to 40 mg esomeprazole

#### Inactive ingredients:

Non Pareil Seeds 40-60 [sugar spheres (sucrose and maize starch); Proprietary ingredient # 108075], hypromellose, croscopovidone, povidone, macrogol 400, purified talc, hypromellose phthalate, diethyl phthalate, macrogol 6000, microcrystalline cellulose, sodium stearyl-fumarate, macrogol 4000 and Opadry Complete Film Coating System 03B86651 Brown (Proprietary Ingredient # 106495).

NEXOLE tablets do not contain lactose, glucose or gluten.

### Supplier

**Arrow Pharma Pty Ltd  
15-17 Chapel Street  
Cremorne Vic 3121  
www.arrowpharma.com.au**

### Australian Registration Numbers

NEXOLE 20 mg tablets: AUST R 184769

NEXOLE 40 mg tablets: AUST R 184755

This leaflet was prepared in July 2020.