

# COMPOUND SODIUM LACTATE (HARTMANN'S) SOLUTION WITH GLUCOSE INTRAVENOUS INFUSION

## Consumer Medicine Information

### What is in this leaflet?

This leaflet answers some common questions about **Compound Sodium Lactate (Hartmann's) Solution with Glucose Intravenous (IV) Infusion**.

It does not contain all of the available information. All medicines have risks and benefits. Your doctor has weighed the risks of you using Hartmann's Solution with Glucose IV Infusion against the benefit they expect it will have for you.

It does not take the place of talking to your doctor or pharmacist.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

### What Hartmann's Solution with Glucose IV Infusion is used for

This medicine is used to replace and balance body fluid and mineral salts that may be lost for a variety of medical reasons and also as an energy source.

### Before you are given Hartmann's Solution with Glucose IV Infusion

*The Hartmann's Solution with Glucose IV Infusion should not be given to you if:*

- you have an allergy to any ingredient listed at the end of this leaflet;
- you have had an allergic reaction to corn or corn products. Some of the symptoms of an allergic reaction may include skin rash, peeling of the skin and swelling of the face, lips or tongue, causing difficulty swallowing or shortness of breath;
- you have congestive heart failure (constant wheezing, shortness of breath);
- you have severely reduced kidney function;
- you have too much sodium and chloride in your blood that is not being corrected with medication
- you have ischaemic stroke or head trauma in the last 24 hours
- it is intended to be administered at the same time and through the same injection equipment as blood preparations (citrate anticoagulated /preserved blood) or ceftriaxone, an antibiotic (if a baby < 1 month of age is to be treated, administration at the same time as ceftriaxone must never occur)
- the expiry date printed on the pack has passed.

*You must tell your doctor if you:*

- have heart problems
- have kidney problems including kidney stones
- have diabetes
- are taking any other medicine including those you can buy without a prescription, in particular digoxin (a medicine for heart problems), medicines for high blood pressure (diuretics, ACE inhibitors or ARAs), immunosuppressant medicines (tacrolimus or cyclosporine), corticosteroids, lithium or vitamin D

- are pregnant
- are breast feeding

### How Hartmann's Solution with Glucose IV Infusion is given

*How much it is given:*

Your doctor will decide how much Hartmann's Solution with Glucose IV Infusion will be given to you, which depends on your need and condition. The medicine is given by a slow rate (drip) injection; therefore it should be given by using special equipment and attended by a health professional (doctor, trained nurse).

*How it is given:*

The Hartmann's Solution with Glucose IV Infusion will be given at a slow rate of injection (drip) by your health professional. Usually, you will need to stay in a health institution (hospital, nursing home, etc.) but in some cases at home, as it requires a special medical equipment to deliver the medicine into your circulation. This delivery should be attended by a health professional. A cannula (administration needle) is placed in a vein by your doctor or nurse. The infusion is for single use, and for one person only. Any unused portion must be discarded and not used later, either for you or anyone else.

### Overdosage

The doctor or nurse giving you the Hartmann's Solution with Glucose IV Infusion has had experience in the use of this sort of medicine, so it is unlikely that you will be given an overdose. However, in case of an overdose, the infusion will be discontinued and another treatment may be needed. You may experience some of the effects listed under "Side Effects" below.

### While you are given Hartmann's Solution with Glucose IV Infusion

Discuss with your doctor the progress you have experienced after the treatment, whether any complication has occurred, especially during the first few days of therapy. Frequent clinical evaluation and laboratory tests may be required. As Hartmann's Solution and Glucose IV Infusion is normally given in a hospital, your nurse provider will take records of the progress and unexpected reactions.

### Side effects

As with any medicines, some side effects may occur. Undesirable side effect such as nausea, vomiting, headache or dizziness may occur. Your body may also retain sodium and this may result in a build-up of fluid and swelling of the hands, ankles and feet. Additionally, as with other preparations similar to Hartmann's Solution with Glucose IV Infusion, pain, irritation, inflammation, and swelling or abnormal clotting at the site of injection is also possible.

Always tell your doctor or nurse if you have any unexpected effects during or after receiving Hartmann's Solution with Glucose IV Infusion and they worry you. In rare circumstances, more serious side effects may occur such as chest pain, fast or slow

heartbeat, or fluid on the lungs or swelling of the face, lips or mouth causing breathing difficulties. If these occur, tell your health professional on duty immediately. These may be symptoms of a serious allergic reaction. Your health professional will take appropriate action promptly, such as stopping the infusion.

### Product description

#### *What Hartmann's Solution with Glucose IV Infusion looks like*

It is a clear, colourless solution filled in plastic bags.

#### *What is in Hartmann's Solution with Glucose IV Infusion?*

The active components are Sodium Lactate, Potassium Chloride, Sodium Chloride, Calcium Chloride and Glucose, formulated and dissolved in water for injection. The bag pack sizes are included below:

AHB2073

Potassium Chloride (0.04%),  
Sodium Chloride (0.6%),  
Sodium lactate (0.322%),  
Calcium Chloride (0.027%),  
Glucose (5%), AUST R 19422,  
500mL

AHB2074

Potassium Chloride (0.04%),  
Sodium Chloride (0.6%),  
Sodium lactate (0.322%),  
Calcium Chloride (0.027%),  
Glucose (5%), AUST R 48490,  
1000mL

#### *Storage*

Hartmann's Solution with glucose IV Infusion will be stored in the pharmacy or the hospital ward. It is recommended that the product be stored below 30°C.

#### *Where can you get more information?*

You can get more information from your doctor or pharmacist.

#### *Name and address of the sponsor*

Baxter Healthcare Pty Ltd  
1 Baxter Drive  
Old Toongabbie NSW 2146

#### *Date of preparation:*

March 2014