

Letroz

Letrozole Tablets 2.5 mg

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Letroz. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

You should ensure that you speak to your pharmacist or doctor to obtain the most up to date information on the medicine.

Those updates may contain important information about the medicine and its use of which you should be aware.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Letroz against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What Letroz is used for

Letroz is used to treat breast cancer in women who are post-menopausal - that is, women who no longer have periods, either naturally due to their age or after surgery or chemotherapy.

Letroz is available in tablets containing 2.5 mg of the active ingredient, letrozole.

Letrozole belongs to a family of medicines called aromatase inhibitors. They are also called "antioestrogens" because they act

by reducing the production of oestrogen in your body.

Oestrogen stimulates the growth of certain types of breast cancer. These cancers are called "oestrogen-dependent." Reducing the production of oestrogen may help to keep the cancer from growing.

This may be the first time you are taking an "antioestrogen" such as Letroz or you may have taken another "antioestrogen" such as tamoxifen in the past.

Ask your doctor if you have any questions about why Letroz has been prescribed for you.

Your doctor may have prescribed Letroz for another reason.

Letroz is only available with a doctor's prescription. It is not addictive.

Before you take Letroz

When you must not use it

Do not take Letroz if you have an allergy to:

- letrozole, the active ingredient in Letroz
- any other ingredients of Letroz listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include rash, itching or hives on the skin; swelling of the face, lips, tongue or other parts of the body; shortness of breath, wheezing or troubled breathing.

Do not take Letroz if you are still having periods.

This medicine is only used in women who are no longer having periods.

Women of child-bearing age who recently became postmenopausal or perimenopausal should use a proven method of birth control to avoid pregnancy, until your postmenopausal status is fully established. Do not take Letroz if you are pregnant or breast feeding.

It may affect your baby if you take it while you are pregnant or breast feeding.

Do not take Letroz after the use by (expiry) date printed on the pack.

If you take this medicine after the expiry date has passed, it may not work or it may make you unwell.

Do not take Letroz if the packaging is torn or shows signs of tampering.

In that case, return it to your pharmacist.

Before you start to take it

Tell your doctor if you have severe kidney or liver disease.

Your doctor may want to take special precautions while you are taking this medicine.

Tell your doctor if you are allergic to any other medicines, foods, dyes or preservatives.

Your doctor will want to know if you are prone to allergies.

If you have not told your doctor about any of these things, tell them before you take Letroz.

Taking other medicines

Tell your doctor if you are taking any other medicines, including medicines that you buy without a prescription from a pharmacy, supermarket or health food shop.

Other medicines may be affected by Letroz or they may affect how well it works.

Your doctor or pharmacist can tell you what to do when taking Letroz with other medicines.

How to take Letroz

Follow the directions given to you by your doctor and pharmacist carefully.

These directions may differ from the information contained in this leaflet.

If you do not understand the instructions on the box, ask your doctor or pharmacist for help.

How much to take

The usual dose is one Letroz tablet daily.

How to take it

Swallow the tablet with a glass of water or other liquid.

If your stomach is upset after taking the tablet, take it with a meal or after a snack.

How long to take it

Your doctor will check your progress to make sure the medicine is working and will decide how long your treatment should continue.

If you are unsure, talk to your doctor.

If you forget to take it

If it is almost time for your next dose (e.g. within 2 or 3 hours), skip the dose you missed and take your next dose when you are meant to. Otherwise, take the dose as soon as you remember,

and then go back to taking your tablet as you would normally.

Do not take a double dose to make up for the one that you missed.

If you have trouble remembering when to take your medicine, ask your pharmacist for some hints.

If you take too much

Immediately telephone your doctor or Poisons Information Centre (telephone 13 11 26) for advice, or go to Accident and Emergency at your nearest hospital if you think that you or anyone else may have taken too much Letroz. Do this even if there are no signs of discomfort or poisoning. Keep the telephone numbers for these places handy.

While you are taking Letroz

Things you must do

If you become pregnant while taking Letroz, tell your doctor immediately.

You should not take this medicine while you are pregnant.

Follow your doctor's instructions carefully.

If you do not follow your doctor's instructions, your treatment may not help or you may have unwanted side effects.

Be sure to keep all of your doctor's appointments so that your progress can be checked.

Your doctor may want you to have blood tests from time to time to check on your progress and detect any unwanted side effects. Your doctor may also decide to monitor your bone health as this medicine may cause thinning or wasting of your bones (osteoporosis).

If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking Letroz.

Tell any other doctor, dentist or pharmacist who treats you that you are taking Letroz.

Things you must not do

Do not use Letroz to treat any other complaints unless your doctor says you can.

Do not give this medicine to anyone else, even if their symptoms seem to be similar to yours.

Things to be careful of

Be careful driving, operating machinery or doing jobs that require you to be alert while you are taking Letroz until you know how it affects you.

This medicine may cause dizziness or tiredness in some people. If you have any of these symptoms, do not drive or do anything else that could be dangerous.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Letroz

It may have unwanted side effects in some people in addition to its beneficial effects. All medicines have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

Do not be alarmed by this list of possible side effects. You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor immediately or go to Accident and Emergency at your nearest hospital if you notice any of the following:

- signs that blood clots may have formed, such as sudden severe headache, sudden loss of coordination, blurred vision or sudden loss of vision, slurred speech, numbness or tingling in

- an arm or leg, painful swelling in the calves or thighs, chest pain, difficulty breathing, coughing blood, rapid heartbeat, bluish skin discolouration, fainting.
- constant "flu-like" symptoms (chills, fever, sore throat, sores in mouth, swollen glands, tiredness or lack of energy) that could be a sign of blood problems.
- swelling mainly of the face and throat (signs of allergic reaction)
- weakness or paralysis of limbs or face, difficulty speaking (signs of stroke)
- crushing chest pain or sudden arm or leg (foot) pain (signs of a heart attack)
- swelling and redness along a vein which is extremely tender, possibly painful to touch (signs of thrombophlebitis)

The above side effects may be serious. You may need urgent medical attention or hospitalisation.

Tell your doctor straight away if you experience any of the following:

- yellow skin and eyes, nausea, loss of appetite, dark coloured urine (signs of hepatitis)
- rash, red skin, blistering of the lips, eyes or mouth, skin peeling, fever (signs of skin disorder)
- blurred vision (sign of cataract)
- swelling of the feet, ankles or other parts of the body due to fluid build up

Tell your doctor if you notice any of the following side effects and they worry you:

- skin rash, itching or dry skin
- pain in the muscles, joints or bones; joint stiffness, arthritis
- high level of cholesterol
- vaginal spotting or bleeding
- whitish, thick vaginal discharge, vaginal dryness
- headache
- fever

- tiredness, sleepiness, weakness or dizziness
- difficulty sleeping
- numbness or tingling in hands or feet
- mood changes such as anxiety, nervousness, irritability and depression (sad mood)
- drowsiness
- forgetfulness
- blurred vision or eye irritation
- stomach upset, nausea (feeling sick) or vomiting, indigestion, pain in the abdomen
- constipation
- diarrhoea
- dry mouth, sore mouth, mouth ulcers and cold sores
- thirst, change in sense of taste, dry mouth
- dry mucous membranes of the mouth, nose, vagina
- breast pain
- hot flushes
- increased sweating
- appetite changes
- increase or decrease in weight
- hair thinning
- urgent need to urinate (pass water)
- pain or burning sensation when urinating, which may be a sign of an infection
- pain or burning sensation in the hands or wrist (carpal tunnel syndrome)
- fast or irregular heartbeats palpitations, high blood pressure (hypertension)
- thinning of bones (osteoporosis), bone fractures
- cough
- trigger finger, a condition in which your finger or thumb catches in a bent position.

Tell your doctor if you notice anything else that is making you unwell.

Other side effects not listed above may happen in some people.

Some of these can only be found by laboratory testing.

After taking Letroz

Storage

- Keep your tablets in the container until it is time to take them.
- Store the tablets below 25°C. Protect from Moisture.
- Do not store Letroz in the bathroom or any other place that is hot or steamy.
- Do not leave the tablets in the car or on window sills.

Heat and dampness can destroy some medicines. Letroz will keep well if it is cool and dry.

Keep this medicine where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking Letroz or the tablets have passed their use by (expiry date), ask your pharmacist what to do with any that are left over.

Product description

What it looks like

Letroz is yellow, round, biconvex, film-coated tablet plain on both sides, supplied in blister pack and each pack contains 30 tablets.

Ingredients

Letroz contains 2.5mg of Letrozole as the active ingredient.

The tablets also contain:

- Lactose
- Maize starch
- Hypromellose
- Microcrystalline cellulose
- Sodium starch glycollate (Type A)
- Colloidal anhydrous silica
- Magnesium stearate

- Opadry 03B82927 yellow
(Hyromellose, Titanium dioxide, Iron oxide yellow CI 77492, Macrogol 400 & Purified talc)

Letroz does not contain sucrose, gluten, tartrazine or any other azo dyes.

Name and Address of the Sponsor

Juno Pharmaceuticals Pty Ltd
Level 2, 6 Bond Street,
South Yarra,
VIC – 3141

Australian Registration Number

AUST R 203220

Date of preparation

July 2017