

Betahistine Lupin

Active ingredient: betahistine dihydrochloride

Consumer Medicine Information

This leaflet provides important information about using Betahistine Lupin. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using Betahistine Lupin.**

Where to find information in this leaflet:

1. Why am I using Betahistine Lupin?
2. What should I know before I use Betahistine Lupin?
3. What if I am taking other medicines?
4. How do I use Betahistine Lupin?
5. What should I know while using Betahistine Lupin?
6. Are there any side effects?
7. Product details

1. Why am I using Betahistine Lupin?

Betahistine Lupin contains the active ingredient betahistine dihydrochloride.

Betahistine Lupin is used to treat a disorder of the working of your inner ear. This disorder may include one or more of the following symptoms, in one or both ears:

- Ringing in the ears (tinnitus)
- Loss of clear hearing
- Problems with balance (vertigo)

These symptoms may also be associated with nausea, vomiting and headache. Often these

symptoms together are referred to as Ménière's Syndrome.

Betahistine Lupin tablets contain the active ingredient betahistine dihydrochloride, which work by improving the blood flow of the inner ear and restoring it to normal. It also acts on the nerve endings in the inner ear to normalize the way in which the nerves respond to outside influences.

Your doctor may have prescribed Betahistine Lupin for another reason. Ask your doctor if you have any questions about why Betahistine Lupin has been prescribed for you.

There is no evidence that Betahistine Lupin is addictive.

What should I know before I use Betahistine Lupin?

Warnings

Do not use Betahistine Lupin if:

- you are pregnant or intend to become pregnant.
Betahistine Lupin may affect your developing baby if taken during pregnancy.