

Tamsil 250 mg Tablets

Terbinafine (as hydrochloride)

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about TAMSIL 250 mg tablets. It does not contain information about other forms of terbinafine hydrochloride that are available without a prescription from your pharmacy.

It does not contain all the available information about TAMSIL. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking TAMSIL against the benefits they expect it will provide.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What TAMSIL tablets are used for

Tamsil tablets are used to treat:

- Fungal infections of the finger nails and toe nails
- Tinea (ringworm) infections of the groin and body
- Tinea infections of the feet, commonly called “athlete’s foot.”

These infections are caused by a group of fungi called dermatophytes.

Terbinafine, the active ingredient in Tamsil tablets, works by killing the dermatophytes.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

Your doctor may have prescribed it for another reason.

Tamsil tablets are only available with a doctor’s prescription. This medicine is not addictive.

There is not enough information to recommend the use of this medicine in children.

Before you take TAMSIL tablets

When you must not take it

Do not take TAMSIL if you have ever had an allergic reaction to:

- Terbinafine, the active ingredient, or to any of the other ingredients listed at the end of this leaflet
- Any other medicines, foods, preservatives or dyes

Your doctor will want to know if you are prone to allergies.

Some of the symptoms of an allergic reaction may include:

- Shortness of breath
- Wheezing or difficulty breathing
- Swelling of the face, lips, tongue or other parts of the body
- Rash, itching or hives on the skin.

Do not take TAMSIL tablets:

- If you have any problems with your kidneys
- If you have or ever had a problem with your liver.

This medicine is not recommended if you currently have a liver problem because it may make the problem worse. If you had a liver problem in the past and your liver is functioning normally now, your doctor may prescribe Tamsil tablets but may want to check your liver function before and during treatment with this medicine. Your doctor might take blood tests to monitor your liver function. In case of abnormal test results, he or she may ask you to stop taking Tamsil.

Do not take TAMSIL tablets after the expiry date printed on the pack, or if the packaging is torn or shows signs of tampering.

In that case, return it to your pharmacist.

Before you start to take it

Tell your doctor if you:

1. Are pregnant or intend to become pregnant

There is no experience with use of TAMSIL tablets during pregnancy. If your doctor thinks it is necessary for you to take it, he/she will discuss with you the benefits and risks involved.

2. Are breast-feeding

Breastfeeding is not recommended since terbinafine, the active ingredient in TAMSIL tablets, passes into breast milk. There is a possibility that your baby could be affected.

3. Have any skin problems such as rash, red skin, blistering of the lips, eyes or mouth, skin peeling, fever (possible signs of serious skin reactions), rash due to high level of a specific type

of white blood cells
(eosinophilia)

4. Have any blood disorders
or experience weakness, unusual
bleeding, bruising or frequent
infections
5. Have or experience thickened
patches of red/silver skin
(psoriasis)
or facial rash, joint pain, muscle
disorder, fever (cutaneous and
systemic lupus erythematosus)

**If you are not sure whether you
should start taking TAMSIL
tablets, talk to your doctor or
pharmacist.**

Taking other medicines

**Tell your doctor if you are taking
any other medicines, including
medicines that you buy without a
prescription from your
pharmacy, supermarket or health
food shop.**

Some medicines and TAMSIL
tablets may interfere with each
other. These include:

- some medicines used to treat
depression and other mental
disorders, including obsessive-
compulsive disorders and panic
attacks (e.g. some
antidepressants such as tricyclic
antidepressants selective
serotonin reuptake inhibitors
including class 1A, 1B AND
1C, monoamine oxidase
inhibitors type B, desipramine)
- Some medicines for parkinson's
disease
- Some medicines used to treat an
irregular heartbeat, heart
problems, high blood pressure
and migraines (e.g. metoprolol)
- Some medicines used to treat
stomach ulcers (e.g. cimetidine)
- Some medicines called
antibiotics used to treat
infectious diseases (e.g.
rifampicin)
- Caffeine
- Cyclosporin, a medicine used to
help prevent organ transplant
rejection or to treat certain

problems with the immune
system

- Oral contraceptives (birth
control pills). You may have
problems, such as bleeding
between periods, while you are
taking TAMSIL tablets.
- Warfarin, a medicine used to
prevent blood clots

You may need to take different
amounts of your medicines or you
may need to take different
medicines.

Your doctor and pharmacist have
more information.

**If you have not told your doctor
about any of these things, tell
him/her before you start taking
this medicine.**

How to take TAMSIL tablets

**Follow all directions given to you
by your doctor and pharmacist
carefully.**

These directions may differ from
the information contained in this
leaflet.

If you do not understand the
instructions on the label, ask your
doctor or pharmacist for help.

How much to take

**Follow your doctor's instructions
on how many TAMSIL tablets to
take.**

The usual dose of TAMSIL is one
tablet (250 mg) each day. If you
have kidney problems, the dose
may be reduced to one-half a tablet
each day.

How to take it

**Take the TAMSIL tablet with a
full glass of water. If you find that
the TAMSIL tablet upsets your
stomach, try taking it
immediately after a light meal.**

**Take the tablet at about the same
time each day.**

Taking your tablet at the same time
each day will have the best effect. It
will also help you remember when
to take it.

How long to take it

The length of your treatment will
depend on the type of infection you
have, what part of the body is
affected and how well you respond
to treatment.

Fungal skin infections (tinea):

If you have a tinea infection of the
feet (athlete's foot), you will
usually take the tablets for 2 to 6
weeks.

If you have a tinea infection of the
body or groin, you will usually take
the tablets for 2 to 4 weeks.

The signs and symptoms of
infection may last for several weeks
after the fungi (dermatophytes) have
been killed.

Fungal nail infections:

Fungal nail infections usually take
longer to heal than fungal skin
infections. You will usually take the
tablets for anywhere from 6 weeks
to 3 months. But, if you have a nail
infection of the big toe or your nails
grow very slowly, you may need to
take the tablets for up to 6 months.

It may take several months after
you stop taking TAMSIL tablets for
your nail to look completely
normal. That is because the
deformed part of the nail has to
grow out and be replaced by a
healthy nail.

If you forget to take it

If it is almost time for your next
dose (within 4 hours), skip the dose
you missed and take your next dose
when you are meant to.

**Otherwise, take it as soon as you
remember, and then go back to
taking your tablet(s) as you would
normally.**

Do not take a double dose to make up for the dose that you missed.

If you are not sure what to do, talk to your doctor or pharmacist.

If you have trouble remembering to take your tablets, ask your pharmacist for some hints.

If you take too much (overdose)

Immediately telephone your doctor or Poisons Information Centre (telephone 13 11 26), or go to accident and emergency at your nearest hospital, if you think that you or anyone else may have taken too much TAMSIL. Do this even if there are no signs of discomfort or poisoning. Keep the telephone numbers for these places handy.

Some of the symptoms of an overdose may include headache, nausea (feeling sick), stomach pain and dizziness.

While you are using TAMSIL tablets

Things you must do

Make sure to take your tablet every day and continue taking it until your doctor tells you to stop.

This will ensure that all of the infection is gone and will lessen the chance of the infection coming back once you stop taking this medicine.

Make sure to have any blood tests done that are ordered by your doctor.

Any side effects on your liver, kidneys or blood can be detected by blood tests.

Tell your doctor immediately if you notice any of the following:

- Fever
- Sore throat
- Mouth ulcers
- “flu-like” symptoms (chills, aching joints, swollen glands, lack of energy)

- **Any other signs of infection, apart from the fungal infection you are being treated for**

If you become pregnant while taking TAMSIL tablets, tell your doctor immediately.

Your doctor can discuss with you the risks and benefits of taking it during pregnancy.

Remind your doctor and pharmacist that you are taking TAMSIL tablets if you are about to be started on any new medicine.

Tell any other doctor, dentist or pharmacist who treats you that you are taking TAMSIL tablets.

Things you must not do

Do not give this medicine to anyone else, even if their condition seems similar to yours.

Do not use it to treat any other complaints unless your doctor tells you to.

Things to be careful of

Be careful driving, operating machinery or doing jobs that require you to be alert while you are taking TAMSIL tablets until you know how it affects you.

This medicine can cause tiredness, sleepiness, dizziness or light-headedness in some people. If you have any of these symptoms, do not drive, use machines, or do anything else that could be dangerous.

Be careful to keep the infected areas dry and cool and change clothing that is in direct contact with the infected areas every day.

This will help to clear up the infection and make sure that it does not return.

Side Effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking TAMSIL, even if you do not think it is connected with the medicine.

All medicines can have side effects. Sometimes they are serious, but most of the time they are not. You may need medical treatment if you get some of the side effects.

Do not be alarmed by these lists of possible side effects. You may not experience any of them. Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor immediately or go to Accident and Emergency at your nearest hospital if you notice any of the following:

- Chest pain
- Signs of a severe allergic reaction such as swelling of the face, lips, tongue, throat or other part of the body; shortness of breath, wheezing or troubled breathing; dizziness, redness, itching or rash on the skin; flushing, crampy abdominal pain, loss of consciousness; joint pain, stiffness, rash, fever or swollen/enlarged lymph nodes
- Possible signs of a serious liver problem such as persistent nausea, loss of appetite, unusual tiredness, vomiting, pain in the upper right abdomen, yellowing of the skin and/or eyes, dark urine or pale bowel motions
- Possible signs of a serious skin reaction such as painful red areas, large blisters, peeling of layers of skin, bleeding in the lips, eyes, mouth, nose or genitals. These signs may be accompanied by fever and chills, aching muscles and feeling generally unwell
- Possible signs of a blood problem such as constant “flu-like” symptoms (fever, sore throat, mouth ulcers, chills, swollen glands, lack of energy)
- Possible signs of diseases that affect certain types of blood cells; unusual bleeding or bruising
- Possible signs of a disease that affects the level of red blood cells including abnormal pale

skin, mucosal lining or nail beds, unusual tiredness or weakness or breathlessness on exertion

- Possible signs of blood vessel inflammation: rash, fever, itching, tiredness or if you notice appearance of purplish-red spots under the skin surface
- Possible signs of pancreas inflammation: severe upper stomach pain with radiation to the back
- Possible signs of muscle necrosis: unexplained muscle weakness and pain or dark (red-brown) urine.

The above are serious side effects that need medical attention. Serious side effects are rare.

Tell your doctor if you notice any of the following and they worry you:

- Nausea (feeling sick) or vomiting
- Upset stomach (heartburn, cramps, wind, belching)
- Loss of appetite
- Diarrhoea
- Aching joints or muscles
- Headache
- Light headedness
- Tiredness, sleepiness
- Skin rash due to high level of a specific type of white blood cells
- Loss of or change in sense of taste, which usually returns to normal within several weeks of stopping TAMSIL tablet treatment
- Blurred vision, decreased sharpness of vision
- Other skin problems
- Psoriasis (thickened patches of red skin, often with silvery scales)
- Hair loss
- Tingling or numbness
- Decreased physical sensitivity
- Smell disorders or loss of smell.

- Anxiety (with symptoms such as sleep disturbances, fatigue, loss of energy or diminished ability to think or concentrate) and depressive symptoms (e.g. depressed mood) due to taste disturbances
- Decreased hearing, impaired hearing and/or perception of noises in the absence of sound (e.g. hissing, ringing) in ears.

Tell your doctor if you notice anything else that is making you unwell.

Some people may have other side effects not yet known or mentioned in this leaflet

After using TAMSIL tablets

Storage

Keep your medicines in the blister pack until it is time to take it.

Store it in a cool, dry place below **25°C and protected from light**

Do not store TAMSIL or any other medicine in the bathroom or near a sink

Do not leave it in the car or on window sills.

Keep the medicine where children cannot reach it.

A locked cupboard at least one-and-a-half meters above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking TAMSIL or the tablets have passed their expiry date, ask your pharmacist what to do with any that are left over.

Product Description

What it looks like

White to off white round tablet embossed “R250” on one side and

score (bisect) line on the other side; pack of 42.

Ingredients

The active ingredient in TAMSIL is terbinafine (as hydrochloride). Each tablet contains 250 mg of terbinafine.

Tamsil tablets also contain:

- Microcrystalline cellulose
- Colloidal anhydrous Silica
- Hypromellose
- Magnesium stearate
- Croscarmellose sodium

If you are concerned about which tablet you are taking or any of the above ingredients, talk to your doctor or pharmacist before starting to take TAMSIL.

Name & Address of Sponsor

Tamsil is supplied in Australia by:

Arrotex Pharmaceuticals Pty Ltd
15 – 17 Chapel Street,
Cremorne, VIC 3121
www.arrotex.com.au

TAMSIL 250 mg Tablets AUST R
167577

This leaflet was prepared in July
2022